

EFFECT OF ALKALINITY ON FECUNDITY OF *CALIGUS MINIMUS* PARASITIZING ON SEABASS (*Lates calcarifer*) IN LABORATORY CONDITION

KHOA DUY TRAN NGUYEN^{1,2*}, FAIZAH SHAROM HARISON³, SUHAIRI MAZELAN³
AND SABRI MUDA³

¹College of Aquaculture and Fisheries Campus II, Can Tho University, 3/2 Street, Ninh Kieu, Can Tho City, Vietnam

²Faculty of Fisheries Shimoarata 4-50-20, Kagoshima University, Kagoshima Shimoarata 890 8580 Japan

³School of Fisheries and Aquaculture Science, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia

*Corresponding author email: tndkhoa@ctu.edu.vn

Abstract: Caligid copepod is one of the most common parasites in marine fish culture, causing negative effects on growth and survival of its host. This study aims to assess the fecundity of *Caligus minimus* parasitizing on seabass (*Lates calcarifer*) and evaluate the parasite's abundance dynamics under various alkalinity levels. Wild *C. minimus* were recorded with 60.9 ± 14.3 eggs in its egg string, which varies in length (3.38 ± 0.27 mm), and egg sizes ranging from 0.34 mm (width) to 0.08 mm (length). Wild *C. minimus* were observed to attach and survive on its host for 50 to 71 days, producing one to three egg strings in its life. Under 60 ppm and 70 ppm of alkalinity condition, the abundance and hatching index of *Caligus* reached a peak of $734 \pm 14.5\%$ and $919 \pm 25.5\%$ of the initial infection. There were no records of hatching at 10, 20, 110 and 120 ppm of alkalinity. Within 21 days, no significant difference in the number of egg and egg size under various alkalinity levels were observed. These results demonstrated that alkalinity strongly affects the population of caligid parasites.

KEYWORDS: Fecundity, *Caligus minimus*, *Lates calcarifer*, alkalinity

Introduction

Sea lice (copepods) have been reported as one of the most critical problems in marine fish culture, causing severe negative effects on stock survival, growth and susceptibility to disease (Johnson *et al.*, 2004; Skovgaard, 2005; Maran *et al.*, 2009). The ensuing losses from fish deaths as a direct or indirect consequence of parasitic copepod infestation may be huge (Torrissen *et al.*, 2013; Lafferty *et al.*, 2015). Sea lice are also known to rapidly reproduce and survive in large populations (Tully *et al.*, 1993), which may affect wild fish populations (Krkošek *et al.*, 2005; Costello, 2009).

Copepod infestation, especially by *Caligus minimus*, is common in southeast Asian seabass (*Lates calcarifer*) and has also been reported in its Mediterranean cousin (*Dicentrarchus labrax*), causing losses to fish farmers there (Khoa *et al.*, 2018; Noor El-Deen *et al.*, 2013). The parasites will damage the host's skin and epidermal layer, leading to stress, osmotic problems and secondary infections (Johnson *et al.*, 2004; Er & Kayış, 2015; Saraiva *et al.*, 2015;

Khoa *et al.*, 2018). Shinn *et al.* (2015) reported that 80% of seabass farms worldwide have been affected by this parasite, which resulted in a 30 to 50% loss of stock. However, the impact of parasitism was limited only to economic and environmental aspects (Heuch *et al.*, 2005).

In Malaysian fish farms, the infestation is also a prevalent problem. Among the reported species of parasites and their hosts are *C. chistos* in *Lutjanus johni*, *C. longipedis* in *Gnathonodon speciosus*; *C. rotundigenitalis* in *L. erythropterus*; *C. epidemicus* in *L. calcarifer* and *Epinephelus coioides*; and, *C. punctatus* and *C. minimus* in *L. calcarifer* (Maran *et al.*, 2009; Leaw *et al.*, 2012; Muhd-Faizul *et al.*, 2012; Khoa *et al.*, 2018).

The impact of environmental factors on parasite population dynamics are rarely documented. Previous studies had observed the reproduction and life cycle of copepods (Lin & Ho, 1993; Piasecki & MacKinnon, 1995; Heuch *et al.*, 2000; González & Carvajal, 2003; Khoa *et al.*, 2018) or created a model population based on fecundity (Tully & Whelan, 1993; Bravo *et*

al., 2009). A good understanding of how the environment affects parasite population may come in handy in predicting outbreaks and managing the disease they cause.

Tully & Whelan (1993) suggested that the fecundity of copepods was decisively associated with water temperature. The body size of female lice had been observed to increase at low temperatures, but egg development took longer and had low hatching index (Nordhagen & Schram, 2000; Heuch *et al.*, 2000). The parasites found on wild fish were larger and had longer egg strings compared with those on farm fish (Tully & Whelan, 1993; Nordhagen & Schram, 2000). Nevertheless, the direct effects of environmental factors were not clear.

Being Crustaceans, caligid copepods require significantly high levels of alkalinity in their environment to harden their new exoskeletons after moulting (Middlemiss *et al.*, 2016). The alkalinity level and Ca²⁺ in water had been shown to play an important role in post-moulting calcification of crustacean exoskeletons, which affects growth performance and survival rate of these organisms (Cameron, 1989; Truchot & Fogue, 1998; Middlemiss *et al.*, 2016). Interruptions to the post-moulting process significantly could significantly reduced the population and productivity of crustaceans (Cowthorn, 1997; Borisov *et al.*, 2007; Middlemiss *et al.*, 2016). However, most studies were conducted on large and economically-important crustaceans, such as lobsters (Middlemiss *et al.*, 2016), crabs (Cameron, 1989) and shrimps (Furtado *et al.*, 2014), and none was ever conducted on copepod parasites.

Therefore, this study aims to evaluate the effects of alkalinity on the fecundity and population dynamics of *C. minimus* parasites found in *L. calcarifer*.

Materials and Methods

Sample collection and fecundity study

Sampling was undertaken every four months throughout the year (June, October, February) with 150 infected seabass sampled. *C. minimus* infested *L. calcarifer* were collected

from fish farms in Bukit Tambun, Penang, Malaysia (5°16'01.83N and 100°27'26.74E) and transferred to the Parasitology laboratory at University Malaysia Terengganu in Kuala Terengganu, Terengganu, Malaysia.

The parasites were removed from the host and observed under a microscope to identify them according to the methods by Boxshall & Defaye (1995). A total of 100 *C. minimus* ovigerous females were isolated and reared in separate beakers containing 50 ml of seawater to collect the copepodids they produced. The copepodids were then introduced to seabass fingerlings in small aquariums until the females matured and produced egg strings.

The new copepodids' lifespan, egg string length, egg size and number of eggs were measured and calculated according to Heuch *et al.* (2000) and Schram & Heuch (2001). The total length of egg strings was averaged from 100 randomly-selected egg strings. For the size of a single egg, the length of 10 egg sections in each egg string were measured under the microscope and averaged. The number of eggs was calculated by dividing the total length of egg strings with the length of the egg sections as shown in Equation 1 (Heuch *et al.* 2000).

$$\text{Number of eggs per female} = \frac{\text{Total length of egg strings}}{\text{length of single egg}} \quad (1)$$

Hatching index at various alkalinity levels

The experimental laboratory set-up was described in Figure 1. Ovigerous *C. minimus* females with mature eggs were placed in three 50 ml beakers. Each beaker was stocked with 10 parasites. Before stocking, the eggs were counted to determine the percentage of viable and non-viable eggs. The latter appeared dark when illuminated from below, were irregularly packed or merged into one long section in the egg string as described by Heuch *et al.* (2000). The hatching index, number of nauplius and copepodids in each treatment were presented in the average percentage of 30 ovigerous females as in Equation 2.

$$\text{Hatching index} = \frac{(\text{Number of eggs introduced initially} - \text{Number of unhatched eggs}) \times 100}{\text{Number of eggs introduced initially}} \quad (2)$$




Figure 1: Diagram of the experimental set

Parasite survivability at various alkalinity levels **Statistical analysis**

A total of 10 ovigerous *C. minimus* parasites with egg strings were introduced to five seabass fingerling hosts weighing 10 ± 2 g in 15 L transparent aquariums. The aquariums were maintained in triplicates at room temperature, with 25 ppt of salinity and low aeration for 21 days. There were 12 levels of alkalinity from 10 to 120 ppm. Each level was applied for 48 hours and NaHCO_3 was used to increase alkalinity while maintaining the pH at 7.8. The fish were fed daily with a controlled amount of commercial pellets and the parasites' egg strings were observed.

Water was changed every three days via a drained pipe installed with a plankton filter (25 μm). At the end of the experiment, all the hosts were killed and the *parasites* collected and examined under a dissecting microscope.

Data were analyzed using IBM SPSS software Version 24.0 (IBM, Armonk, New York, USA) and Origin 2018 (OriginLab, Northampton, Massachusetts, USA). One-way ANOVA was used to determine the significant variations ($p < 0.05$) between fecundity and fluctuation of *Caligus* presence at different alkalinity levels. Significant differences in treatments were further subjected to Duncan's post-hoc test.

Results and Discussion
Fecundity under different alkalinity levels

Throughout the experiment, the water used to transport the fish and parasites was kept between 25.5 and 31 °C, at a pH of 7.8, 25 to 30.5 ppt of salinity and 50 to 70 ppm of alkalinity.

The length of egg strings from 100 females was observed to be 3.38 ± 0.27 mm, and each

brood was carrying 60.9 ± 14.3 eggs in its string. The egg string length was shortened after matured eggs were released. Matured eggs normally ruptured from the distal sac. However, the hatching caused the breaking of egg strings at un-sequenced parts (Figure 2).

Compared with other studies in Table 1, the egg string length and number of eggs in

C. minimus were higher than *C. rogercresseyi*, *Lepeophtheirus pectoralis* and other freshwater copepods. However, the egg size of this parasite was smaller. This reproductive characteristics gave *C. minimus* the ability to quickly multiply and cause damage in a fish population within a short time.

Table 1: Fecundity of *Caligus minimus* and some crustacean parasites

Species	Egg string length (mm)	Number of egg/female	Egg size (mm)	Host	Source
<i>Caligus rogercresseyi</i>	-	45 ± 16	-	<i>Salmo salar</i>	Bravo <i>et al.</i> (2009)
<i>Caligus rogercresseyi</i>	1.5-3.1	12-56	-	<i>Eleginops maclovinus</i>	González <i>et al.</i> (2012)
<i>Caligus rogercresseyi</i>	4.9-6.1	19-95	-	<i>Oncorhynchus mykiss</i>	González <i>et al.</i> (2012)
<i>Caligus rogercresseyi</i>	5.07	59 ± 19	-	<i>Salmo salar</i>	Bravo <i>et al.</i> (2013)
<i>Lepeophtheirus salmonis</i>	-	1000	-	-	Costello (1993)
<i>Calanidae</i> sp	-	40	0.152	-	Poulin (1995)
<i>Pseudodiaptomidae</i> sp	-	23	0.075	-	Poulin (1995)
<i>Diaptomidae</i> sp	-	29	0.086	-	Poulin (1995)
<i>Porcellidiidae</i> sp	-	11	0.085	-	Poulin (1995)
<i>Lepeophtheirus pectoralis</i>	3.36	45.5 ± 18.7	0.077	<i>Platichthys flesu</i>	Frade <i>et al.</i> , (2015)
<i>Lepeophtheirus salmonis</i>	8-40	50-500	-	<i>Salmo trutta</i>	Tully & Whelan (1993)

The ovigerous *C. minimus* survived up to three or four days without a host under laboratory conditions. Adult *C. minimus* females produced one to three egg strings in its life. The egg size was 0.34 mm wide and 0.08 mm long. The copepod lifecycle began as nauplius larvae, which metamorphosized into copepodids,

chalimi, pre-adult and adults through molting. The parasites started to attach to the host from the copepodid stage. Through observation of 100 specimens, the lifespan of *Caligus minimus* was between 50 and 71 days on their host (61.5 ± 10.5 days) (Figure 2).




Figure 2: *C. minimus* could survive with a fish host/or without host by attaching on temporary substrate

Based on initial egg string length and egg size of *Caligus*, the number of eggs and percentage of viable and non-viable eggs were not significantly different between aquariums. The alkalinity levels did not significantly affect egg size and number.

The interaction between parasites and host in an ecosystem had been studied to formulate strategies to control infestations. Several studies were conducted on *C. rogercresseyi* (Bravo *et al.*, 2009; Gonzalez *et al.*, 2012; Sandra Bravo *et al.*, 2013), *L. simplex* (Serna *et al.*, 2015), *L. salmonis* (Tully & Whelan, 1993; Ugelvik *et al.*, 2017) and *L. pectoralis* (Frade *et al.*, 2015). However, these studies were conducted in temperate zones and there were very few studies in tropical waters.

Heuch *et al.* (2000) noted that a higher water temperature positively affected the copepods' egg production and development, whereby greater fecundity and shorter life cycle were

observed in *L. salmonis*. When cultured at 8.7 °C, the parasites' egg strings were longer by seven per cent than their counterparts at 12.2 °C. However, it took approximately four times longer for *L. salmonis* to produce an egg string at 7.2 °C than 12.2 °C (Heuch *et al.*, 2000).

This study showed a great fecundity of *C. minimus* in warm waters. The ovigerous female carried 50 to 70 eggs on a pair of egg strings, similar to *C. rogercresseyi* found parasitizing on *S. salar* (Sandra Bravo *et al.*, 2013). But that figure was far lower compared to *L. salmonis* and *L. pectoralis*, which had up to 500 to 1000 eggs in each brood.

The fecundity difference could be influenced by many environmental factors, such as temperature (Sarvala, 1979; Tully, 1989; Heuch *et al.*, 2000), female body size (Tully & Whelan, 1993), type of host (farmed or wild type) (Tully & Whelan, 1993; Nordhagen & Schram, 2000) and salinity level (Muhd-Faizul *et al.*, 2012).

Therefore, the significant difference in water temperature between temperate and tropical zones could lead to greater egg production and metamorphosis of copepods in warm conditions.

The life cycle of *C. rogercresseyi* could be prolonged from 31 to 45 days in waters ranging from 10.3 to 12.8 °C. *C. elongates* could survive for 43.3 days at 10 °C. In warmer temperatures, *C. pageti* could only live for 10 to 11 days at 26 °C (Russell, 1925), while the lifespan of *C. minimus* was even shorter at five to seven days at 29.5 °C (Khoa, 2015; Khoa *et al.*, 2018). Additionally, *L. latis* took just 222 hours to complete its life cycle at 30 °C (Brazenor & Hutson, 2013; Qamarina *et al.*, 2014). This explained the challenges posed by crustacean parasites in tropical waters due to rapid multiplication.

Hatching index (%)

A higher alkalinity level could negatively affect the hatching index of copepods. The highest rate of hatching was observed between 60 ppm (81.99%) and 70 ppm (85.27%) of alkalinity levels (Figure 3), which was significantly different compared to other levels of treatment. The eggs were observed to take up water through osmosis, enlarging their internal membranes, but no hatching was recorded at 10, 20, 110 and 120 ppm of alkalinity. The result also showed a limited hatching of eggs at 30 ppm and 100 ppm of alkalinity. Plot analysis suggested that hatching index could peak at 65 ppm, and the approximate range of alkalinity for *C. minimus* to hatch was from 45 ppm to 80 ppm, where the hatching index was above 50%.




Figure 3: Hatching index of *Caligus minimus* egg in various alkalinity levels

The presence of *C. minimus* strongly fluctuated at different alkalinity levels (Figure 4). The highest infection was recorded at 60 and 70 ppm ($734 \pm 14.5\%$ and $919 \pm 25.5\%$ of the initial infection, respectively), and were significantly different compared to other levels. The high mortality rate of *C. minimus* in 10, 20, 30, 110, 120 ppm of alkalinity caused a

reduction in parasite population, and none of the organism was observed on the fish at the end of the experiment. At between 40 and 100 ppm, the parasite population slightly increased while a higher population dynamics was observed in 50, 80 and 90 ppm of alkalinity. However, there were no significant differences observed among these treatments.




Figure 4: Population dynamic of *Caligus minimus* in various alkalinity levels

In the marine ecosystem, pH is an important factor that affects the viability of marine life (Doney *et al.*, 2009; Omstedt *et al.*, 2010), and is characterized by total alkalinity (Omstedt *et al.*, 2010). The rise in water alkalinity could negatively affect copepod populations (Garzke *et al.*, 2016) and this imbalance would lead to disorder in the organisms’ metabolism (Pörtner *et al.*, 2004; Michaelidis *et al.*, 2005). Metabolic depression had been suggested to affect copepod fecundity (Mayor *et al.*, 2007). In egg development and hatching, a decrease in pH and alkalinity would interfere in egg membrane depolarization during fertilization (McCulloh *et al.*, 1987), depress respiration and protein synthesis in gametes and embryos (Spikes, 1983), and cause genetic abnormalities in egg cells (Pagano *et al.*, 1985; Stack *et al.*, 2006). Hence, egg mortality dramatically affects the hatching index and abundance of copepod parasites.

Female copepods would suffer from low egg production (Vehmaa *et al.*, 2012; Zervoudaki *et al.*, 2014) and produce a high number of unfertilized, non-viable or viable eggs in quiescent states (Cripps *et al.*, 2014). Mass

mortality of parasites had also been observed (Pedersen *et al.*, 2013) as well as impairment of the crustacean’s post-moulting stage (Borisov *et al.*, 2007; Michaelidis *et al.*, 2005). This study was limited by its short duration (21 days) and it could not evaluate the effect of alkalinity on egg number and size.

Conclusion

The current study evaluated the fecundity of *C. minimus* and the effect of alkalinity on the parasite’s fertility. The optimal alkalinity levels for copepods to thrive on seabass hosts was between 60 and 70 ppm. However, further studies need to investigate the population dynamics of *C. minimus* under various environmental factors to model epidemiological progress and develop effective management strategies for commercial aquaculture.

Acknowledgements

This study was funded by the Exploratory Research Grant Scheme (ERGS, Vote number 55081), Ministry of Education, Malaysia. We

would like to thank the Institute of Tropical Aquaculture (AKUATROP), Universiti Malaysia Terengganu, and the National Fish Health Research Center (NAFISH) for their support.

References

- Borisov, R. R., Epelbaum, A. B., Kryakhova, N. V., Tertitskaya, A. G., & Kovatcheva, N. P. (2007). Cannibalistic behavior in red king crabs reared under artificial conditions. *Russian Journal of Marine Biology*, 33(4), 227–231.
- Boxshall, G. A., D. D. (1995). *Pathogens of wild and farmed fish: sea lice. Aquaculture* (Vol. 133).
- Bravo, S., Erranz, F., & Lagos, C. (2009). A comparison of sea lice, *Caligus rogercresseyi*, fecundity in four areas in southern Chile. *Journal of Fish Diseases*, 32(1), 107–113.
- Bravo, S., Pozo, V., Silva, M. T., & Abarca, D. (2013). Comparison of the fecundity rate of *Caligus rogercresseyi* infesting Atlantic salmon (*Salmo salar* L.) on farms in two regions of Chile. *Aquaculture*, 404–405, 55–58.
- Brazenor, A. K., & Hutson, K. S. (2013). Effect of temperature and salinity on egg hatching and description of the life cycle of *Lernanthropus latis* (Copepoda: Lernanthropidae) infecting barramundi, *Latescalcarifer*. *Parasitology International*, 62(5), 437–447.
- Cameron, B. Y. J. N. (1989). Post-Moult Calcification in the Blue Crab, *Callinectes Sapidus*: Timing and Mechanism. *The Journal of Experimental Biology*, 143, 285–304.
- Cawthorn, R. J. (1997). Overview of “bumper car” disease—Impact on the North American lobster fishery. *International Journal for Parasitology*, 27, 167–172.
- Costello, M. J. (2009). How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *Proceedings of the Royal Society B: Biological Sciences*, 276(1672), 3385–3394.
- Cripps, G., Lindeque, P., & Flynn, K. (2014). Parental exposure to elevated pCO₂ influences the reproductive success of copepods. *J. Plankton Res.*, 36(5), 1165–1174.
- Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). Ocean Acidification: The Other CO₂ Problem. *Annual Review of Marine Science*, 1(1), 169–192.
- Er, A., & Kayış, Ş. (2015). Intensity and prevalence of some crustacean fish parasites in Turkey and their molecular identification. *Turkish Journal of Zoology*, 39(6), 1142–1150.
- Frade, D. G., Santos, M. J., & Cavaleiro, F. I. (2015). The reproductive effort of *Lepeophtheirus pectoralis* (Copepoda: Caligidae): insights into the egg production strategy of parasitic copepods. *Parasitology*, 143(1), 87–96.
- Francisco, N. M. S., Rivas-Salas, A. I., Gomez, S., & Fajer-Avila, E. J. (2015). Developmental stages and fecundity of *Lepeophtheirus simplex* (Copepoda: Caligidae) parasitic on bullseye puffer fish (*Sphoeroides annulatus*). *Folia Parasitologica*, 62(004).
- Furtado, P. S., Poersch, L. H., & Wasielesky, W. (2014). The effect of different alkalinity levels on *Litopenaeus vannamei* reared with biofloc technology (BFT). *Aquaculture International*, 23(1), 345–358.
- Garzke, J., Hansen, T., Ismar, S. M. H., & Sommer, U. (2016). Combined effects of ocean warming and acidification on copepod abundance, body size and fatty acid content. *Plos One*, 11(5), e0155952.
- González, L., & Carvajal, J. (2003). Life cycle

- of *Caligus rogercresseyi*, (Copepoda: Caligidae) parasite of Chilean reared salmonids. *Aquaculture*, 220(1–4), 101–117.
- González, M. T., Molinet, C., Arenas, B., Asencio, G., & Carvajal, J. (2012). Fecundity of the sea louse *Caligus rogercresseyi* on its native host *Eleginops maclovinus* captured near salmon farms in southern Chile. *Aquaculture Research*, 43(6), 853–860.
- Heuch, P. A., Bjørn, P. A., Finstad, B., Holst, J. C., Asplin, L., & Nilsen, F. (2005). A review of the Norwegian “National Action Plan Against Salmon Lice on Salmonids”: The effect on wild salmonids. *Aquaculture*, 246(1–4), 79–92.
- Heuch, P. A., Nordhagen, J. R., & Schram, T. A. (2000). Egg production in the salmon louse [*Lepeophtheirus salmonis* (Kroyer)] in relation to origin and water temperature. *Aquaculture Research*, 31(11), 805–814.
- Johnson, S. C., Treasurer, J. W., Bravo, S., Nagasawa, K., & Kabata, Z. (2004). A review of the impact of parasitic copepods on marine aquaculture. *Zoological Studies*, 43(2), 229–243.
- Khoa, T. N. . (2015). *Some Aspects of Caligus spp. Infestation On Cultured Asian Seabass (Lates calcarifer)*. Universiti Malaysia Terengganu, Malaysia.
- Khoa T. N. D; Suhairi M; Sabri M; Faizah Harison. (2018). The life cycle of *Caligus minimus* on Seabass (*Lates calcarifer*) from floating cage culture. *Thalassas: An International Journal of Marine Sciences, In press*, 1–9.
- Krkošek, M., Lewis, M. A., & Volpe, J. P. (2005). Transmission dynamics of parasitic sea lice from farm to wild salmon. *Proceedings of the Royal Society B: Biological Sciences*, 272(1564), 689–696.
- Lafferty, K. D., Harvell, C. D., Conrad, J. M., Friedman, C. S., Kent, M. L., Kuris, A. M., ... Saksida, S. M. (2015). Infectious Diseases Affect Marine Fisheries and Aquaculture Economics. *Annual Review of Marine Science*, 7(1), 471–496.
- Leaw, Y. Y., Faizah, S., Anil, C., & Kua, B. C. (2012). Prevalence, mean intensity and site preference of *Caligus rotundigenitalis* Y, 1933 (Copepoda: Caligidae) on cage cultured crimson snapper (*Lutjanus erythropterus* Bloch, 1790) from Bukit Tambun, Penang, Malaysia. *Veterinary Parasitology*, 187(3–4), 505–510.
- Lin, C. L., & Ho, J. S. (1993). Life history of *Caligus epidemicus* Hewitt parasitic on tilapia (*Oreochromis mossambicus*) cultured in brackish water.
- Maran, B. a V., Seng, L. T., Ohtsuka, S., Nagasawa, K., & Nagasawa, K. (2009). Records of *Caligus* (Crustacea: Copepoda: Caligidae) from Marine Fish Cultured in Floating Cages in Malaysia with a Redescription of the Male of *Caligus longipedis* Bassett-Smith, 1898. *Zoological Studies*, 48(6), 797–807.
- Mayor, D. J., Matthews, C., Cook, K., Zuur, A. F., & Hay, S. (2007). CO₂-induced acidification affects hatching success in *Calanus finmarchicus*. *Marine Ecology Progress Series*, 350, 91–97.
- McCulloh, D. H., Lynn, J. W., & Chambers, E. L. (1987). Membrane depolarization facilitates sperm entry, large fertilization cone formation, and prolonged current responses in sea urchin oocytes. *Developmental Biology*, 124(1), 177–190.
- Michaelidis, B., Ouzounis, C., Palaras, A., & Portner, H. O. (2005). Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, 293 (Lackner 2003), 109–118.
- Middlemiss, K. L., Urbina, M. A., & Wilson, R. W. (2016). Effects of seawater alkalinity on calcium and acid-base regulation in juvenile

- European lobster (*Homarus gammarus*) during a moult cycle. *Comparative Biochemistry and Physiology -Part A : Molecular and Integrative Physiology*, 193, 22–28.
- Muhd-Faizul, H. A. H., Kua, B. C., & Leaw, Y. Y. (2012). Caligidae infestation in Asian seabass, *Lates calcarifer*, Bloch 1790 cultured at different salinity in Malaysia. *Veterinary Parasitology*, 184(1), 68–72.
- Noor El-Deen, a. I. E., Abeer, E. M., & Azza, H. M. H. (2013). Field studies of caligus parasitic infections among cultured seabass (*Dicentrarchus labrax*) and mullet (*Mugil cephalus*) in marine fish farms with emphasis on treatment trials. *Global Veterinaria*, 11(5), 511–520.
- Nordhagen, J. R., & Schram, T. A. (2000). Size as indicator of origin of salmon lice *Lepeophtheirus salmonis* (Copepoda: Caligidae). *Contributions to Zoology*, 69(1), 99–108.
- Omstedt, A., Edman, M., Anderson, L. G., & Laudon, H. (2010). Factors influencing the acid-base (pH) balance in the Baltic Sea: A sensitivity analysis. *Tellus, Series B: Chemical and Physical Meteorology*, 62(4), 280–295.
- Pagano, G., Cipollaro, M., Corsale, G., Esposito, A., Ragucci, E., & Giordano, G. G. (1985). pH-Induced Changes in Mitotic and Developmental Patterns in Sea-Urchin Embryogenesis .1. Exposure of Embryos. *Teratogenesis Carcinogenesis and Mutagenesis*, 5(2), 113–121.
- Pedersen, S. A., Hansen, B. H., Altin, D., & Olsen, A. J. (2013). Medium-term exposure of the North Atlantic copepod *Calanus finmarchicus* (Gunnerus, 1770) to CO₂-acidified seawater: Effects on survival and development. *Biogeosciences*, 10(11), 7481–7491.
- Piasecki, W., & MacKinnon, B. M. (1995). Life cycle of a sea louse, *Caligus elongatus* von Nordmann, 1832 (Copepoda, Siphonostomatoida, Caligidae). *Canadian Journal of Zoology*, 73(1), 74–82.
- Pörtner, H. O., Langenbuch, M., & Reipschläger, A. (2004). Biological impact of elevated ocean CO₂ concentrations: lessons from animal physiology and earth history. *Journal of Oceanography*, 60, 705–718.
- Poulin, R. (1995). Clutch size and egg size in free living and parasitic copepods a comparative analysis. *Evolution*, 49(2), 325–336.
- Qamarina, N., Khalid, A., & Shaharoum-Harrison, F. (2014). The life cycle of the parasitic crustacean, *Lernanthropus latis Yamaguti*, 1954 (Copepoda : Lernanthropidae), on marine-cultured fish, *Lates calcarifer*, from Setiu Wetland, Terengganu. *Journal of Parasitology Research*, 2014(2014), 1–6.
- Russell, F. S. (1925). A new species of *Caligus* from Egypt, *Caligus pageti*, sp. n. *Journal of Natural History Series* 9, 15(90), 611–618.
- Saraiva, A., Costa, J., Serrão, J., Eiras, J. C., & Cruz, C. (2015). Study of the gill health status of farmed sea bass (*Dicentrarchus labrax* L., 1758) using different tools. *Aquaculture*, 441, 16–20.
- Sarvala, J. (1979). Effect of temperature on the duration of egg, nauplius and copepodite development of some freshwater benthic Copepoda. *Freshwater Biology*, 9, 515–534.
- Schram, T. A., & Heuch, P. A. (2001). The egg string attachment mechanism of selected pennellid copepods. *Journal of the Marine Biological Association of the United Kingdom*, 81(1), 23–32.
- Shinn, A. P., Pratoomyot, J., Bron, J. E., Paladini, G., Brooker, E. E., & Brooker, A. J. (2015). Economic costs of protistan and metazoan parasites to global mariculture. *Parasitology*, 142(1), 196–270.

- Skovgaard, A. (2005). Infection with the dinoflagellate parasite *Blastodinium* spp. in two Mediterranean copepods. *Aquatic Microbial Ecology*, 38(1), 93–101.
- Spikes, J. D. (1983). Metabolism of Sea Urchin Sperm. *The American Naturalist*, 258(9), 5392–5399.
- Stack, C., Lucero, A. J., & Shuster, C. B. (2006). Calcium-responsive contractility during fertilization in sea urchin eggs. *Developmental Dynamics*, 235(4), 1042–1052.
- Torrissen, O., Jones, S., Asche, F., Guttormsen, A., Skilbrei, O. T., Nilsen, F., Jackson, D. (2013). Salmon lice - impact on wild salmonids and salmon aquaculture. *Journal of Fish Diseases*, 36(3), 171–194.
- Truchot, J. P., & Fogue, J. (1998). Effect of water alkalinity on gill CO₂ exchange and internal PCO₂ in aquatic animals. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 119(1), 131–136.
- Tully, O. (1989). The succession of generations and growth of the caligid copepods *Caligus elongatus* and *Lepeophtheirus salmonis* parasitising farmed Atlantic salmon smolts (*Salmo salar* L.). *Journal of the Marine Biological Association of the United Kingdom*, 69(2), 279–287.
- Tully, O., Poole, W. R., & Whelan, K. F. (1993). Parameters and possible causes of epizootics of *Lepeophtheirus salmonis* (Kroyer) infesting sea trout (*Salmo trutta* L.) off the west coast of Ireland. In D. D. Geoffrey A. Boxshall (Ed.), *Pathogens Of Wild And Farmed Fish: Sea Lice* (p. 378). Ellis Horwood Limited.
- Tully, O., & Whelan, K. F. (1993). Production of nauplii of *Lepeophtheirus salmonis* (Kroyer) (Copepoda: Caligidae) from farmed and wild salmon and its relation to the infestation of wild sea trout (*Salmo trutta* L.) off the west coast of Ireland in 1991. *Fisheries Research*, 17(1–2), 187–200.
- Ugelvik, M. S., Skorping, A., & Mennerat, A. (2017). Parasite fecundity decreases with increasing parasite load in the salmon louse *Lepeophtheirus salmonis* infecting Atlantic salmon *Salmo salar*. *Journal of Fish Diseases*, 40(5), 671–678.
- Vehmaa, A., Brutemark, A., & Engström-Öst, J. (2012). Maternal effects may act as an adaptation mechanism for copepods facing pH and temperature Changes. *PLoS ONE*, 7(10), 1–8.
- Zervoudaki, S., Frangoulis, C., Giannoudi, L., & Krasakopoulou, E. (2014). Effects of low pH and raised temperature on egg production, hatching and metabolic rates of a Mediterranean copepod species (*Acartia clausi*) under oligotrophic conditions. *Mediterranean Marine Science*, 15(1), 74–83.