

## TEXTURE AND COLOUR CHANGES IN *Penaeus vannamei* AFTER COATINGS WITH *Piper betle* (L) LEAVES EXTRACT

AL HAFIZ ZAINI<sup>1,2</sup>, KHAIRUL NAIM AZIZ<sup>3</sup> AND NURUL ULFAH KARIM<sup>1\*</sup>

<sup>1</sup>Higher Institution Centre of Excellence, Institute of Tropical Aquaculture and Fisheries, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia. <sup>2</sup>Greater Asia Agro Berhad, Plo 545 and 678, Jalan Keluli 7, Kawasan Perindustrian Pasir Gudang, 81700 Pasir Gudang, Johor, Malaysia. <sup>3</sup>Marine Research Station (MARES), Faculty of Applied Science, Universiti Teknologi MARA, Perlis Branch, 02600 Arau, Perlis, Malaysia.

\*Corresponding author: [ulfah@umt.edu.my](mailto:ulfah@umt.edu.my)  
Submitted final draft: 21 August 2023

<http://doi.org/10.46754/jssm.2024.01.001>  
Accepted: 10 September 2023 Published: 15 January 2024

**Abstract:** *Penaeus vannamei* is an important aquaculture product heavily exported to countries such as Europe and China due to its high consumer demand. However, commercialisation of the product involves various levels of cold chain and time to reach consumers which can lead to rapid deterioration. The study aims to evaluate the texture and colour of *P. vannamei* after being coated with 0.5, 1.0 and 2.0% betel leaf extract for 16 days of chilled storage. *P. vannamei* was found to have slightly random insignificant changes in texture. However, the colour changes were slowly decreased especially in *P. vannamei* coated with 2.0% betel leaves extract and supported by sensory evaluation for consumer acceptance. *P. vannamei* coated with 2.0% betel leaves extract resulted in a significantly lower ( $p < 0.05$ ) hardness ( $20646.11 \pm 480.10$  N) compared to others on the initial day and showed a slightly decreasing trend during storage but not significantly ( $p > 0.05$ ). After 12 days of chilled storage, *P. vannamei* coated with 2.0% betel leaf extract showed a significant increase ( $p < 0.05$ ) in springiness from  $0.53 \pm 0.15\%$  at the initial day to  $0.63 \pm 0.06\%$  at the end of storage day, indicating a firmer texture. *P. vannamei* coated with 2.0% betel leaves extract had the highest cohesiveness ( $0.35 \pm 0.05\%$ ) and resilience ( $0.58 \pm 0.06$ ) value on the initial day but not significantly different ( $p > 0.05$ ). However, over time, cohesiveness and resilience values showed a decreasing trend, indicating that the shrimp became less cohesive and resilient with prolonged storage. The colour of the *P. vannamei* coated with 2.0% betel leaf extract recorded a lightness  $L^*$  value of  $29.23 \pm 0.14$ ; redness  $a^*$  value of  $0.37 \pm 0.25$ ; and blueness  $b^*$  value of  $5.12 \pm 0.37$  and significantly different ( $p < 0.05$ ) compared to *P. vannamei* coated with 0.5%, and 1.0% betel leaves extract and 1.25% SMS throughout 16 days of chilled storage. The organoleptic test showed that *P. vannamei* coated with 2.0% betel leaf extract was highly accepted by consumers. Therefore, coating *P. vannamei* with a 2.0% concentration of betel leaf extract promises a significantly improved quality of *P. vannamei* at prolonged storage.

Keywords: Whiteleg shrimp, betel leaves, quality, sensory analysis.

### Introduction

*Penaeus vannamei* or commonly known as whiteleg shrimp is an important food source for humans, as it is rich in nutrients and contains polyunsaturated fatty acids. However, *P. vannamei* is easily perishable and has a limited shelf-life which depends on its pre-treatment, conditions, and duration of storage (Sae-Leaw & Benjakul, 2019). The poor appearance of *P. vannamei* can reduce its market value and lead to consumer rejection, resulting in economic loss (Gonçalves & de Oliveira, 2016). Edible coating is a promising and emerging technique for maintaining the quality of *P. vannamei*,

especially in terms of texture, colour, and shelf-life extension.

A previous study conducted by Bao (2014) clearly demonstrated that the extract derived from straw mushrooms is a positive source of natural antioxidants, which can be utilized to prevent melanosis in shrimp during iced storage. Meanwhile, the extraction of moringa leaves has also demonstrated its antioxidant properties and its ability to slow down the melanosis process in shrimp (Karim *et al.*, 2018). Edible coating provides various benefits such as delaying lipid oxidation, preserving protein functionality,

reducing off-odours, discolouration, and delaying spoilage (Khaledian *et al.*, 2021). Additionally, it can prolong the product's freshness by delaying bacteria growth, reducing moisture loss, and acting as an antioxidant and antibacterial agent (Kumari & Thakur, 2020). Edible coating substitutes the natural layers at the product surface to prevent solute movements out of the shrimp, gas aroma and moisture losses while allowing for a controlled exchange of important gases such as oxygen, carbon dioxide, and ethylene, which are involved in food respiration (Dehghani *et al.*, 2018).

In order to control discolouration and lipid oxidation in fish meat and other crustaceans, various studies related to the use of antioxidants have been conducted and widely used in the market (Encarnacion *et al.*, 2012). Antioxidants or inhibitors of oxidation are compounds which retard or prevent oxidation and generally prolong the life of the oxidisable matter (Chirag *et al.*, 2013). Betel leaf is one of the natural herbs that comes from the betel plant, an evergreen climber that is a good source of antioxidants and antimicrobials (Sarma *et al.*, 2018). The use of natural additives is expected to replace synthetic materials such as sodium metabisulphite (SMS), which can have many negative effects on consumers (Naureen *et al.*, 2021). SMS is the cheapest, easiest, and most effective method available today to preserve shrimp. It is also considered an antimicrobial food additive (Omar, 1998).

Therefore, the objective of this study is to determine the effectiveness of betel leaf extract on the texture and colour changes of *P. vannamei* during 16 days of chilled storage (4°C) and to evaluate consumer acceptance of the product. This bio-preservation technique may be introduced as a new natural alternative to maintain the quality and safety of *P. vannamei*.

## Materials and Methods

### *Piper Betle Sample Collection and Extract Preparation*

Approximately  $2.0 \pm 0.1$  kg of matured *Piper betle* leaves freshly collected from Pagoh, Johor

were used in the experiment. The mature leaves are commonly larger, darker green in colour, have a more distinct flavour and aroma, and usually with an estimated age of 6 months. The betel leaves were dried in the shade at room temperature for 7 days, with an average temperature of  $29 \pm 0.1^\circ\text{C}$ . The leaves were pulverised into powder by using a Waring blender (Waring Commercial, Malaysia). Sample extractions were performed according to the method described by Ibrahim *et al.* (2022), with some modifications. Approximately  $30 \pm 0.1$  g of betel powder was mixed with 200 ml solvent (6:4) methanol-distilled water. The mixture was shaken at room temperature for 24 hours using a Stuart Orbital Shaker (Fisher Scientific, Malaysia), followed by filtration with Whatman filter paper number 1. The mixture was dried using a rotary evaporator R-210 (Buchi, Switzerland) to remove excess solvent. The extract was kept in a container at room temperature prior to further analysis.

### *Penaeus vannamei Preparation*

Fresh *P. vannamei* of approximately  $15 \pm 0.5$  g was immersed at different concentrations of 0.5, 1.0 and 2.0% betel leaves extract and 1.25% SMS treatment for 20 minutes, respectively. The controls were left untreated. All samples were stored at chilled storage at 4°C for 16 days. All analysis was carried out in triplicate at intervals of 4 days during 16 days of storage.

### *Texture Analysis of Penaeus vannamei*

Texture profile analysis of *P. vannamei* including hardness, springiness, cohesiveness, and resilience was done by using TA.XT Plus Texture Analyser (Stable Micro System, United Kingdom). The instrument was set up to a pre-test speed of  $1.0 \text{ mm s}^{-1}$ , test speed of  $0.10 \text{ mm s}^{-1}$ , post-test speed of  $1.0 \text{ mm s}^{-1}$ , and trigger force of 5.0 g. The samples were compressed to 50% of their height using an aluminium cylinder probe of 4 mm diameter and the test speed was at  $1 \text{ mm s}^{-1}$ .

**Colour Analysis of *Penaeus vannamei***

A CR-400 colourimeter (Konica Minolta, Japan) was used to determine the colour changes in *P. vannamei* throughout 16 days of chilled storage. The readings were recorded in a CIE L\*, a\*, and b\* colour space, where L\* represents the brightness from black (0) to white (100), a\* represents from green to red, and b\* represents from blue colour to yellow. The instrument is calibrated using a white standard plate. The readings were taken in triplicate at intervals of 4 days during 16 days of chilled storage.

**Sensory Analysis of *Penaeus vannamei***

Sensory analysis was performed by the method of Balfour *et al.* (2014) with some modifications. This sensory test was included to measure consumer acceptance of obvious changes in the shrimp by experienced panellists. A total of 20 panellists were involved in this experiment with experience in the field of shrimp processing between 2 to 10 years at Greater Asia Agro Berhad. The panellists performed the sensory evaluations at daily intervals during the 16 days of chilled storage. The observed criteria involve colour, texture, and odour (Table 1).

**Statistical Analysis**

All analysis was performed in triplicate ( $n = 3$ ) and the data were represented as mean  $\pm$  standard deviation. The collected data was tabulated and analyzed using IBM SPSS Statistic Version 20 software (IBM Corporation, New York). The data was subjected to two-way ANOVA with Ryan, Einot, Gabriel, Welsh Range Q (REGWQ) post hoc test. In all analyses by ANOVA, the residuals were inspected for normality and homogeneity of variance across the treatments. Different superscripts (A, B, C, D, E) were used to indicate significant differences ( $p < 0.05$ ) between storage days. Meanwhile, different superscripts (a, b, c) were used to indicate significant differences ( $p < 0.05$ ) between treatments.

**Results and Discussion**

**Texture Analysis of *Penaeus vannamei***

*P. vannamei* coated with 1.25% SMS was recorded as the lowest hardness value compared to others on the initial day. Meanwhile, the controls showed the highest hardness value compared to others but were not significantly ( $p > 0.05$ )

Table 1: Grading description scheme on the quality of whole shrimp Balfour *et al.* (2014)

		Criteria		
		Colour	Texture	Odour
Grading	Excellent	Body has dark red to bright pink	Firm, body has a cleaned defined shape	Strong seaweedy, marine odour
	Good	Body has a natural light pink to white	The flesh has slight firmness; the body still has a defined shape	Weak characteristic shrimp odour
	Moderate	Naturally light pink with grey-greenish or yellowish discolouration	Flesh soft to the touch, may easily be pushed by a finger	Marine odours are diminishing, weak fishy odours, even slight ammonia
	Borderline	Naturally pink with grey-greenish or yellowish discolouration	Flesh may be easily pushed by a finger	Weak ammonia odour
	Unfit	Naturally pink with grey-greenish or yellowish discolouration. Lack defined colour	Flesh soft and pliable	Ammonia odour

different. Hardness value showed a fluctuating trend throughout the 16 days of storage in most samples, except for *P. vannamei* coated with 1.0% betel leaves extract, where it was uniformly degrading throughout the storage days. However, the hardness results of all samples showed no significant difference ( $p > 0.05$ ) to one another during the 16 days of storage.

The changes in *P. vannamei* hardness are due to the proteolysis in the flesh, which is caused by endogenous and microbial enzymes (Jeyakumari *et al.*, 2019) (Table 2). The effect observed on the *P. vannamei* was minimal based on the results. A higher hardness value of *P. vannamei* indicates a low quality, good quality *P. vannamei* meat should have a firm texture and

be consistent in appearance and should not be mushy or overly soft (Kim *et al.*, 2020).

On the initial day, *P. vannamei* coated with 2.0% betel leaf extract showed a lower springiness value compared to *P. vannamei* coated with 0.5% betel leaf extract (Table 3). Interestingly, *P. vannamei* coated with 2.0% betel leaf extract showed an increasing trend from  $0.53 \pm 0.15\%$  at 0 day to  $0.63 \pm 0.06\%$  at 12<sup>th</sup> days of storage. Meanwhile, *P. vannamei* coated with 1.25% SMS showed a lower value compared to the control, with no significant difference ( $P > 0.05$ ) on the 16 days of the experiment. The springiness of the *P. vannamei* provides a sensation of comfort when chewed and provides satisfaction for its connoisseurs

Table 2: Hardness of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	25578.78 ± 4420.80 <sup>Aa</sup>	22646.10 ± 3396.52 <sup>Aa</sup>	22080.97 ± 1233.26 <sup>Aa</sup>	20646.11 ± 480.10 <sup>Ab</sup>	20093.15 ± 1235.35 <sup>Aa</sup>
4	23045.16 ± 938.38 <sup>Aa</sup>	19394.65 ± 852.80 <sup>Aa</sup>	21750.53 ± 1920.50 <sup>Aa</sup>	17795.19 ± 2168.88 <sup>Ba</sup>	23369.41 ± 1302.94 <sup>Aa</sup>
8	9398.54 ± 2207.53 <sup>Aa</sup>	21214.23 ± 2853.49 <sup>Ab</sup>	20605.47 ± 502.44 <sup>Ab</sup>	20397.18 ± 109.47 <sup>Ab</sup>	22060.89 ± 3615.23 <sup>Bb</sup>
12	23063.41 ± 1735.53 <sup>Aa</sup>	20477.28 ± 4189.14 <sup>Aa</sup>	19016.90 ± 1135.33 <sup>Aa</sup>	18776.97 ± 1039.44 <sup>ABa</sup>	17947.75 ± 2482.63 <sup>Aa</sup>
16	19263.99 ± 5077.74 <sup>Aa</sup>	21097.99 ± 5832.10 <sup>Aa</sup>	18632.53 ± 7457.90 <sup>Aa</sup>	17013.86 ± 5111.05 <sup>Aa</sup>	22067.91 ± 598.73 <sup>Aa</sup>

Note: Capital letters indicate significant differences between different storage days for the same treatment and small letters indicate significantly different treatments for the same storage day.

Table 3: Springiness of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	0.72 ± 0.16 <sup>Aab</sup>	0.93 ± 0.03 <sup>Ab</sup>	0.81 ± 0.20 <sup>Aab</sup>	0.53 ± 0.15 <sup>Aa</sup>	0.64 ± 0.20 <sup>ABab</sup>
4	0.74 ± 0.19 <sup>Aa</sup>	0.88 ± 0.10 <sup>Aa</sup>	0.85 ± 0.05 <sup>Aa</sup>	0.70 ± 0.12 <sup>ABa</sup>	0.87 ± 0.02 <sup>Ba</sup>
8	0.64 ± 0.25 <sup>Ba</sup>	0.66 ± 0.27 <sup>Aab</sup>	0.65 ± 0.06 <sup>Ab</sup>	0.65 ± 0.21 <sup>Bab</sup>	0.70 ± 0.14 <sup>ABab</sup>
12	0.65 ± 0.22 <sup>Aab</sup>	0.70 ± 0.15 <sup>Aab</sup>	0.68 ± 0.05 <sup>Aab</sup>	0.63 ± 0.06 <sup>Bb</sup>	0.54 ± 0.30 <sup>Aa</sup>
16	0.77 ± 0.15 <sup>Aa</sup>	0.77 ± 0.25 <sup>Aa</sup>	0.72 ± 0.24 <sup>Aa</sup>	0.64 ± 0.23 <sup>ABa</sup>	0.74 ± 0.12 <sup>ABa</sup>

Note: Capital letters indicate significant differences between different storage days at the same treatment and small letters indicate significantly different treatments at the same storage day.

(Salasiah & Jirarat, 2018). If the texture of the *P. vannamei* is too soft or easily crumbled, the food can taste unpleasant and has no satisfying taste. On the other hand, if the texture of the *P. vannamei* is too hard, the food can be difficult to chew and uncomfortable in the mouth.

The cohesiveness value of *P. vannamei* coated with 2.0% betel leaf extract showed the highest value but was not significantly different ( $p < 0.05$ ) compared to other treatments in the initial days. Coatings with 0.5 and 1.0% betel leaf extract showed no significant effects ( $p > 0.05$ ) on the cohesiveness value of *P. vannamei* during 16 days of storage (Table 4). Meanwhile, the cohesiveness value of *P. vannamei* coated with 1.25% SMS showed a significant increase ( $p < 0.05$ ) value after being stored for 12 days. Cohesiveness texture refers to the ability of *P. vannamei* texture to maintain its integrity and not fall apart when bitten, as well as provide a special taste and texture sensation when chewed.

The consumer indicates that *P. vannamei* is fresh and highly acceptable to consume. The higher the cohesiveness coefficient, the better the *P. vannamei* texture is to maintain the integrity and roundness of the texture during chewing or processing in the mouth.

The resilience value was significantly higher ( $p < 0.05$ ) in *P. vannamei* coated with 2.0% betel leaf extract compared to other treatments on the initial day (Table 5). However, the value showed a significant decreasing trend ( $p < 0.05$ ) with prolonged storage. In contrast, *P. vannamei* coated with 1.25% SMS showed an increasing resilience value with prolonged storage, up to 12 days. This indicates that *P. vannamei* meat underwent some internal structural changes during storage. Some studies state that temperature might have the potential to affect the myofibrillar protein to bring about immediate to intense contractions of muscle of the *P. vannamei* meat (Okpala, 2015).

Table 4: Cohesiveness of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	0.24 ± 0.04 <sup>Aa</sup>	0.24 ± 0.01 <sup>Aa</sup>	0.25 ± 0.04 <sup>Aa</sup>	0.35 ± 0.05 <sup>Ab</sup>	0.21 ± 0.07 <sup>Aa</sup>
4	0.19 ± 0.02 <sup>Aa</sup>	0.23 ± 0.01 <sup>Aa</sup>	0.23 ± 0.04 <sup>Aa</sup>	0.25 ± 0.04 <sup>Ba</sup>	0.26 ± 0.00 <sup>ABa</sup>
8	0.23 ± 0.04 <sup>B</sup>	0.25 ± 0.03 <sup>Aab</sup>	0.26 ± 0.02 <sup>Ab</sup>	0.27 ± 0.02 <sup>Bab</sup>	0.28 ± 0.06 <sup>ABab</sup>
12	0.24 ± 0.05 <sup>ABab</sup>	0.24 ± 0.06 <sup>Aab</sup>	0.24 ± 0.03 <sup>Aab</sup>	0.25 ± 0.04 <sup>Ba</sup>	0.32 ± 0.10 <sup>Bb</sup>
16	0.24 ± 0.04 <sup>ABa</sup>	0.25 ± 0.05 <sup>Aa</sup>	0.28 ± 0.07 <sup>Aa</sup>	0.30 ± 0.09 <sup>ABa</sup>	0.27 ± 0.02 <sup>ABa</sup>

Note: Capital letters indicate significant differences between different storage days at the same treatment and small letters indicate significantly different treatments at the same storage day.

Table 5: Resilience of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	0.38 ± 0.10 <sup>Aa</sup>	0.39 ± 0.05 <sup>Aa</sup>	0.42 ± 0.04 <sup>Aa</sup>	0.58 ± 0.06 <sup>Aa</sup>	0.40 ± 0.11 <sup>Aa</sup>
4	0.33 ± 0.01 <sup>Aa</sup>	0.38 ± 0.02 <sup>Aa</sup>	0.41 ± 0.05 <sup>Aa</sup>	0.44 ± 0.06 <sup>Ba</sup>	0.47 ± 0.01 <sup>ABa</sup>
8	0.56 ± 0.05 <sup>Ba</sup>	0.42 ± 0.02 <sup>Aa</sup>	0.43 ± 0.02 <sup>Aa</sup>	0.49 ± 0.03 <sup>Ba</sup>	0.48 ± 0.08 <sup>ABa</sup>
12	0.42 ± 0.11 <sup>Aa</sup>	0.40 ± 0.04 <sup>Aa</sup>	0.43 ± 0.02 <sup>Aa</sup>	0.46 ± 0.03 <sup>Ba</sup>	0.53 ± 0.13 <sup>Ba</sup>
16	0.45 ± 0.06 <sup>ABa</sup>	0.46 ± 0.09 <sup>Aa</sup>	0.49 ± 0.10 <sup>Aa</sup>	0.53 ± 0.11 <sup>ABa</sup>	0.46 ± 0.05 <sup>ABa</sup>

Note: Capital letters indicate significant differences between different storage days at the same treatment and small letters indicate significantly different treatments at the same storage day.

Resilience in *P. vannamei* texture is important in food processing because it can affect the final texture of the product. A more resilient *P. vannamei* can be more easily to cut and process and can maintain its shape during the processing process. *P. vannamei* that has rotted or is not fresh generally has low resilience, as the meat has softened and cannot return to its original shape after being stressed.

### Colour Analysis of *Penaeus vannamei*

*P. vannamei* coated with 2.0% betel leaf extract showed a significantly lower ( $p < 0.05$ )  $L^*$  value throughout 16 days of storage compared to the controls. A similar result was found with regard to the *P. vannamei* coated with 0.5% betel leaf extract. However, the  $L^*$  value of *P. vannamei* coated with a 1.25% SMS showed a significant ( $p < 0.05$ ) higher  $L^*$  value compared to others at 16 days of storage (Table 6). The colour of *P. vannamei* was observed from light pink to deep

red. A higher  $L^*$  value indicates a lighter shade of pink while a lower  $L^*$  value indicates a darker shade of pink or red.

Initially,  $a^*$  value of control was significantly lower ( $p < 0.05$ ) compared *P. vannamei* coated with 2.0% betel leaf extract. However, after being stored for 16 days in chilled storage, the  $a^*$  value of control showed a significantly higher ( $p < 0.05$ ) value compared to the 0 day. In contrast,  $a^*$  value of *P. vannamei* coated with 2.0% betel leaf extract showed a significant decrease ( $p < 0.05$ ) value after 16 days of storage. The least  $a^*$  value was found in *P. vannamei* coated with 1.25% SMS at the end of the storage day (Table 7). A positive  $a^*$  value indicates that the colour of *P. vannamei* has more red tones and a negative value indicates that the colour has more green tones.

The  $b^*$  value of control at 0 day of storage showed a significantly lower ( $p < 0.05$ ) value compared to *P. vannamei* coated with SMS.

Table 6: Lightness of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	36.15 ± 0.60 <sup>aAB</sup>	37.63 ± 0.29 <sup>aA</sup>	36.93 ± 0.55 <sup>aAB</sup>	35.95 ± 0.12 <sup>aB</sup>	36.94 ± 0.59 <sup>aAB</sup>
4	37.85 ± 3.14 <sup>ba</sup>	37.35 ± 0.79 <sup>aA</sup>	38.24 ± 1.51 <sup>aA</sup>	31.08 ± 0.65 <sup>bB</sup>	29.10 ± 0.45 <sup>cb</sup>
8	30.24 ± 0.32 <sup>ca</sup>	29.81 ± 0.38 <sup>ba</sup>	32.28 ± 2.22 <sup>bb</sup>	24.50 ± 1.57 <sup>dc</sup>	34.33 ± 1.49 <sup>bd</sup>
12	34.37 ± 0.73 <sup>da</sup>	31.31 ± 0.15 <sup>bb</sup>	26.43 ± 0.37 <sup>dc</sup>	30.56 ± 0.58 <sup>bcB</sup>	33.37 ± 0.39 <sup>ba</sup>
16	32.01 ± 0.12 <sup>ca</sup>	31.30 ± 0.42 <sup>baB</sup>	29.90 ± 0.09 <sup>dbC</sup>	29.23 ± 0.14 <sup>cc</sup>	34.98 ± 0.44 <sup>bd</sup>

Note: Capital letters indicate significant differences between different storage days at the same treatment and small letters indicate significantly different treatments at the same storage day.

Table 7:  $a^*$  value of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	0.16 ± 0.08 <sup>aA</sup>	0.93 ± 0.15 <sup>aB</sup>	0.44 ± 0.25 <sup>aA</sup>	0.90 ± 0.02 <sup>aBC</sup>	0.47 ± 0.15 <sup>baC</sup>
4	2.22 ± 0.78 <sup>ba</sup>	2.01 ± 0.39 <sup>baB</sup>	0.52 ± 0.16 <sup>ac</sup>	0.12 ± 0.25 <sup>bc</sup>	1.58 ± 0.05 <sup>ab</sup>
8	1.24 ± 0.34 <sup>caB</sup>	0.54 ± 0.15 <sup>aBC</sup>	1.49 ± 0.40 <sup>ba</sup>	1.603 ± 0.17 <sup>ca</sup>	0.85 ± 0.04 <sup>bcB</sup>
12	0.71 ± 0.16 <sup>daC</sup>	2.63 ± 0.27 <sup>cb</sup>	0.49 ± 0.10 <sup>aA</sup>	1.12 ± 0.25 <sup>ac</sup>	0.40 ± 0.05 <sup>ba</sup>
16	0.59 ± 0.09 <sup>ca</sup>	1.50 ± 0.19 <sup>db</sup>	0.79 ± 0.17 <sup>ac</sup>	0.37 ± 0.25 <sup>bcd</sup>	0.11 ± 0.51 <sup>cd</sup>

Note: Capital letters indicate significant differences between different storage days at the same treatment and small letters indicate significantly different treatments at the same storage day.



However, the  $b^*$  value showed no significant difference ( $p > 0.05$ ) in *P. vannamei* coated with 2.0% betel leaf extract. The  $b^*$  value showed a significantly higher ( $p < 0.05$ ) value in *P. vannamei* coated with 2.0% betel leaves extract at day 16<sup>th</sup> of storage compared to 0 day of storage. A similar result was found in regards to *P. vannamei* coated with 1.25% SMS (Table 8). A positive  $b^*$  value indicates that the colour of the *P. vannamei* has more yellow tones and a negative value indicates that the colour has more blue tones.

In general, the colour of the *P. vannamei* tends to turn dark due to melanosis, this is due to the polyphenol oxidase (PPO) enzyme biochemical reaction. PPO is certainly not harmful to humans. PPO enzyme oxidizes phenols (colourless) to quinones (coloured), to form melanin (Illera *et al.*, 2019). In addition, based on other studies, the higher the concentration of betel leaves, the darker the colour will be;  $a^*$  and  $b^*$  values also increased after the application of the betel leaves extract (Hoppy *et al.*, 2018). The colour changes were observed where the  $L^*$  value of *P. vannamei* decreased in value but remained positive, indicating that the colour of *P. vannamei* is becoming darker while still retaining its basic hue.

Additionally, the  $a^*$  value indicates that the colour of the *P. vannamei* is becoming redder until day 8 but showed a decreased value at the end of the experiment. The  $b^*$  value indicates that the colour of the *P. vannamei* is

becoming more yellow until day 8 but becomes less yellowish at the end of the experiment. Furthermore, decomposed shrimp will release volatile amine compounds during storage, causing the anthocyanin structure to change to its basic form, resulting in colour change (from pink to greenish colour) (Suratsawadee *et al.*, 2022) as observed at the end of the experiment.

**Sensory Analysis**

The quality of the *P. vannamei* coated with 0, 0.5, 1.0 and 2.0% betel leaves extract and 1.25% SMS showed that the quality of *P. vannamei* declined from the initial day to day 16 of storage. The characteristics used to describe the *P. vannamei* are based on grading criteria for the sensory test set in Table 1. On the fourth day of storage, *P. vannamei* coated with 2.0% betel leaf extract and 1.25% SMS showed good colour compared to other treatments, which showed natural light pink to white (Figure 1). The other treatments showed a moderate colour that was natural light pink with a grey-greenish hue. On the last day of the experiment, *P. vannamei* coated with 2.0% betel leaves extract, 1.0% betel leaves extract and 1.25% SMS showed natural light pink with yellowish discolouration. Meanwhile, other treatments showed a lack of defined colour. During storage, the colour of the shrimp deteriorated, indicating the occurrence of lipid and phenol oxidation (Li *et al.*, 2020).

*P. vannamei* texture was not significantly ( $p > 0.05$ ) changed during the first 4 days of storage, where the body had a defined shape and was

Table 8:  $b^*$  value of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage

Storage Time (days)	Control	0.50% Betel Leaves Extract	1% Betel Leaves Extract	2% Betel Leaves Extract	1.25% SMS
0	0.45 ± 0.14 <sup>4A</sup>	1.19 ± 0.06 <sup>4B</sup>	0.09 ± 0.10 <sup>4A</sup>	0.15 ± 0.04 <sup>4A</sup>	0.56 ± 0.18 <sup>4C</sup>
4	5.45 ± 0.50 <sup>4A</sup>	3.07 ± 0.60 <sup>4B</sup>	1.55 ± 0.49 <sup>4C</sup>	1.07 ± 0.32 <sup>4C</sup>	3.68 ± 0.47 <sup>4D</sup>
8	3.47 ± 0.25 <sup>4A</sup>	3.15 ± 0.09 <sup>4A</sup>	5.24 ± 0.24 <sup>4B</sup>	4.35 ± 0.54 <sup>4C</sup>	4.69 ± 0.17 <sup>4C</sup>
12	3.76 ± 0.38 <sup>4A</sup>	7.55 ± 0.40 <sup>4B</sup>	3.91 ± 0.08 <sup>4A</sup>	1.76 ± 0.18 <sup>4C</sup>	0.49 ± 0.08 <sup>4D</sup>
16	1.54 ± 0.16 <sup>4A</sup>	2.3 ± 0.09 <sup>4B</sup>	1.18 ± 0.02 <sup>4A</sup>	5.12 ± 0.37 <sup>4C</sup>	1.56 ± 0.07 <sup>4A</sup>

Note: Capital letters indicate significant differences between different storage days at the same treatment and small letters indicate significantly different treatments at the same storage day.

firm (Figure 2). *P. vannamei* texture maintains a firm texture for up to 4 days of storage after harvesting (Nunak & Schleining, 2011). On the fifth day, the texture of *P. vannamei* showed a slight change, where the flesh had slight firmness but the texture was in a defined shape. *P. vannamei* coated with 2.0% betel leaf extract and 1.25% SMS showed a slower process of quality texture deterioration compared to other treatments. On the last day of the experiment, the texture of *P. vannamei* coated with 2.0% betel leaves extract and 1.25% SMS showed an easy push with a finger but was not pliable, compared to other treatments where the meat was soft and pliable. Chemical and biological reaction in shrimp, even in frozen temperatures, including oxidative changes of lipids and protein in the muscle causes the quality of shrimp texture to deteriorate (Valencia-Perez et al., 2015).

Spoilage *P. vannamei* emits unpleasant odours. On the second day, *P. vannamei* odour's was diminished, with a weak fishy odour, and a slight ammonia smell could be detected in all treatment samples (Figure 3). However, *P. vannamei* coated with 2.0% betel leaf extract showed the slowest odour deterioration process compared to other treatments. On the last day of storage, *P. vannamei* coated with 2.0% and 1.0% betel leaves extract and 1.25% SMS emitted a slight odour of ammonia compared to other treatments. The emission is a strong ammonia odour. Bacterial contamination and enzymatic activity on shrimp during the storage process cause protein, fat and carbohydrates to decompose into ammonia, hydrogen sulphite, ethyl mercaptan, aldehydes, aldehydes acids, alcohol, ketones, and carboxylic acid gas cause of unpleasant odour (Du et al., 2015).




Figure 1: Colour quality of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage based on sensory test




Figure 2: Texture quality of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage based on sensory test






Figure 3: Odour quality of *P. vannamei* coated at different concentrations of betel leaf extract and 1.25% SMS in 16 days of chilled storage based on sensory test

One possible compound in betel leaves that have biological properties in *P. vannamei* is eugenol. Eugenol is a phenolic compound found in betel leaves (Hadzir & Hadzuin, 2012). Eugenol has been shown to have antimicrobial activity against various foodborne pathogens. When applied to *P. vannamei*, it may reduce the bacterial load on the surface of the shrimp, which can improve their shelf life (Marchese *et al.*, 2017). Eugenol can interact with proteins in the *P. vannamei* tissue, causing changes in their physical and chemical properties. For example, it may alter the surface charge or hydrophobicity of the *P. vannamei* proteins, which could affect their solubility, stability, and functionality. Eugenol could also affect the lipid content and composition of shrimp, which may lead to changes in texture, flavour, and aroma. Eugenol also could potentially interact with the pigments in the shrimp, resulting in changes in their colour (Aara *et al.*, 2020).

*P. vannamei* colour, texture, and odour can influence consumer acceptance. Consumers prefer *P. vannamei* with natural and translucent colours ranging from greyish to light pink. Brightly coloured or unnatural-looking *P. vannamei* may be perceived as less desirable. Furthermore, *P. vannamei* texture should be firm and tender and the odour of *P. vannamei* should have a mild and salty smell. The use of 2.0% betel leaf extract gives the best score in prolonging the shelf life of *P. vannamei* comparable to SMS and other treatments until

the 10<sup>th</sup> day of the experiment. While on the 11<sup>th</sup> day of the experiment, all treatments showed that the *P. vannamei* was unacceptable for the consumer.

## Conclusion

The texture attributes of *P. vannamei*, along with body colour and sensory evaluation, provide comprehensive information about *P. vannamei*'s quality. By considering these factors, consumers can assess the freshness, taste, and overall desirability of *P. vannamei* they are consuming. This study suggested that coating *P. vannamei* with 2.0% betel leaf extract can help maintain the quality during chilled storage for up to 16 days. The coating resulted in a decrease in hardness, springiness, cohesiveness, and resilience, which indicates that the texture of the *P. vannamei* became firmer and less cohesive and resilient with prolonged storage. However, *P. vannamei* coated with 2.0% betel leaf extract had significantly different colour values (Lowest L\* value and a\* and highest b\* value) compared to other treatments throughout the storage day. *P. vannamei* were preferred to have a higher L\* value, lower a\* value and lower b\* value within the acceptable range for shrimp's characteristic pinkish or reddish colour. The sensory evaluation showed that the *P. vannamei* coated with 2.0% betel leaf extract was highly accepted by the consumers. Therefore, the results suggested that coating *P. vannamei* with 2.0% betel leaf extract

is a promising approach to maintaining the quality of the shrimp during commercialization and prolonging its shelf-life.

### Acknowledgements

The authors would like to thank the Higher Institution of Excellence (HiCoE) Institute of Tropical Aquaculture and Fisheries, UMT Programme (Vot No 56054) and Greater Asia Agro Berhad for laboratory facilities support.

### Conflict of Interest Statement

The authors declare that they have no conflict of interest.

### References

- Aara, A., Chappidi, V., & Ramadas, M. N. (2020). Antioxidant activity of eugenol in Piper betel leaf extract. *Journal of Family Medicine and Primary Care*, 9(1), 327.
- Balfour, S. T., Badrie, N., Chang Yen, I., & Chatergoon, L. (2014). Microbiological, physical, and sensory quality of marine shrimp (*Peneaus* spp.) sold by vendors in Trinidad, West Indies. *International Food Research Journal*, 21(4), 1279-1288.
- Bao, H. N. D. (2014). Antioxidant activity of hydrophilic extract from straw mushroom and its effect on shrimp melanosis. *Khon Kaen Agriculture Journal*, 42(4), 79-83.
- Chirag, P. J., Tyagi, S., Halligudi, N., Yadav, J., Pathak, S., Singh, S. P., & Kamboj, P. S. (2013). Antioxidant activity of herbal plants; A recent review. *Journal of Drug Discovery and Therapeutics*, 1(8), 01-08.
- Dehghani, S., Hosseini, S. V., & Regenstein, J. M. (2018). Edible films and coatings in seafood preservation: A review. *Food Chemistry*, 240, 505-513.
- Du, L., Chai, C., Guo, M., & Lu, X. (2015). A model for discrimination freshness of shrimp. *Sensing and Bio-sensing Research*, 6, 28-32.
- Encarnacion, A. B., Fagutao, F., Jintasataporn, O., Worawattanamateekul, W., Hirono, I., & Ohshima, T. (2012). Application of ergothioneine-rich extract from an edible mushroom *Flammulina velutipes* for melanosis prevention in shrimp, *Penaeus monodon* and *Litopenaeus vannamei*. *Food Research International*, 45(1), 232-237.
- Gonçalves, A. A., & de Oliveira, A. R. M. (2016). Melanosis in crustaceans: A review. *LWT-Food Science and Technology*, 65, 791-799.
- Hadzir, N., & Hadzuin, N. (2012). *Extraction of eugenol from betel leaves (Piper betle L.) using hydrodistillation and supercritical carbon dioxide technique*. [Thesis Master]. Selangor: Universiti Putra Malaysia. <http://psasir.upm.edu.my/id/eprint/48483/>
- Hoppy, D., Noerdin, A., Irawan, B., & Soufyan, A. (2018). Effect of betel leaf extract gel on color change in the dental enamel. *Journal of Physics*, 1073(3), 032028.
- Ibrahim, A., Che Ibrahim, M. S., Bakar, K., Bakar, J., Ikhwanuddin, M., & Karim, N. U. (2022). Effects of *Annona muricata* extraction in inhibition of polyphenoloxidase and microbiology quality of *Macrobrachium rosenbergii*. *Journal of Food Science and Technology*, 59(3), 859-868.
- Illera, A. E., Sanz, M. T., Beltrán, S., & Benito-Román, O. (2019). Effect of high-pressure carbon dioxide on polyphenoloxidase from *Litopenaeus vannamei*. *LWT-Food Science and Technology*, 109, 359-365.
- Jeyakumari, A., Debbarma, J., Remya, S., Ravishankar, C. N., Zynudheen, A. A., Ninan, G., & Lalitha, K. V. (2019). Quality changes of white shrimp (*Litopenaeus vannamei*) treated with Potassium Sorbate and Sodium Metabisulphite: A comparative study. *International Journal of Current Microbiology and Applied Sciences*, 8(11), 1331-1339.
- Karim, N. U., Siddiq, U. S. A. A., Razak, M. R. M., Zainol, M. K. M., & Abdullah,

- M. I. (2018). Effect of moringa leaves (*Moringa oleifera*) extraction on quality changes and melanosis of giant freshwater prawn (*Macrobrachium rosenbergii*) during chilled storage. *Italian Journal of Food Safety* 2018, 7, 6846.
- Khaledian, S., Basiri, S., & Shekarforoush, S. S. (2021). Shelf-life extension of pacific white shrimp using tragacanth gum-based coatings containing Persian lime peel (*Citrus latifolia*) extract. *LWT-Food Science and Technology*, 141, 110937.
- Kim, S., Jung, E., Hong, D., Lee, S., Lee, Y., Cho, S., & Kim, S. B. (2020). Quality assessment and acceptability of whiteleg shrimp (*Litopenaeus vannamei*) using biochemical parameters. *Fisheries and Aquatic Sciences*, 23(1), 1-10.
- Kumari, S., & Thakur, R. R. (2020). Edible coating- to enhance shelf-life fruits and vegetables. *Food and Scientific Reports*, 1(5), 56-59.
- Li, D. Y., Yuan, Z., Liu, Z. Q., Yu, M. M., Guo, Y., Liu, X. Y., & Zhou, D. Y. (2020). Effect of oxidation and maillard reaction on color deterioration of ready-to-eat shrimps during storage. *LWT-Food Science and Technology*, 131, 109696.
- Marchese, A., Barbieri, R., Coppo, E., Orhan, I. E., Daglia, M., Nabavi, S. F., Izadi, M., & Ajami, M. (2017). Antimicrobial activity of eugenol and essential oils containing eugenol: A mechanistic viewpoint. *Critical Reviews in Microbiology*, 43(6), 668-689.
- Naureen, I., Saleem, A., Nawaz, A., & Javed, M. (2021). Study the harmful effects of sodium metabisulphite on liver, testes, and lipid peroxidation in male mice (*Mus musculus*). *International Journal of Engineering Applied Science and Technology*, 6(5), 30-38.
- Nunak, N., & Schleining, G. (2011). Instrumental textural changes in raw white shrimp during iced storage. *Journal of Aquatic Food Product Technology*, 20(4), 350-360.
- Okpala, C. O. R. (2015). The physicochemical changes of farm-raised Pacific white shrimp (*Litopenaeus vannamei*) as influenced by iced storage. *Food and Nutrition Sciences*, 6(10), 906.
- Omar, M. I. V. (1998). Utilization of sodium metabisulphite for preservation of frozen-thawed shrimp (*Pandalus borealis*). *Fisheries Training Programme. Final Project* 1998.
- Sae-leaw, T., & Benjakul, S. (2019). Prevention of melanosis in crustaceans by plant polyphenols: A review. *Trends in Food Science and Technology*, 85, 1-9.
- Salasiah, M. N., & Jirarat, T. (2018). Effect of food additives on the quality of white shrimp (*Litopenaeus vannamei*). *Food Research*, 2(6), 546-554.
- Sarma, C., Rasane, P., Kaur, S., Singh, J., Singh, J., Gat, Y., & Dhawan, K. (2018). Antioxidant and antimicrobial potential of selected varieties of *Piper betle* L. (Betel leaf). *Anais da Academia Brasileira de Ciências*, 90, 3871-3878.
- Suratsawadee, A., Wangmo, L., Ratvijitvech, T., & Siripinyanond, A. (2022). A spoilage indicator card based on distance-based color change of paper impregnated with acid-base indicator for freshness monitoring of shrimp. *Microchemical Journal*, 175, 107110.
- Valencia-Perez, A. Z., Soto-Valdez, H., Ezquerra-Brauer, J. M., Márquez-Ríos, E., & Torres-Arreola, W. (2015). Quality changes during frozen storage of blue shrimp (*Litopenaeus stylirostris*) with antioxidant,  $\alpha$ -tocopherol, under different conditions. *Food Science and Technology*, 35, 368-374.