

## ANTIOXIDANT ACTIVITY OF TANNIN-RICH EXTRACT FROM WATER LETTUCE (*Pistia stratiotes*) LEAF

SABRI SUDIRMAN<sup>1\*</sup>, LOUSIANA JUNI ARTHA SITUNGKIR<sup>1</sup>, HERPANDI<sup>1</sup>, INDAH WIDIASTUTI<sup>1</sup> AND MIFTAHUL JANNA<sup>2</sup>

<sup>1</sup>Fisheries Product Technology, Faculty of Agriculture, Universitas Sriwijaya, 30862 Indralaya, Ogan Ilir Regency, South Sumatra, Indonesia. <sup>2</sup>Master Program in Agribusiness, Faculty of Agriculture, Universitas Sriwijaya, 30139 Palembang City, South Sumatra, Indonesia.

\*Corresponding author: [sabrisudirman@unsri.ac.id](mailto:sabrisudirman@unsri.ac.id)

<http://doi.org/10.46754/jssm.2024.10.001>

Received: 6 September 2023

Accepted: 23 April 2024

Published: 15 October 2024

**Abstract:** Oxidative stress is a condition in which the free-radical substance is higher than the potential antioxidant in the human body. Therefore, exogenous antioxidants are needed to resolve this problem. Water lettuce (*Pistia stratiotes*) is an aquatic floating plant that contains several bioactive compounds, including tannins and exhibits antioxidant activity. However, since organic solvent extraction results in a crude polyphenol extract, purification is needed to remove the undesirable compounds. This study was aimed at determining the total tannin content and antioxidant activity of water lettuce leaf extract before and after the purification process. The purification was done using a solid-phase extraction method, the total tannin contents were analysed using Folin-Ciocalteu's method, and the antioxidant activity was analysed using 2,2-diphenyl-1-picrylhydrazyl. The results showed that the purification process successfully removed undesirable compounds and increased the total tannin content of the extract from  $8.71 \pm 1.50$  mg TAE/g of the dry sample to  $13.84 \pm 0.28$  mg TAE/g of the dry sample. The antioxidant activities also increased after the purification from  $IC_{50} 135.14 \pm 22.44$   $\mu$ g/mL to  $IC_{50} 78.45 \pm 5.42$   $\mu$ g/mL. According to these results, the tannin from water lettuce extract can be used as a source of natural antioxidants and formulated as an ingredient in food supplements.

Keywords: Antioxidant, *Pistia stratiotes*, polyphenols, purification, tannins.

### Introduction

Oxidative stress is a condition in which the free radical substance is higher than the potential antioxidant in the human body (Vona *et al.*, 2021). Free radicals are normally produced in the body and are usually destroyed by the body's natural antioxidant system. However, this substance can increase more than antioxidants due to external factors such as radiation, smoking, alcohol intake, industrial residue, and antibiotic residue. Free radicals can attack and damage some important molecules such as lipids and DNA, as well as cause some diseases (Rad *et al.*, 2020). Therefore, exogenous antioxidants are needed to resolve this problem. Generally, this antioxidant contains some bioactive compounds such as polyphenols, flavonoids, and tannins (Spiegel *et al.*, 2023). These compounds can be extracted from some plants, including water lettuce (*Pistia stratiotes*) (Herpandi *et al.*, 2021; Sudirman *et al.*, 2022a).

Water lettuce (*P. stratiotes*) is an aquatic floating plant widely found in tropical areas, including Indonesia. A previous study reported that this plant extract contains several bioactive compounds such as phenolics, tannins, and flavonoids (Sudirman *et al.*, 2017; Sudirman *et al.*, 2022a). Tannins are polyphenol compounds widely found in terrestrial and aquatic plants (Tong *et al.*, 2022). Polyphenols are categorised as secondary metabolites and exhibit excellent antioxidant activity (Das *et al.*, 2020). Generally, tannins can be extracted from plants using organic solvents such as ethanol, acetone, and methanol solutions (Sirisangsawang & Phetyim, 2023). Since the extraction using organic solvent resulted in a crude polyphenol extract; therefore, purification was needed to remove the undesirable compounds such as lipids, sugars, and organic acids (Zeka *et al.*, 2017).

A solid-phase extraction (SPE) method is widely used for the purification process due to its rapidity, simplicity, and economy (Dai & Mumper, 2010). Previous studies reported that purification increases the total polyphenol content of *Inga edulis* (Souza *et al.*, 2007) and the flavonoid compound (luteolin) also increases after the purification process (Zhao *et al.*, 2022). Additionally, purified polyphenol compounds exhibit stronger antioxidant activity than crude extract (Wang *et al.*, 2019). The purification process also increases the inhibition of 3-hydroxy-3-methylglutaryl coenzyme-A (HMG-CoA) reductase and the pancreatic lipase activity of polyphenol extract from water lettuce (Sudirman *et al.*, 2022b; Sudirman *et al.*, 2023). According to these conditions, we hypothesised that purification would also increase tannin content and antioxidant activity of the ethanol extract of water lettuce leaf. Therefore, this study aims to determine the total tannin content and antioxidant activity of water lettuce leaf extract before and after the purification process.

## Materials and Methods

### Extraction and Purification

The young and mature leaves of water lettuce (*Pistia stratiotes*) were collected from Sukaraja

Village, South Sumatra, Indonesia. The plants were transported to the laboratory to be cleaned and undesirable substances removed. They were cut into small pieces and dried in the oven at 45°C for 24 hours. After drying, the sample was ground into powder and kept at a cold temperature for further use. The tannin extraction was done by maceration method and 70% ethanol was used as a solvent according to the previous methods (Markom *et al.*, 2007; Sirisangsawang & Phetyim, 2023). Briefly, 20 g of dried sample was extracted using 200 mL of 70% ethanol in an Erlenmeyer flask. The extraction was performed at room temperature for three hours and stirred with a magnetic stirrer (IKA C-MAG HS 7) at 100 rpm. After the extraction, the mixture was filtered through Whatman No. 42 filter paper to separate the filtrate and residue. The filtrate was transferred to a new collection tube, whereas the residue was extracted using a fresh solvent in the same condition as the first extraction and was performed five times in total. After the extraction, the filtrate was mixed in new tubes and the solvent was evaporated using a rotary vacuum evaporator at 40°C. The extraction was fully dried using a freeze dryer (Biobase BK-FD10S, Shandong, China) to obtain powder (crude extract). The yield of extraction was calculated according to this formula:

$$\text{Yield (\%)} = \frac{\text{Extract weight (g)}}{\text{Dried sample (g)}} \times 100\% \quad (1)$$

The purification process used the solid-phase extraction (SPE) method using HyperSep Retain PEP Cartridges (Part. No. 60107-212, Thermofisher Scientific) according to the previous method (Sudirman *et al.*, 2022b). Briefly, 2 mL of distilled water (dH<sub>2</sub>O) and then 2 mL of absolute methanol were used to wash the cartridge in the preconditioning step. After preconditioning the cartridge, 2 mL of the crude extract (1 mg/10 mL) was pipetted into the cartridge and was eluted using 2 mL of *n*-hexane and 2 mL of a low concentration of sulfuric acid (H<sub>2</sub>SO<sub>4</sub> 1 N), respectively. The absolute methanol was then used to wash the cartridge to collect the purified extract, which

was dried by a freeze dryer to obtain extract powder (purified extract).

### Tannin Content Analysis

The total tannin content was analysed by Folin-Ciocalteu's method according to the previous methods (Vijaytha *et al.*, 2020; Rajkumar *et al.*, 2022). Briefly, 0.1 mL of sample solution (1 mg/mL) was pipetted into the reaction tube. The tube was added with 7.5 mL of distilled water (dH<sub>2</sub>O), 0.5 mL of Folin-Ciocalteu's reagent, and 1 mL of 35% NaCO<sub>3</sub>, the volume was made up to 10 mL by dH<sub>2</sub>O. The mixture was allowed to react at room temperature for 30 minutes. After the reaction time, the absorbance

was measured immediately by using a UV-Vis spectrophotometer (Genesys 150 Thermo-Scientific) at 700 nm. Tannic acid was used as a standard for total tannin determination. Therefore, the total tannin was expressed as mg total tannin equivalent per g of dried weight of the sample (mg TAE/g DW).

### Antioxidant Activity Assay

The antioxidant activity was determined by using the 2,2-diphenyl-1-picrylhydrazyl (DPPH)

method according to the previous methods (Chew *et al.*, 2011; Sudirman *et al.*, 2022a). Briefly, the extract was dissolved in ethanol to make a serial concentration (0-1,000  $\mu\text{g/mL}$ ). Then, 1 mL of sample solution at each concentration was mixed with 1 mL of 0.2 mM DPPH in a reaction tube. The mixture was then incubated at 37°C for 30 minutes. The absorbance was measured at 517 nm using a UV-Vis spectrophotometer. The antioxidant activity was expressed as the ability of the sample to inhibit the DPPH radical according to this formula:

$$\text{Percentage (\%)} \text{ of inhibition} = \frac{\text{Abs}_{\text{blank}} - \text{Abs}_{\text{sample}}}{\text{Abs}_{\text{blank}}} \times 100\% \quad (2)$$

where:

$\text{Abs}_{\text{blank}}$  = Absorbance at 517 nm without sample

$\text{Abs}_{\text{sample}}$  = Absorbance at 517 nm with sample

### Functional Groups Analysis by FT-IR

The functional groups of tannin-rich extract were detected using a Fourier-transform infrared (InfraRed Bruker Tensor 37) according to the previous reference (Bhateja *et al.*, 2020). The FT-IR spectra of the extract were obtained by mixing the extract with potassium bromide, further, it was pressed into pellets.

### Data Analysis

All the data were presented as mean  $\pm$  standard deviation (SD). The data were analysed using an independent t-test with SPSS 22.0 (IBM Corporation, Armonk, NY, USA). All graphics were produced using the GraphPad Prism 5.0 software (GraphPad Software, Inc., San Diego, CA, USA).

## Results and Discussion

### Yield of Extraction and Purification

The yield of extraction (crude extract) and purified extract of the water lettuce leaf is shown in Figure 1. The yield of the crude extract is about  $14.46 \pm 1.07\%$ , whereas, after purification, the yield was significantly ( $p < 0.05$ ) reduced to  $5.91 \pm 0.57\%$ . A previous study reported that the




Figure 1: The yields of extraction (crude extract) and purified extract from water lettuce (*Pistia stratiotes*) leaf. The data are shown as the mean  $\pm$  SD ( $n = 3$ ).

Statistical significance at  $*p < 0.05$  versus crude extract

yield of crude extract was reduced from 1.57% to 0.81% after purification using the solid-phase extraction (SPE) method (Sudirman *et al.*, 2023). Additionally, a previous study also reported that the purified extract of *Quercus crassifolia* was lower than the crude extract (Valencia-Avilés *et al.*, 2018). This condition indicated that some compounds were successfully removed by the purification process, resulting in a lower yield of the purified extract when compared to the crude extract. During the purification process, *n*-hexane was used to remove lipoidal substances. In contrast, the low concentration of sulfuric acid was used to elude organic acids, sugar, and other non-polyphenol polar compounds (Pérez-Magariño *et al.*, 2008). The

SPE method is widely used during polyphenol purification due to its rapidity, simplicity, and economy (Dai & Mumper, 2010).

### Tannin Content

The total tannin contents of the crude and purified extracts are shown in Figure 2. The total tannin of the purified extract was significantly higher when compared to the crude extract, at about  $13.84 \pm 0.28$  mg TAE/g of the dry sample and  $8.71 \pm 1.50$  mg TAE/g of the dry sample, respectively. Tannins are a group of polyphenolic substances that are widely found in plants (Tong *et al.*, 2022). A previous study reported that the total polyphenol content of *Inga edulis* increased from 496.5 mg GAE/g to 518.8 mg GAE/g (Souza *et al.*, 2007). Also, the flavonoid compound (luteolin) increases after the purification process from 0.68 mg/g to 3.52 mg/g (Zhao *et al.*, 2022). This condition indicated that some unwanted compounds such as sugar impurities were removed during the purification process. Similar to this present study, the purified extract contains a high concentration of tannin-polyphenol compounds due to some unwanted compounds being removed. *N*-hexane and a low concentration of sulfuric acid are used to remove undesirable compounds during SPE purification such as non-polar compounds (lipid) and non-polyphenol polar compounds (sugar and organic acids), respectively. A previous study also reported that purification using the SPE method

successfully increased the total polyphenol and flavonoid contents of *P. stratiotes* ethanol extract (Sudirman *et al.*, 2022b).

### Antioxidant Activity

The antioxidant activity of crude and purified tannin from water lettuce leaf extract is shown in Figure 3. Both of the extracts exhibit antioxidant activity. However, the antioxidant activity of the purified extract was significantly ( $p < 0.05$ ) higher than that of the crude extract, as indicated by the low half-maximum inhibition concentration ( $IC_{50}$ ) of the purified extract. A low  $IC_{50}$  value indicated that the substances exhibit higher antioxidant activity and vice-versa (Goutzourelas *et al.*, 2023). The antioxidant activity of the purified extract is about  $IC_{50} 78.45 \pm 5.42$   $\mu$ g/mL and that of the crude extract is about  $IC_{50} 135.14 \pm 22.44$   $\mu$ g/mL. In this study, we found that the purification process successfully increased the antioxidant activity of the tannin-polyphenol compound of water lettuce extract. Increased antioxidant activity in the purified extract is because this extract also contains high levels of tannin-polyphenol compounds when compared to crude extract. A previous study also reported that the purification process increased the antioxidant activity of distiller grains (Wang *et al.*, 2019). The bioactivity or quality of the crude extract increased after purification (Barbosa-Pereira *et al.*, 2013). This is because undesirable




Figure 2: The tannin content of the crude and purified extracts from water lettuce (*Pistia stratiotes*) leaf. The data are shown as the mean  $\pm$  SD ( $n = 3$ ).

Statistical significance at  $*p < 0.05$  versus crude extract




Figure 3: The antioxidant activities of the crude and purified tannin extracts from water lettuce (*Pistia stratiotes*) leaf. The data are shown as the mean  $\pm$  SD ( $n = 3$ ).

Statistical significance at  $*p < 0.05$  versus crude extract

compounds have been removed, thereby improving the overall quality of the crude extract. The polyphenol compounds, including tannin, act as antioxidants by donating the hydrogen (H) atom to the free radicals (Matyas *et al.*, 2019) and reducing the oxidation reaction by transferring electrons (Lee *et al.*, 2015).

### Functional Groups of the Tannin

The functional groups of the tannin-rich extract are shown in Figure 4. In this present study, the stretching of a polymeric hydroxyl (O–H) group and H-bonded stretching were detected at 3,334.12  $\text{cm}^{-1}$ . A previous study reported that phenolic compounds, including tannin are characterised by asymmetric and symmetric stretching of the polymeric hydroxyl group and H-bonded stretching, which is detected in the region of 3,400 to 3,200  $\text{cm}^{-1}$  (Wongsa *et al.*, 2022). Whereas, 1,618.33  $\text{cm}^{-1}$  indicates the C–O stretching vibration absorption bands and also indicates the presence of a phenolic group in the extract (1,760–1,600  $\text{cm}^{-1}$ ) (Kannan *et al.*, 2011).

### Conclusions

Overall, the tannin compounds were successfully extracted from water lettuce (*P. stratiotes*) using an ethanol solution. The purification using the

solid-phase extraction method increases the total tannin content of this extract and also increases its antioxidant activity. Therefore, tannin from water lettuce can be used as a source of natural antioxidants and formulated as a food supplement.

### Acknowledgements

This work was financially supported by the Universitas Sriwijaya within the framework of the Indonesia Directorate General of Higher Education, Research and Technology Project, Ministry of Education, Culture, Research, and Technology, SPPK No. 059/E5/PG.02.00PL/2023 and SP DIPA-023-17.1.690523/2023.

### Conflict of Interest Statement

The authors declare that they have no conflict of interest.

### References

Barbosa-Pereira, L., Pocheville, A., Angulo, I., Paseiro-Losada, P., & Cruz, J. M. (2013). Fractionation and purification of bioactive compounds obtained from a brewery waste stream. *BioMed Research International*, 2013, 1-11.




Figure 4: The FT-IR spectra of the purified tannin-rich extract from water lettuce (*Pistia stratiotes*) extract

- Bhateja, P. K., Kajal, A., & Singh, R. (2020). Amelioration of diabetes mellitus by modulation of GLP-1 via targeting alpha-glucosidase using *Acacia tortilis* polysaccharide in streptozotocin-nicotinamide induced diabetes in rats. *Journal of Ayurveda and Integrative Medicine*, *11*, 405-413.
- Chew, K. K., Ng, S. Y., Thoo, Y. Y., Khoo, M. Z., Wan Aida, W. M., & Ho, C. W. (2011). Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of *Centella asiatica* extracts. *International Food Research Journal*, *18*, 571-578.
- Dai, J., & Mumper, R. J. (2010). Plant phenolics: Extraction, analysis, and their antioxidant and anticancer properties. *Molecules*, *15*, 7313-7352.
- Das, A. K., Islam, M. N., Faruk, M. O., Ashaduzzaman, M., & Dungan, R. (2020). Review on tannins: Extraction processes, applications, and possibilities. *South African Journal of Botany*, *135*, 58-70.
- Goutzourelas, N., Kevrekidis, D. P., Barda, S., Malea, P., Trachana, V., Savvidi, S., Kevrekidou, A., Assimopoulou, A. N., Goutas, A., Liu, M., Lin, X., Kollatos, N., Amoutzias, G. D., & Stagos, D. (2023). Antioxidant activity and inhibition of liver cancer cells' growth of extracts from 14 marine macroalgae species of the Mediterranean Sea. *Foods*, *12*.
- Herpandi, Lestari, S. D., Bastian & Sudirman, S. (2021). Antioxidant activity of the fractions from water lettuce (*Pistia stratiotes*) extract. *Food Research*, *5*, 451-455.
- Kannan, R. R. R., Arumugam, R., & Perumal, A. (2011). Fourier transform infrared spectroscopy analysis of seagrass polyphenols. *Current Bioactive Compounds*, *7*, 118-125.
- Lee, C. Y., Nanah, C. N., Held, R. A., Clark, A. R., Huynh, U. G. T., Maraskine, M. C., Uzarski, R. L., Mccracken, J., & Sharma, A. (2015). Effect of electron donating groups on polyphenol-based antioxidant dendrimers. *Biochimie*, *111*, 125-134.
- Markom, M., Hasan, M., Daud, W., Singh, H., & Jahim, J. (2007). Extraction of hydrolysable tannins from *Phyllanthus niruri* Linn.: Effects of solvents and extraction methods. *Separation and Purification Technology*, *52*, 487-496.
- Matyas, M., Hasmasanu, M. G., & Zaharie, G. (2019). Antioxidant capacity of preterm neonates assessed by hydrogen donor value. *Medicina*, *55*, 720.
- Pérez-Magariño, S., Ortega-Heras, M., & Cano-Mozo, E. (2008). Optimization of a solid-phase extraction method using copolymer sorbents for isolation of phenolic compounds in red wines and quantification by HPLC. *Journal of Agricultural and Food Chemistry*, *56*, 11560-11570.
- Rad, M. S., Anil Kumar, N. V., Zucca, P., Varoni, E. M., Dini, L., Panzarini, E., Rajkovic, J., Tsouh Fokou, P. V., Azzini, E., Peluso, I., Prakash Mishra, A., Nigam, M., El Rayess, Y., Beyrouthy, M. E., Polito, L., Iriti, M., Martins, N., Martorell, M., Docea, A.O., Setzer, W. N., Calina, D., Cho, W. C., & Sharifi-Rad, J. (2020). Lifestyle, oxidative stress, and antioxidants: Back and forth in the pathophysiology of chronic diseases. *Frontiers in Physiology*, *11*.
- Rajkumar, G., Panambara, P. A. H. R., & Sanmugarajah, V. (2022). Comparative analysis of qualitative and quantitative phytochemical evaluation of selected leaves of medicinal plants in Jaffna, Sri Lanka. *Borneo Journal of Pharmacy*, *5*, 93-103.
- Sirisangsawang, R., & Phetyim, N. (2023). Optimization of tannin extraction from coconut coir through response surface methodology. *Heliyon*, *9*.
- Souza, J. N. S., Silva, E. M., Silva, M. N. D., Arruda, M. S. P., Larondelle, Y., & Rogez, H. (2007). Identification and antioxidant

- activity of several flavonoids of *Inga edulis* leaves. *Journal of the Brazilian Chemical Society*, 18, 1276-1280.
- Spiegel, M., Cel, K., & Sroka, Z. (2023). The mechanistic insights into the role of pH and solvent on antiradical and prooxidant properties of polyphenols — Nine compounds case study. *Food Chemistry*, 407.
- Sudirman, S., Herpandi, Nopianti, R., Lestari, D. S., Wasahla, W., & Mareta, H. (2017). Phenolic contents, tannin, vitamin C, and vitamin E of water lettuce (*Pistia stratiotes*). *Oriental Journal of Chemistry*, 33, 3173-3176.
- Sudirman, S., Herpandi, Safitri, E., Apriani, E. F., & Taqwa, F. H. (2022a). Total polyphenol and flavonoid contents and antioxidant activities of water lettuce (*Pistia stratiotes*) leave extracts. *Food Research*, 6, 205-210.
- Sudirman, S., Janna, M., Herpandi & Widiastuti, I. (2022b). *In vitro* inhibitory HMG-CoA reductase activity of purified polyphenol compounds from water lettuce (*Pistia stratiotes*) leaf extract. *Tropical Journal of Natural Product Research*, 6, 1131-1134.
- Sudirman, S., Pujiastuti, N., Janna, M., Herpandi & Widiastuti, I. (2023). Antihyperlipidemic activity of purified polyphenol extracted from water lettuce (*Pistia stratiotes*) leaf: An *in vitro* analysis. *Tropical Journal of Natural Product Research*, 7, 3177-3181.
- Tong, Z., He, W., Fan, X., & Guo, A. (2022). Biological function of plant tannin and its application in animal health. *Frontiers in Veterinary Science*, 8.
- Valencia-Avilés, E., García-Pérez, M., Garnica-Romo, M., Figueroa-Cárdenas, J., Meléndez-Herrera, E., Salgado-Garciglia, R., & Martínez-Flores, H. (2018). Antioxidant properties of polyphenolic extracts from *Quercus Laurina*, *Quercus Crassifolia*, and *Quercus Scytophylla* Bark. *Antioxidants*, 7.
- Vijaytha, V., Anupama, R. V., & Haridas, M. (2020). Phytochemical profiling and anti-oxidant, anti-bacterial, and anti-inflammatory properties of *Viburnum coriaceum* Blume. *Future Journal of Pharmaceutical Sciences*, 6.
- Vona, R., Pallotta, L., Cappelletti, M., Severi, C., & Matarrese, P. (2021). The impact of oxidative stress in human pathology: Focus on gastrointestinal disorders. *Antioxidants*, 10.
- Wang, X., Wang, S., Huang, S., Zhang, L., Ge, Z., Sun, L., & Zong, W. (2019). Purification of polyphenols from distiller's grains by macroporous resin and analysis of the polyphenolic components. *Molecules*, 24.
- Wongsa, P., Phatikulrungsun, P., & Prathumthong, S. (2022). FT-IR characteristics, phenolic profiles and inhibitory potential against digestive enzymes of 25 herbal infusions. *Scientific Reports*, 12.
- Zeka, K., Ruparella, K., Arroo, R., Budriesi, R., & Micucci, M. (2017). Flavonoids and their metabolites: Prevention in cardiovascular diseases and diabetes. *Diseases*, 5.
- Zhao, H., Avena-Bustillos, R. J., & Wang, S. C. (2022). Extraction, purification, and *in vitro* antioxidant activity evaluation of phenolic compounds in California olive pomace. *Foods*, 11, 174.