

## MITIGATION N<sub>2</sub>O EMISSION THROUGH BEEF CATTLE WASTE FERTILISATION APPLICATION IN CORN FIELD

SRI ARNITA ABU TANI<sup>1\*</sup>, FACHROERROZI HOESNI<sup>1</sup>, ALI PURNOMO<sup>2</sup>, SAITUL<sup>1</sup> AND ANURAGA JAYANEGARA<sup>3</sup>

<sup>1</sup>Department of Animal Production, Faculty of Animal Science, Jambi University Mendalo Darat, Muarojambi, 36361 Jambi, Indonesia. <sup>2</sup>Indonesian Agricultural Environment Standardisation Institute, Jl. Jakenan-Jaken Km. 5, 59182 Pati, Central Java, Indonesia. <sup>3</sup>Department of Nutrition and Feed Technology, Faculty of Animal Science, IPB University, 16680 Bogor, West Java, Indonesia.

\*Corresponding author: sriarnita.1963@gmail.com

<http://doi.org/10.46754/jssm.2024.10.012>

Received: 22 June 2024

Accepted: 14 August 2024

Published: 15 October 2024

**Abstract:** This study is aimed at explaining the pattern of N<sub>2</sub>O emission from the use of Cattle Manure Fertiliser (bio-slurry, bio-urine, and Trichocompost) on sweet corn (*Zea mays L. Saccharata*). This study used a Randomised Block Design (RBD) with four treatment groups and four replications. Each test consisted of 16 experimental units. The treatments consisted of combinations of Chemical Fertiliser (SP36, KCl, urea) and Organic Fertiliser (bio-urine and bio-slurry, Trichocompost), i.e., 100% Chemical Fertiliser (C<sub>100</sub>), 75% Chemical Fertiliser + 25% Organic Fertiliser (C<sub>75</sub>O<sub>25</sub>), 50% Chemical Fertiliser + 50% Organic Fertiliser (C<sub>50</sub>O<sub>50</sub>), and 25% Chemical Fertiliser + 75% Organic Fertiliser (C<sub>25</sub>O<sub>75</sub>). The observed variables were the pattern of N<sub>2</sub>O flux after the first and second fertilisation and the emission of N<sub>2</sub>O. The results showed that the highest N<sub>2</sub>O flux in the first fertilisation resulted from C<sub>75</sub>O<sub>25</sub> treatment while in the second fertilisation, the highest N<sub>2</sub>O flux was emitted from C<sub>100</sub> and the lowest was emitted from C<sub>25</sub>O<sub>75</sub>. The lowest N<sub>2</sub>O emission was from C<sub>25</sub>O<sub>75</sub> treatment. In conclusion, using beef cattle waste fertiliser could decrease N<sub>2</sub>O flux and emissions by 75% in sweet corn cropping.

Keywords: Beef cattle, manure, fertiliser, emission, global warming.

### Introduction

The accumulation of greenhouse gases (GHG) such as carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) in the atmosphere causes climate change, also referred to as global warming. This increase can be caused by industrial activities, fossil fuel combustion, and the agricultural and livestock sectors (IPCC, 2013). In the livestock sector, beef cattle waste contributes up to 40% of greenhouse gases (GHG) in the atmosphere (Grossi *et al.*, 2019) and agricultural sector also contributes around 10 to 12% to greenhouse gas emissions (Pramono *et al.*, 2021). These greenhouse gas emissions are expected to continue to increase until 2030 as food demand increases. This increase in greenhouse gas emissions will contribute to global warming as CH<sub>4</sub> in the atmosphere is 25 times greater than CO<sub>2</sub> while N<sub>2</sub>O is 298 times higher (Valero, 2019).

GHG emissions from the agriculture and livestock sectors are closely related

to conventional farming. At present, beef cattle waste in the form of solid waste (cow manure) and liquid (urine) are not disposed of properly. The beef cattle waste will pollute the environment and potentially increase GHG emissions. The accumulation of beef cattle waste can produce high amounts of CH<sub>4</sub> gas, which contributes to global warming (Chadwick *et al.*, 2011). Furthermore, the intensive use of chemicals by farmers increases the production costs and has the potential to contribute to GHG emissions in the form of N<sub>2</sub>O gas (Nenobesi *et al.*, 2017).

The large contribution of the livestock and agriculture sectors to global warming must be mitigated. By making solid and liquid beef cattle waste as sources of organic fertilisers in the form of trichocompost, bio-urine, and bio-slurry, GHG emissions could be reduced. As an organic material, trichocompost can improve agricultural land by adding nutrients that plants

need. This is expected to increase productivity, lower the cost of chemical fertilisers, and preserve environmental quality (Hartati *et al.*, 2016). Liquid fertiliser in the form of bio-urine, in addition to containing high amounts of nutrients, also contains natural growth regulators and pest and disease-repellent compounds. The natural growth regulator contains hormones from the IAA, Gibberellin, and cytokinin groups (Sirajuddin *et al.*, 2015). Bio-slurry is the final product after processing beef cattle solid waste for biogas energy. The liquid bio-slurry has no damaging effect on the soil or plants and can act as a binding agent so that the fertiliser solution applied to the soil surface can be directly utilised by plants. It also contains microbes that facilitate the fertilisation of the soil (Kabedey *et al.*, 2023).

Therefore, it is necessary to assess how much of N<sub>2</sub>O gas emissions are mitigated through the application of organic fertiliser based on beef cattle waste compared with chemical fertilisers.

## Materials and Methods

### Materials

The research was conducted in Pudak Community Economic Zone, Kumpeh Ulu Village, Muaro Jambi Regency, Jambi Province, Indonesia. The materials used were Bonanza F1 seed (sweet corn type), bio-urine, bio-slurry, Trichocompost as Organic Fertiliser and KCL (*Kalium Chloride*), Dolomite, and Urea as Chemical Fertiliser. The gas measurement apparatus comprised N<sub>2</sub>O gas closed chamber (50 cm x 30 cm x 25 cm), vial storage, bubble wrap, septum rubber, thermometer, aluminium foil paper, and 10 ml syringe.

### Research Method

#### Land Preparation

A 0.25 ha plot planted with sweet corn was prepared. The ground was tilled using a hand tractor that was divided into 16 pillows around 100 cm in height of ± 10 cm. Two rows of

plants were planted in each pillow designed in a randomised complete block design, consisting of four treatments and eight replications. The soil type was Alluvial with a pH of 4.7 and chemical composition of N 0.05%; P 5.3 ppm; K 0.05%; and C 1.0%.

### Planting Process

Corn seeds were planted into the soil at a depth of 3 cm using the *legowo* row system. Each pillow had two rows of plants with a spacing 25 cm x 50 cm. If a seed did not grow or the seedling died, a replacement plant prepared in the reserve yard was planted.

### Fertilisation

Chemical and organic fertilisation applications in sweet corn fields according to the treatments are presented in Table 1.

### Chemical Fertiliser Applications

The chemical fertilisation was done by digging a hole about 5 cm from the plants. Dolomite is applied three days before planting. For KCl and SP36 fertilisation, they were applied at 14 DAPs, Urea were applied two times at 14 DAPs and 30 DAPs.

### Organic Fertilisers

The organic fertiliser used in this study is a combination of solid and liquid waste from beef cattle, which includes Trichocompost, bio-urine, and bio-slurry. Trichocompost is processed beef cattle feces that have been treated with the decomposer *Trichoderma harzianum* sp. which was isolated at the Jambi Province Food Plant Pest Disease Laboratory. Trichocompost was applied by sprinkling it on the planting holes that had been prepared according to the treatment given, three days before planting at the beginning of planting. Bio-urine is beef cattle urine that has undergone fermentation for approximately one month while bio-slurry is the final product of biogas. Bio urine is applied first, followed by bio slurry by spraying it onto each

Table 1: Application of chemical and organic fertilisers in sweet corn fields (*Zea mays* L.Saac)/0.25 ha

No.	Treatment	3 Days Before Planting	Total (kg <sup>-1</sup> 0.25 ha)	14 Days After Planting (DAPs)	Total (kg <sup>-1</sup> 0.25 ha)	30 Days After Planting (DAPs)	Total (kg <sup>-1</sup> 0.25 ha)		
1	CF <sub>100</sub>	Dolomite	500	SP <sub>36</sub>	50	Urea <sub>2</sub>	37.5		
				KCL	25				
				Urea <sub>1</sub>	37.5				
2	C <sub>75</sub> O <sub>25</sub>	Trichocompost	125	SP <sub>36</sub>	37.5	Urea <sub>2</sub>	28.13		
				KCL	18.75				
				Urea <sub>1</sub>	28.13				
3	C <sub>50</sub> O <sub>50</sub>	Trichocompost	250	SP <sub>36</sub>	25	Urea <sub>2</sub>	18.75		
				KCL	12.5				
				Urea <sub>1</sub>	18.75				
				Bio-urine <sub>1</sub>	12.5 ltr (Kg <sup>-1</sup> ha)			Bio-urine <sub>2</sub>	12.5 ltr (Kg <sup>-1</sup> ha)
4	C <sub>25</sub> O <sub>75</sub>	Trichocompost	375	SP <sub>36</sub>	12.5	Urea <sub>2</sub>	9.375		
				KCL	6.25				
				Urea <sub>1</sub>	9.375				
				Bio-urine <sub>1</sub>	12.5 ltr (kg <sup>-1</sup> ha)			Bio-urine <sub>2</sub>	12.5 ltr (kg <sup>-1</sup> ha)
				Bio-slurry <sub>1</sub>	12.5 ltr (kg <sup>-1</sup> ha)			Bio-slurry <sub>2</sub>	12.5 ltr (kg <sup>-1</sup> ha)

Note: \*Recommended use of chemical and organic fertilisers in sweet corn crops: Super Phosphate (SP36) 200 kg<sup>-1</sup>ha, Potassium Chloride (KCL) 100 kg<sup>-1</sup>ha, Urea 300 kg<sup>-1</sup>ha, and dolomite 2,000 kg<sup>-1</sup>ha (Nurmegawati *et al.*, 2015). Trichocompost 2,000 kg<sup>-1</sup>ha, bio-urine 100 ltr<sup>-1</sup>ha, and bio-slurry 100 ltr<sup>-1</sup>ha (Tani, 2017).

plant from the tip of the leaves to the ground. Bio urine and bio slurry applications were applied at 14 DAPs and 30 DAPs.

### Nitrous Oxide Gas (N<sub>2</sub>O) Measurement

During this research, the N<sub>2</sub>O gas sampling was implemented in two stages. The first stage consisted of three, six, and nine days after the first fertilisation with 14 DAPs and the second stage also being done in three, six, and nine days after the second fertilisation with 30 DAPs. The stages of N<sub>2</sub>O emission were running in these steps: Placing N<sub>2</sub>O gas capture chamber (50 cm x 30 cm x 25 cm) between two corn plants

within the same pillow and the gas capturing process was conducted in the morning.

- Installing a thermometer to measure the temperature after the gas taking has been completed. The range of time that being set as benchmark were in 10, 20, 30, 40, and 50 minutes.
- N<sub>2</sub>O gas sampling process was running by injection of 10 ml with wrapping the syringe surface by aluminium foil to prevent gas leaking during the process.
- Immediately closing the syringe that contains the gas to ensure the leakage of captured gas.

(d) Analysing the gas concentration in Greenhouse Gas Laboratory by using chromatography gas.

**Research Design**

This research was carried out by using a randomised complete block design (RCBD), where it has been implemented in four treatment groups where each group included two experimental units. The treatment process consisted of these elements:

- (1) C<sub>100</sub> : 100% Chemical Fertiliser
- (2) C<sub>75</sub>O<sub>25</sub> : 75% Chemical Fertiliser + 25% Organic Fertiliser
- (3) C<sub>50</sub>O<sub>50</sub> : 50% Chemical Fertiliser + 50% Organic Fertiliser
- (4) C<sub>25</sub>O<sub>75</sub> : 25% Chemical Fertiliser + 75% Organic Fertiliser

**Observed Variables**

The flux of N<sub>2</sub>O gas in the first and second fertilisation pattern measured in kg ha<sup>-1</sup>day<sup>-1</sup> and N<sub>2</sub>O gas emissions measured in kg ha<sup>-1</sup>season<sup>-1</sup>.

**Data Analysis**

Flux data and N<sub>2</sub>O emissions are calculated with the (IAEA, 1992):

$$E = \frac{dc}{dt} \times V_{ch} / A_{ch} \times mW / mV \times 273.2 / (273.2 + T)$$

where:

- E = N<sub>2</sub>O gas emission (mg/m<sup>2</sup>/day)
- dc/dt = the difference N<sub>2</sub>O per time given (ppm/minute)
- V<sub>ch</sub> = the volume of box (m<sup>3</sup>)
- A<sub>ch</sub> = the box area (m<sup>2</sup>)
- mW/mV = gas molecular weight/molecular volume constant N<sub>2</sub>O (22.41 l)
- T = average temperature measured during sampling (°C)
- 273.2 value = Kelvin temperature constant degree

Greenhouse gas emission (N<sub>2</sub>O) were being calculated and measured in Analysis of Variance and if there were discrepancies between treatments, they were tested further by using Duncan’s test. The data processing in this research was conducted in Statistical Analysis System (SAS) software.

**Results and Discussion**

***Patterns of N<sub>2</sub>O Flux from Application of Chemical Fertiliser and Organic Fertiliser at First Fertilisation in Sweet Corn Fields***

The N<sub>2</sub>O flux emitted from various levels of Trichocompost, bio-urine, and bio-slurry applications on sweet corn plantation (*Zea mays L. Saccharata*) in the first fertilisation is presented in Figure 1.




Figure 1: N<sub>2</sub>O flux in first fertilisation with four treatments: 100% Chemical (C<sub>100</sub>); 75% Chemical + 25% Organic (C<sub>75</sub>O<sub>25</sub>); 50% Chemical + 50% Organic (C<sub>50</sub>O<sub>50</sub>); and 25% Chemical + 75% Organic (C<sub>25</sub>O<sub>75</sub>), respectively

The N<sub>2</sub>O flux from the application of a combination of chemical fertilisers and organic fertilisers (Trichocompost, bio-urine, and bio-slurry) on sweet corn fields was quite varied, where at the beginning of fertilisation (1<sup>st</sup> fertilisation) the combination of CF+OF (75% and 25%) emitted higher N<sub>2</sub>O than the application of chemical fertiliser (100%), CF+OF (50+50%) and CF+OF (25% and 75%). This shows that after the first fertilisation, the N<sub>2</sub>O flux from the plots C<sub>75</sub>O<sub>25</sub> was highest than others the plots, may caused by the organic fertiliser was still undergoing decomposition and still supplies energy for denitrification. N from fertiliser, dissolved organic carbon and inorganic N from soil organic matter decomposition and N mineralisation also contribute to N<sub>2</sub>O production (Wu *et al.*, 2017). Fertilisation is a major factor that contributes to the release and addition of N<sub>2</sub>O. The higher N<sub>2</sub>O flux pattern in C<sub>75</sub>O<sub>25</sub> is due to chemical fertilisers that can cause pollution and global warming. Chemical fertilisers themselves contain nitrogen needed by plants but not all nitrogen is absorbed by plants. Some of the nitrogen will be broken down by microorganisms or flows with water and binds with oxygen to form nitrous oxide gas (N<sub>2</sub>O) (Dalman *et al.*, 2021).

### N<sub>2</sub>O Production Pattern at the Time of Second Fertilisation in Sweet Corn Field

In the second fertilisation, the use of organic fertiliser (beef cattle waste) consisting of trichocompost, bio-urine, and bio-slurry with a combination of C<sub>75</sub>O<sub>25</sub>, C<sub>50</sub>O<sub>50</sub>, and C<sub>25</sub>O<sub>75</sub> was seen to decrease while the use of 100% chemical fertiliser (C<sub>100</sub>), N<sub>2</sub>O flux was seen to be the highest of the other treatments. This result shows that at the age of 15 days after fertilisation, the absorption of fertilisers by plants has gone well. The pattern of the decline in N<sub>2</sub>O flux is presented in Figure 2.

The results showed that the application of 100% chemical fertiliser (C<sub>100</sub>) on sweetcorn fields increased the highest N<sub>2</sub>O flux compared to the treatment with organic fertiliser (beef cattle waste). The application of NPK fertiliser can increase the level of nitrogen (N) in the soil so that the demand for N for the bacterial transformation process can be met in the long term (Jain *et al.*, 2010). Nitrogen is the main component in the formation of N<sub>2</sub>O, where N<sub>2</sub>O is formed from the conversion of ammonium to nitrate through the process of nitrification and then to N<sub>2</sub>O through the process of denitrification (Suntoro *et al.*, 2013). N<sub>2</sub>O fluxes increase with increasing soil N (Weitz *et al.*, 2001). N is required for plant growth in two forms,




Figure 2: N<sub>2</sub>O flux in second fertilisation with four treatments: 100% Chemical (C<sub>100</sub>); 75% Chemical + 25% Organic (C<sub>75</sub>O<sub>25</sub>); 50% Chemical + 50% Organic (C<sub>50</sub>O<sub>50</sub>); and 25% Chemical + 75% Organic (C<sub>25</sub>O<sub>75</sub>), respectively

ammonium ( $\text{NH}_4^{+}$ ) when the soil is wet and nitrate ( $\text{NO}_3^-$ ) when the soil is dry. In addition,  $\text{N}_2\text{O}$  can be formed by natural processes, namely the activity of microorganisms.  $\text{N}_2\text{O}$  is formed naturally through the processes of nitrification and denitrification as by-products of the nitrogen cycle (Bracmort, 2010).

Nitrification is the aerobic reduction of  $\text{NH}_3$  to nitrite and nitrate. The first step in the nitrification process is the oxidation of  $\text{NH}_3$  to  $\text{NO}_2$  by ammonia oxidising bacteria such as Nitrosomonas. The next stage is the oxidation of  $\text{NO}_2$  to  $\text{NO}_3$  by nitrite oxidising bacteria such as Nitrobacter. The formation of  $\text{N}_2\text{O}$  occurs when Anammox bacteria reduce ammonia to  $\text{N}_2$  gas with the electron acceptor  $\text{NO}_2$  (Bernhard, 2010).

Denitrification is the process of reducing  $\text{NO}_3^-$  to  $\text{NO}_2^-$  and further to  $\text{NO}$ ,  $\text{N}_2\text{O}$ , and  $\text{N}_2$ , with  $\text{N}_2$  being the main product of the denitrification process.  $\text{N}_2\text{O}$  can be formed by natural processes, namely the activity of microorganisms.  $\text{N}_2\text{O}$  is formed naturally through nitrification and denitrification processes as by-products of the nitrogen cycle (Bracmort, 2010).

Nitrification is the aerobic reduction of  $\text{NH}_3$  to nitrite and nitrate. The first step in the nitrification process is the oxidation of  $\text{NH}_3$  to  $\text{NO}_2$  by ammonia oxidising bacteria such as Nitrosomonas. The next stage is the oxidation of  $\text{NO}_2$  to  $\text{NO}_3$  by nitrite oxidising bacteria such as Nitrobacter. The formation of  $\text{N}_2\text{O}$  occurs when Anammox bacteria reduce ammonia to  $\text{N}_2$  gas with the electron acceptor  $\text{NO}_2$  (Bernhard, 2010).

Denitrification is the process of reducing  $\text{NO}_3^-$  to  $\text{NO}_2^-$  and further to  $\text{NO}$ ,  $\text{N}_2\text{O}$ , and  $\text{N}_2$ , with  $\text{N}_2$  being the main product of the denitrification process. This process is also referred to as enzymatic denitrification, which is different from the assimilatory reduction of  $\text{NO}_3^-$  carried out by various biota for their growth and also different from the dissimilatory reduction of  $\text{NO}_3^-$  to  $\text{NH}_4^{+}$  carried out by some microbes in the absence of oxygen (Hadisudarmo, 2009). The process of nitrification by producing  $\text{N}_2\text{O}$  in small amounts. Whereas  $\text{NO}_3^-$  can be reduced

through denitrification in a slightly aerobic state to  $\text{N}_2\text{O}$ , in this process  $\text{N}_2\text{O}$  are formed (Ramos *et al.*, 2019).

The decrease in  $\text{N}_2\text{O}$  flux production in the second fertilisation in sweet corn land, the application of organic fertiliser based on beef cattle waste (Trichocompost, bio-urine, and bio-slurry) by 25% ( $\text{C}_{25}\text{O}_{75}$ ), 50% ( $\text{C}_{50}\text{O}_{50}$ ), and 75% ( $\text{C}_{75}\text{O}_{25}$ ), the higher the amount of organic fertiliser used, the lower the  $\text{N}_2\text{O}$  flux value. Shao *et al.* (2017) stated that the soil  $\text{N}_2\text{O}$  emission flux decreased as the proportion of organic fertiliser increased. Organic fertilisers are able to maintain land fertility and productivity in a sustainable manner. The combination of inorganic and organic fertiliser applications can increase land productivity sustainably, increase N use efficiency, and reduce environmental pollution. The combination of inorganic and organic N fertilisation can maintain soil fertility and crop productivity in the long term (Eche *et al.*, 2013).

### ***Effect of the Organic Fertiliser Application on Greenhouse Gas ( $\text{N}_2\text{O}$ ) Emissions***

The  $\text{N}_2\text{O}$  emissions generated from the four application patterns of chemical fertiliser and beef cattle waste fertiliser (organic fertiliser), it can be seen that the higher the organic fertiliser applied to sweet corn fields (75%) can reduce  $\text{N}_2\text{O}$  emissions. Results are presented in Figure 3.

The results showed that using 100% chemical fertiliser ( $\text{C}_{100}$ ) produced the highest  $\text{N}_2\text{O}$  emissions, followed by application of Organic Fertiliser 75% ( $\text{C}_{75}\text{O}_{25}$ ), 50% ( $\text{C}_{50}\text{O}_{50}$ ), and 25% ( $\text{C}_{25}\text{O}_{75}$ ). The high  $\text{N}_2\text{O}$  emissions at 100% Chemical Fertiliser application ( $\text{C}_{100}$ ) are due to  $\text{N}_2\text{O}$  emitted from nitrogen-fertilised soils due to nitrification and denitrification processes. Agricultural land has the potential to increase its  $\text{N}_2\text{O}$  gas emissions when the amount of N available for microbial transformation is increased through inorganic N fertilisation, return of organic fertilisers and crop residues to the soil, mineralisation of soil biomass and other forms of organic matter (Efosa *et al.*, 2023). Besides fertilisation, some factors that




Figure 3: N<sub>2</sub>O emission (kg CO<sub>2</sub>-e ha<sup>-1</sup> season<sup>-1</sup>) on sweet corn fields with four treatments: 100% Chemical (C<sub>100</sub>); 75% Chemical + 25% Organic (C<sub>75</sub>O<sub>25</sub>); 50% Chemical + 50% Organic (C<sub>50</sub>O<sub>50</sub>); and 25% Chemical + 75% Organic (C<sub>25</sub>O<sub>75</sub>), respectively

can affect N<sub>2</sub>O emissions are temperature, soil type, vegetation type, different climatic and soil conditions, location, and sampling time.

The amount of soil-borne N<sub>2</sub>O emissions correlates with the amount of nitrogen (N) available in the soil. Adding organic matter such as compost and manure to the soil in maize crops has the potential to increase soil C storage while providing a source of N. Replacing synthetic N fertilisers with organic matter will also reduce the environmental costs associated with N fixation by recycling large amounts of available N resources. Inorganic fertilisation causes an increase in N<sub>2</sub>O flux (Mishra *et al.*, 2012). Organic fertilisation with biochar can reduce greenhouse gas emissions and increase land productivity. Charcoal is beneficial for soil management, carbon sequestration, and immobilising pollutants (Kajitani *et al.*, 2013). Charcoal can reduce the rate of N<sub>2</sub>O production by improving soil physical properties (diffusion, aggregation, and water-holding capacity), soil chemical properties (acidity, available nitrogen nutrients from minerals and organic matter, dissolved organic carbon), and soil biological properties (number of microorganisms and activity of macrofauna) (Lorenz & Lal, 2014). The organic fertiliser used in this study contains charcoal, which is very useful for improving soil

fertility and increasing plant growth. Charcoal has specific properties that, when incorporated into the soil, increase soil pH and nutrient holding capacity, making fertilisation more efficient (Wang *et al.*, 2017). The use of solid and liquid beef cattle waste in food crops on tidal swamp can reduce N<sub>2</sub>O emissions (Tani, 2017). N<sub>2</sub>O production increases, both through the process of nitrification and through the process of denitrification, when the soil is given an excessive application of N fertiliser (Tracy *et al.*, 2017).

## Conclusions

The highest N<sub>2</sub>O flux in the first fertilisation was resulted from C<sub>75</sub>O<sub>25</sub> treatment while in the second fertilisation, the highest N<sub>2</sub>O flux was emitted from C<sub>100</sub> and the lowest was C<sub>25</sub>O<sub>75</sub>. By applying organic fertiliser (beef cattle waste) to substitute N fertiliser rate on corn cropping could reduce N<sub>2</sub>O emission 23 to 75% in our field. The highest reduction was reached by combining 25% chemical fertiliser and 75% organic fertiliser. Fertiliser management could be key factors to reducing GHG emissions in tropical regions. The application of beef cattle waste fertiliser could reduce N<sub>2</sub>O flux and emission up to 75% from sweet corn cropping.

## Acknowledgements

The authors are grateful to the Indonesian Agricultural Environment Standardisation Institute for their technical assistance and IPB University during the experiment and University of Jambi for its facilities. We are also very grateful to two anonymous reviewers who provided us with invaluable suggestions.

## Conflict of Interest Statement

The authors declare that they have no conflict of interest.

## References

- Bernhard, A. (2010). The nitrogen cycle: Processes, players, and human impact. *Nature Education Knowledge*, 2(2), 1-9.
- Bracmort, K. (2010). Nitrous Oxide from agricultural sources: Potential role in greenhouse gas emission reduction and ozone recovery. *Congressional Research Service*, (2), 1-9.
- Chadwick, D., Sommer, S., Thorman, R., Fongueiro, D., Cardenas, L., Amon, B., & Misselbrook, T. (2011). Manure management: Implications for greenhouse gas emissions. *Animal Feed Science and Technology*, 166-167, 514-531. <https://doi.org/10.1016/j.anifeedsci.2011.04.036>
- Eche, N. M., Iwuafor, E. N. O., Amapui, I. Y., & Bruns, M. V. (2013). Effect of application of organic and mineral soil amendment in continuous cropping system for 10 years on chemical and physical properties of an Alfisol in Northern Guinea savanna zone: Experimental investigation. *International Journal of Agricultural Policy and Research*, 1(4), 116-123. <https://doi.org/10.9734/bpi/rppsr/v3/1705C>
- Efosa, N., Krause, H-M., Hüppi, R., Krauss, M., Vaucher, N., Zourek, F., Maye, J., Six, J., & Bünemann, E-K. (2023). Emissions of Nitrous Oxide and Methane after field application of liquid organic fertilisers and biochar. *Agriculture, Ecosystems & Environment*, 356, 1-11. <http://dx.doi.org/10.1016/j.agee.2023.108642>
- Dalmau, J. M., Berbel, J., & Fernandez, R. O. (2021). Nitrogen fertilisation. A review of the risk associated with the inefficiency of its use and policy responses. *Sustainability*, 13(10), 5625. <https://doi.org/10.3390/su13105625>
- Grossi, G., Goglio, P., Vitali, A., & Williams, A. G. (2019). Livestock and climate change: Impact of livestock on climate and mitigation strategies. *Animal Frontiers*, 9(1), 69-76. <https://doi.org/10.1093/af/vfy034>
- Hadisudarmo, Purwanto. (2009). Soil Biology Environmentally Sound Soil. Management Studies. Indonesia Cerdas. Yogyakarta.
- Hartati, R., Yetti, H., & Puspita, F. (2016). The provision of Trfichocompost various organic substances to growth and the production of Sweet Corn (*Zea mays Saccharata Strt.*). *Jurnal Online Mahasiswa Fakultas Pertanian*, 3(1), 1-15.
- International Atomic Energy Agency. (1992). *Manual of measurement of Methane and Nitrous Oxide Emission from agricultural*. International Energy Agency. pp. 56-57.
- IPCC. (2013). *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK & New York, NY, USA: Cambridge University Press.
- Jain, M. C., Kumar, S., Wassmann, S., Mitra, S., Singh, S. D., Singh, J. P., Singh, S., Yadav, A. K., & Gupta, S. (2010). Methane emissions from irrigated rice fields in Northern India (New Delhi). *Nutrient Cycling in Agroecosystems*, 58(4), 75-83. <http://dx.doi.org/10.1023/A:1009882216720>
- Kabedey, T., Kanenia, Y. G., Senbetaa, A. F., & Simea, G. (2023). Effect of bioslurry and chemical fertiliser on the agronomic performances of maize. *Heliyon*, 9(1), 1-8.

- <https://doi.org/10.1016/j.heliyon.2023.e13000>
- Kajitani, S., Tay, L. H., Zhang, S., & Li, Z. C. (2013). Mechanisms and kinetic modelling of steam gasification of brown coal in the presence of volatile-char interactions. *Fuel*, *103*, 7-13. <https://doi.org/10.1016/j.fuel.2011.09.059>
- Lehmann, J., Gaunt, J., & Rondon, M. (2006). Bio-char sequestration in terrestrial ecosystems – A review. *Mitigation and Adaptation Strategies for Global Change*. Springer. 11: 403-427. 1-96. <https://doi.org/10.1007/s13593-012-0081-1>
- Lorenz, K., & Lal, R. (2014). Biochar application to soil for climate change mitigation by soil organic carbon sequestration. *Journal of Plant Nutrition and Soil Science*, *177*(5), 651-670. <http://doi.org/10.1002/jpln.201400058>
- Lu, L., Wassmann, R., Neue, H. U., Huang, C., & Bueno, C. S. (2000). Methanogenic responses to exogenous substrates in anaerobic rice soil. *Soil Biology and Biochemistry*, *32*, 1683-1690. [https://doi.org/10.1016/S0038-0717\(00\)00085-7](https://doi.org/10.1016/S0038-0717(00)00085-7)
- Meng, L., Ding, W., & Cai, Z. (2005). Long-term application of organic manure and nitrogen fertiliser on N<sub>2</sub>O emissions, soil quality and crop production in sandy loam soil. *Soil Biology and Biochemistry*, *37*(11), 2037-2045. <https://doi.org/10.1016/j.soilbio.2005.03.007>
- Mishra, S. N., Mitra, S., Rangan, L., Dutta, S., & Singh, P. (2012). Exploration of 'hot-spots' of methane and nitrous oxide emission from the agriculture fields of Assam, India. *Agriculture & Food Security*, *1*, 16. <http://dx.doi.org/10.1186/2048-7010-1-16>
- Nenobesi, D., Mella, W., & Soetedjo, P. (2017). Utilisation of solid waste manure compost in improving environmental supportability and biomass of green bean (*Vigna Radiata* L). *Jurnal Pangan*, *26*(1), 43-56. <https://doi.org/10.33964/jp.v26i1.344>
- Nurmegawati, Yahumri & Afrizon. (2015). Recommendation on fertiliser of corn and soybean crops in Kaur District, Bemgkulu. *Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia*, *1*(4), 914-917. <https://doi.org/10.13057/psnmbi/m010446>
- Pramono, A., Andriany, T. A., Susilawati, T. H., & Sutriadi, M. T. (2021). Global warning potential from maize cultivation as affective by organic urea fertiliser in rainfed lowland. *International Conference on Agro-Industry. IOP Conference Series: Earth and Environmental Science*, *733* 012144. <https://doi.org/10.1088/1755-1315/733/1/012144>
- Ramos, T. M., Jay-Russel, M. T., Miller, P. D., Shade, J., Misslewicz, T., Sorge, U. S., Hutchinson, M., Lilley, J., & Pires, A. F. A. (2019). Assesment of biological soil amandements of animal, origin use, research needs and extension oppurtunities in organic production. *Frontiers in Sustainable Food Systems*, *3*(73), 4-11. <https://doi-org/103389/fstufs.2019.00073>
- Shao, J., Chen, J., Wang, L., Hou, M., & Chen, D. (2021). Effects of fermented organic fertiliser application on soil N<sub>2</sub>O emission under the vegetable rotation in polyhouse. *Environmental Research*, *200*, 1-9. <https://doi.org/10.1016/j.envres.2021.111491>
- Sirajuddin, S. N., Ikrar, M. S., & Kasniyati, K. (2015). Perception of farmers against liquid fertiliser benefit of beef cattle urine. *Proceeding of 38<sup>th</sup> IIER International Conference, Zurich, Switzerland. 26<sup>th</sup> September 2015*, 56-59. <https://doi.org/10.2166/wst.2003.0013>
- Smith, K. A., Ball, T., Connan, F., Dobbie, K. E., Massheder, Rey A (2003). Exchange of greenhouse gases between soil and atmosphere: Interactions of soil physical factors and biological process. *Europeon. Journal of Soil Sci.ence*, *54*, 779-791. <https://doi.org/10.2166/wst.2003.0013>
- Suntoro, J., Syamsiah, J., & Tiyanto, F. A. A. (2013). Potential emission of N<sub>2</sub>O from various types of soil with organic

- material addition. *Jurnal Ilmu Tanah dan Agroklimatologi*, 10(1), 2013. <https://doi.org/10.15608/stjssa.v10i1.133>
- Tani, S. A. A., Bagus, P., Wonny, A. R., Asnath, M. F., & Munif, G. (2017). Integration of Bali Cattle and soybean in Tidal Swamp Land. *Pakistan Journal of Nutrition*, 16(4), 193-199. <https://doi.org/10.3923/pjn.2017.193.199>
- Tracy, M. W., Waren, J., & Almas, B. (2017). Nitrous Oxide production by nitrification and denitrification in soil aggregates as affected by O<sub>2</sub> concentration. Division of Agriculture Science and Natural Resources Oklahoma State University.
- Vallero, D. (2019). Air pollution biodeochemistry. In *Air pollution calculation* (pp. 175-206). <http://dx.doi.org/10.1016/B978-0-12-814934-8.00008-9>
- Wang, T. W., Tu, A. Y., Tsung, L. C., & Juang, K. W. (2017). Effects of biochar amendment and fertilisation on soil properties and maize plant growth. Paper presented in 13<sup>th</sup> International Conference of the East and Southeast Asia Federation of Soil Sciences Societies (13<sup>th</sup> ESAFS), 12-15 December 2017, Nong Nooch Tropical Garden, Pattaya, Pattaya, Thailand. <http://dx.doi.org/10.2136/sssaspecpub63.2014.0040>
- Weitz, A. M., Linder, E., Frohling, S., Crill, P. M., & Keller, M. (2001). N<sub>2</sub>O emissions from humid tropical agricultural soils: Effects of soil moisture, texture and nitrogen availability. *Soil Biology and Biochemistry*, 33(7-8), 1077-1093. [https://doi.org/10.1016/S0038-0717\(01\)00013-X](https://doi.org/10.1016/S0038-0717(01)00013-X)
- Wrage, M. N., Marcus, A. H., Well, R., Muller, C., Velthof, G., & Oenema, O. (2018). Role of nitrifier in the production of nitrous. *Soil Biology and Biochemistry*, 123, A3-A16. <https://doi.org/10.1016/j.soil.bio.2018.03.02>
- Wu, L., Tang, S., He, D., Wu, X., Shaaban, M., Wang, M., Zhao, J., Khan, I., Zheng, X., Hu, R., & Horwath, W. R. (2017). Conversion from rice to vegetable production increases N<sub>2</sub>O emission via increased soil organic matter mineralisation. *Science of the Total Environment*, 583, 190-201. <https://doi.org/10.1016/j.scitotenv.2017.01.050>
- Zhu, N. (2006). *System of rice intensification*. Beijing: China Agricultural Science and Technology Press.