

## GENETIC DIVERSITY AND POPULATION GENETICS OF TROPICAL SHADS (*Tenualosa toli*) FROM SARAWAK, MALAYSIA

NOR AIFFA WAHYU ABU BAKAR<sup>1</sup>, MUHAMMAD SYAFIQ AIMAN MOHD NASIR<sup>1</sup>, TUN NURUL AIMI MAT JAAFAR<sup>1</sup>, FAKHRUDDIN BOKHARI<sup>2</sup>, SARMIZA SAPERI<sup>2</sup> AND NUR ASMA ARIFFIN<sup>1\*</sup>

<sup>1</sup>Faculty of Fisheries and Food Science, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.

<sup>2</sup>Department of Agriculture Sarawak, 12<sup>th</sup>-17<sup>th</sup> Floor, Menara Pelita, Jalan Tun Abdul Rahman Yakub, Petra Jaya, 93050 Kuching, Sarawak, Malaysia.

\*Corresponding author: nurasma@umt.edu.my

<http://doi.org/10.46754/jssm.2025.01.005>

Received: 29 May 2024

Revised: 5 September 2024

Accepted: 15 September 2024

Published: 15 January 2025

**Abstract:** The tropical shads (*Terubok*), *Tenualosa toli* catch has been in decline for the past 15 years owing to the extensive exploitation and high commercial demand. There is limited genetic information on *Terubok* research using a molecular approach, leaving its population's genetic status unassessed. A total of 180 *T. toli* samples from Mukah, Daro, and Pusa in Sarawak, Malaysia were collected for this study. The DNA of samples was then amplified, sequenced, and analysed using Basic Local Alignment Search Tool (BLAST) and Molecular Evolutionary Genetics Analysis (MEGA). The findings indicate that all the samples were *T. toli* with six haplotypes for Cytochrome Oxidase I (*COI*) and 11 haplotypes for Cytochrome B (*CytB*). Neighbour-Joining (NJ) and Maximum Likelihood (ML) phylogenetic trees for the *CytB* gene revealed that all haplotypes formed only a single cluster. The population was discovered to have a low genetic diversity. The ( $F_{ST}$ ) Hierarchical Analysis of Molecular Variance (AMOVA) and pairwise  $\Phi_{ST}$  results indicate no significant genetic differences. The gene flow (Nm) also exhibits the presence of high Nm from the three populations. The Isolation by Distance (IBD) has proven insignificant, indicating that the *T. toli* populations came from similar origins. The neutrality test confirmed that all the samples experienced population expansion and the population is currently in equilibrium. This study provides comprehensive genetic information that can be used for the sustainability management of this valuable species in Sarawak.

Keywords: *Tenualosa toli*, Sarawak, population genetic, genetic diversity, population structure.

### Introduction

Tropical shads, *Tenualosa* spp. locally known as *Terubok* are one of the most economically vital fish species in Malaysia, especially in Sarawak. There are three species of *Tenualosa* recorded in Malaysia: *T. toli*, *T. macrura*, and *T. ilisha* (Abdul Aziz *et al.*, 2015; Arjunaidi *et al.*, 2016; Puvanasundram *et al.*, 2018). *Empirit* is another name that refers to *Terubok*; however, this term is used for small-sized *T. toli* (Khairul Adha *et al.*, 2014; Puvanasundram *et al.*, 2018). *Tenualosa* species is considered an endemic species in Sarawak, inhabiting coastal areas (Blaber *et al.*, 1996; Arjunaidi *et al.*, 2016; Puvanasundram *et al.*, 2018). Notably, *T. toli* belongs to the family Clupeidae, a protandrous hermaphrodite species. *T. toli* is also classified as an anadromous fish and spends most of its juvenile stage in freshwater. Furthermore, *T. toli*

is categorised as a well-adapted fast swimmer among pelagic fish. In addition, this fish has a very thin ventral edge with a long caudal fin and a streamlined body that helps the species survive in waters with high levels of salinity. *T. toli* can be discovered in lagoons and estuarine waters and in shallow muddy water (Blaber *et al.*, 1996; Ambak, 2010). Moreover, this fish is in high demand because of its unique taste and roe (Abdul Aziz *et al.*, 2015; Puvanasundram *et al.*, 2018).

Kasuma *et al.* (2019) highlighted the significance of *Terubok* fish in Sarawak as salted *Terubok* is considered a signature product of the state. The importance of *Tenualosa* spp. to the Sarawak economy can be observed, since some salted *Terubok* fish are claimed to be able to

sell for between RM3,000.00 and RM4,000.00 daily. This is attributable to the local Sarawak *Terubok* fish that can command prices of between RM50.00 to RM70.00 per fish while prices for some of the *Terubok* fish weighing between 800 g and 1 kg can reach RM150.00 for a fish (Mohd Radzi, 2019; Nor Ain, 2021). Furthermore, Mohd Radzi (2019) and Nor Ain (2021) also reported that the roe of the *Tenualosa* spp. sells for between RM200.00 and RM500.00 per kilogram and has high demand from buyers in Singapore and Brunei. This can be observed by referring to the Department of Fisheries Malaysia (DOF) (2021) fish landing statistics, where 2,168 tonnes of *Tenualosa* spp. were landed in the year 2020. The same report also stated that the fishing gear used to capture these fish include gill nets, bag nets, hooks, lines, and trawl nets.

However, there is another species of *Tenualosa* spp. in Sarawak which is *T. macrura*. This shad species has an almost identical morphology to that of *T. toli*, which can lead to misidentification, especially in the juvenile stages (Puvanandram *et al.*, 2018). It was also reported that the *Tenualosa* spp. population has been heavily exploited, including *T. toli* (Blaber *et al.*, 1996; Khairul Adha *et al.*, 2014). Consequently, this situation can affect genetic diversity by reducing the genetic diversity of the population (Kenchington, 2003; Gandra *et al.*, 2021; Taboun *et al.*, 2021).

Various molecular markers have been employed to assess the genetic diversity and populations genetic for both freshwater and marine species of fish (Madduppa *et al.*, 2018; Panprommin *et al.*, 2019; Zhai *et al.*, 2019;

Chen *et al.*, 2020; Firidin *et al.*, 2020; Sun *et al.*, 2021; Prasertlux *et al.*, 2022). Among the markers, mitochondrial genes such as Cytochrome Oxidase I (*COI*) and Cytochrome B (*CytB*) were the genes used for the genetic diversity evaluation for marine fish species (Zheng *et al.*, 2015; Moreira *et al.*, 2019; Liu *et al.*, 2020). *COI* and *CytB* genes were used in the identification of marine fishes since these two genes are capable of identifying unambiguous sequences due to low intraspecific variation and high conservative region possessed by the two genes. Accordingly, it increases the accuracy of species delimitation in fish (Kochzius *et al.*, 2010; Bingpeng *et al.*, 2018; Naeem *et al.*, 2020; Panprommin *et al.*, 2023; Sout *et al.*, 2023).

This study aims to identify the *Tenualosa* species from Daro, Mukah, and Pusa, Sarawak, Malaysia using mitochondrial DNA (mtDNA) *COI* and *CytB* genes. This study also aims to assess the genetic diversity and population structure of the identified *Tenualosa* spp. using the *CytB* gene. The outcome of this study will enable the establishment of *Tenualosa* spp. database from Sarawak waters, in addition to have baseline data on the genetic diversity and population structure of *Tenualosa* spp. in terms of its migratory patterns. This can later be used to conserve the habitat of this highly prized fish species.

## Materials and Methods

### *Samples Collection*

A total of 180 samples of *T. toli* (Figure 1) were collected from three locations of Sarawak River systems (Table 1, Figure 2). These locations



Figure 1: External morphology of (A) *Tenualosa toli* and (B) *Tenualosa macrura*

Table 1: Sampling locations of *T. toli* in the Sarawak River system

River	Sampling Site	Geographical Coordinate	Number of Individuals (n)
Batang Saribas	Pusa	01°37'31.0" N, 111°15'38.4 E	60
Batang Lassa	Daro	02°28'53.0" N, 111°27'07.1 E	60
Gigis River	Mukah	02°54'59.0" N, 112°05'37.6 E	60




Figure 2: Sampling area of *T. toli* in (A) Batang Lassa River, Daro, (B) Gigis River, Mukah, and (C) Batang Saribas River, Pusa

include Mukah (n = 60), Daro (n = 60), and Pusa (n = 60) and the samples were caught using a gill net. All specimens were morphologically identified using taxonomic keys and fish identification references from Whitehead (1985), Munroe *et al.* (1999), Narejo *et al.* (2008), and Hong *et al.* (2013). A small portion of pectoral fins and muscles from each individual was cut, preserved in 95% ethanol, and stored in a 1.5 mL centrifuge tube at 4°C until needed for further analysis.

#### DNA Extraction and Quantification

Genomic material and DNA from the 180 samples was extracted using Cell-free DNA (cfDNA) Extraction Kit (BioTeke Corporation, China) following manufacturer protocol and the DNA was eluted into the final concentration of 70 µL. The concentration and purity of the extracted DNA stock was determined using a Bio Photometer (Eppendorf, Germany) and then stored at -20°C prior to amplification.

#### Polymerase Chain Reaction (PCR) Amplification and Sequencing

Polymerase Chain Reaction (PCR) amplification of *COI* was performed according to Lim *et al.* (2011) while *CytB* according to Faria *et al.* (2006). Polymerase reactions for both genes was prepared in 25.0 µL reaction volumes separately, which contained 1X of 10X PCR buffer with MgCl<sub>2</sub> (Merck Group, Germany), 0.008 mM of Thermo Scientific™ dNTP Mix (Thermo Fisher Scientific, United States), 0.24 mM of forward and reverse primer, respectively, 0.06 mM of Invitrogen Taq DNA Polymerase Recombinant (Thermo Fisher Scientific, United States), 8 ng of DNA template, and sterile deionised water filled up to 25.0 µL. The sequences of the primers used were FishF1 and FishR1 for *COI* (Ward *et al.*, 2005) while Alocytfb1 and Alocytrb1 for *CytB* (Faria *et al.*, 2006). The thermocycling profiles for the *COI* gene consisted of an initial denaturation step of 3 minutes at 94°C, followed by 35 cycles of denaturation step at 94°C for

30 seconds, annealing at 54°C for 40 seconds, and extension at 72°C for 40 seconds, and final extension at 72°C for 10 minutes.

Meanwhile, the thermocycling profiles for the *CytB* gene consisted of an initial denaturation step of 5 minutes at 95°C, followed by 25 cycles of denaturation step at 94°C for 40 seconds, annealing at 54°C for 45 seconds, and extension at 72°C for 1 minute, and final extension at 72°C for 7 minutes. The PCR products were visualised on a 1.2% (w/v) agarose gel stained with SYBR Safe and the clear and single bands of each product proceeded with the purification procedure using QIAquick® PCR Purification Kit (Qiagen, Germany) following the manufacturer's protocol prior to submit it to Apical Scientific Sdn. Bhd. (Selangor, Malaysia) for DNA sequencing.

Samples were sequenced bi-directionally using the BigDye® Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, California, United States). Sequencing reactions were conducted on an ABI 3730xl DNA Analyser (Thermo Fisher Scientific, United States).

#### **Identification of *Tenualosa* spp. using mtDNA *COI* and *CytB***

All *COI* and *CytB* sequences were edited and aligned using the software Chromas (Technelysium Pty Ltd) and BioEdit (Ibis Bioscience). This process was essential to validate the sequences by removing the gaps, noise, and ambiguous sites. Confirmation of species identification was performed by comparing the clean sequences to the sequences in the National Centre for Biotechnology Information (NCBI) database (<http://www.ncbi.nlm.nih.gov>) using the nucleotide Basic Local Alignment Search Tool (BLAST) option to avoid misidentification of specimens collected. The highest percentages of identity, 97% and above were considered as maximum identification and most similar to the sample sequences (Aaron *et al.*, 2018; Mwita & Chuhila, 2023). Expect (E) value for each sample was also provided in the result; the lower the E value or closer to zero,

the more significant the alignment (Ghouri *et al.*, 2020).

#### **Data Analyses**

##### **Genetic Diversity**

The DnaSP software version 5.0.1.1 (Librado & Rozas, 2009) was first used to evaluate the presence of haplotypes in each of the three localities. Representative sequences from each of the haplotypes of *COI* and *CytB* genes were deposited in the GenBank, National Centre for Biotechnology Information (NCBI). GenBank accession number for *COI* haplotypes was OQ380665 – OQ380670 while the haplotypes for *CytB* was OQ421124 – OQ421134. DnaSP was also used to calculate genetic diversity, including analysis levels of DNA polymorphism at each site, including haplotype or gene diversity (h) and nucleotide diversity ( $\pi$ ). Nucleotide diversity is the average pairwise nucleotide differences per site among DNA sequences (Nei, 1987). Estimation of Gene Flow (Nm) was also determined using DnaSP software. Furthermore, DnaSP was again used to evaluate the mismatch distribution estimation of pairwise differences between all individual haplotypes with 10,000 parametric bootstrap iterations to provide a clear picture of previous population demography.

##### **Phylogenetic and Population Level Analyses**

The phylogenetic relationship among haplotypes was determined based on Neighbour-Joining (NJ) and the Maximum Likelihood (ML) methods implemented in MEGA 7.0 (Kumar *et al.*, 2016). Both phylogenetic trees were constructed based on the suggested best-fit DNA substitution models. *CytB* gene analysis was conducted based on the Kimura 2 Parameter with Gamma distribution (K2+G) model with 1,000 bootstrap replications (Kumar *et al.*, 2016). Next, all the *CytB* sequences were aligned with sequences of other *Tenualosa* spp. and some outgroup species available in the GenBank. All the sequences retrieved from the GenBank used for the phylogenetic tree construction were as follows: *T. toli* (KX859103.1), *T. thibaudeaudi* (MG958229.1), *T. ilisha* (MN748961.1), *T.*

*macrura* (KY628753.1), and *Dussumieria elopoides* (MH380250.1).

Population growth was assessed using two selective neutrality tests, Tajima's D (Tajima, 1989) and Fu's  $F_s$  (Fu, 1997) using the software Arlequin ver. 3.5 (Excoffier & Lischer, 2010), with 1,000 permutations under the assumption of neutral evolution in population equilibrium. Hierarchical Analysis of Molecular Variance (AMOVA) was also conducted using the same software. The  $F_{ST}$  value was calculated by incorporating the variance value among groups within the total (FCT), populations within groups (FSC), and populations within the total (FCT). Minimum Spanning Network (MSN) was developed using Arlequin based on the calculation of *CytB* Haplotype Data Files of the three *T. toli* populations generated by DNAsp software. HapStar ver. 0.6 (Teacher & Griffiths, 2010) was used to visualise the network (Elbrecht et al., 2014; Gwak et al., 2014; Osyczka et al., 2014; Williford et al., 2014). Isolation by Distance (IBD) was evaluated using the Mantel Test with 10,000 permutations as implemented in Arlequin ver. 3.5 software using the *CytB* sequence data. The association between geographical distances, measured as linear distance (km) between two sites using Google Earth and genetic distances across all groups was examined (Bohonak, 2002; Winters et al., 2010; Dong et al., 2012; Drew & Barber, 2012; Sá-Pinto et al., 2012; Orsini et al., 2013; Derycke et al., 2013; Li et al., 2013; Wang et al., 2013; Torquato et al., 2019).

## Results and Discussion

### **Identification of *T. toli* using mtDNA *COI* and *CytB***

A total of 141 and 108 individuals from three different populations were successfully sequenced for the partial mtDNA *COI* and *CytB* gene, respectively. The percentage of similarities (BLAST) when compared with sequences from GenBank was 100% and 98.6% for *CytB* and *COI*, respectively, which was above the 97% acceptance percentage. Therefore, this

confirmed that all specimens collected were correctly identified as *T. toli*.

A precise species identification must include molecular analysis, as misidentification potentially occurs due to similarities and overlapping characteristics in the morphology of tropical shads (Arai & Amalina, 2014; Robert et al., 2019). In addition, correct identification is also crucial for the conservation and management of the organisms, as fishes exhibit various patterns of shapes, colours, and sizes (Omer et al., 2017). The samples from Daro, Mukah, and Pusa were confirmed as *T. toli*. The gene fragments *COI* (499 bp) and *CytB* (469 bp) were successfully amplified and sequenced from all specimens extracted in this study. The efficiency of *COI* and *CytB* was proven as a wise marker in species identification and have been extensively used in fish barcoding studies in the Barcode of Life Data System (BOLD Systems) (Kochzius et al., 2010; Ratnasingham & Hebert, 2013; Fernandes et al., 2017; Puvanasundram et al., 2018; Panprommin et al., 2019; Basith et al., 2022; Zafar et al., 2024). It also indicates that at least 27 samples from each population were successfully amplified using *COI* and *CytB* genes. The sampling size of a minimum of  $n = 25$  was also used by several previous genetic population studies such as Hale et al. (2012), Villanueva et al. (2022), and Groeneveld et al. (2024), indicating the results obtained from such a sampling size is considered as valid and acceptable.

Past research has revealed that the taxonomy of Clupeiformes has been extensively studied (Whitehead, 1985). However, the molecular phylogenetic relationships are still poorly understood, indicating that the genetic characteristics such as the genetic species concept had never been tested (Baker & Bradley, 2006; Puvanasundram et al., 2018). In parallel with previous research, findings from this study on the *CytB* gene and *COI* gene reported the genetic relationship between *T. toli* and *T. macrura* (Puvanasundram et al., 2018) and *T. toli* with *T. ilisha* (Arjunaidi et al., 2016). According to Blaber et al. (1999) and Puvanasundram

*et al.* (2018), *T. macrura* closely resembles *T. toli* but has a smaller head and longer tail compared with the latter. Meanwhile, according to Whitehead (1985) and Puvanasundram *et al.* (2018), *T. toli* resembles *T. ilisha*, which has a shorter caudal fin (25% to 31% standard length), longer head (28% to 32%), more gill rakers, and spots along its flank. Therefore, it was suggested that molecular genetic analysis should always be included for precise species identification and confirmation of shad species (Robert *et al.*, 2019).

**Genetic Diversity of *Tenualosa toli* Inferred by mtDNA *CytB***

**Haplotypes Distribution**

A total of 6 and 11 unique haplotypes were identified for *COI* (Table 2) and *CytB* (Table 3), respectively. In *COI* sequence analysis, Hap01 was identified as common haplotype shared by a total of 126 individuals with the highest relative frequency of 0.960. The other five haplotypes (Hap02 to Hap06) demonstrated distinctive unique composite nucleotides

and only represented a single individual with a relatively low frequency of 0.008. Four haplotypes (Hap02, Hap03, Hap 04, and Hap05) were discovered within the Daro locality and another haplotype (Hap06) was recorded within the Mukah locality.

*CytB*, Hap01 is a common haplotype shared by a total of 98 individuals with the highest relative frequency of 0.910. The other 10 haplotypes (Hap02 to Hap11) exhibited unique composite nucleotides. They only represented a single individual with a relatively low frequency of 0.009. Two haplotypes (Hap02 and Hap03) were discovered within the Pusa locality and five haplotypes (Hap04, Hap05, Hap06, Hap07, and Hap08) were detected within the Daro locality. In comparison, another three haplotypes (Hap09, Hap10, and Hap11) were recorded within the Mukah locality.

*CytB* gene has been used widely in many studies on closely related fish species such as *Epinephelus coioides* (Waludin *et al.*, 2018), *Channa striata* (Duong *et al.*, 2019), and *Lates calcarifer* (Bakri & Esa, 2021). In

Table 2: Numbers of haplotype and relative frequency of *COI* of *Tenualosa* spp. from three study sites in Sarawak

Samples			Haplotypes	Frequency
Daro	Mukah	Pusa		
D1, D2, D3, D4, D5, D6, D7, D8, D10, D12, D13, D14, D15, D16, D17, D19, D21, D22, D23, D24, D27, D28, D29, D30, D32, D34, D35, D36, D37, D38, D39, D41, D43, D44, D45, D46, D47	M1, M2, M3, M4, M5, M6, M7, M9, M10, M11, M12, M13, M14, M15, M16, M17, M19, M22, M23, M24, M26, M27, M28, M29, M30, M32, M33, M34, M36, M38, M39, M40, M43, M44, M45, M46, M47, M48, M49, M50, M51, M52, M53, M54, M55, M56, M57, M58, M59	P1, P2, P3, P4, P5, P6, P7, P8, P9, P10, P11, P12, P13, P14, P15, P16, P17, P18, P19, P20, P21, P22, P23, P24, P25, P26, P27, P28, P29, P30, P31, P32, P33, P34, P35, 36, P37, P38, P39, P40, P41, P42, P43, P44, P45, P46, P47, P48, P49, P50	Hap01	0.960
D9			Hap02	0.008
D20			Hap03	0.008
D26			Hap04	0.008
D31			Hap05	0.008
	M25		Hap06	0.008

Table 3: Numbers of haplotype and relative frequency of *CytB* of *Tenualosa* spp. from three study sites in Sarawak

Samples			Haplotypes	Frequency
Daro	Mukah	Pusa		
D1, D2, D3, D4, D5, D6, D7, D8, D9, D11, D12, D13, D14, D15, D16, D17, D18, D19, D22, D23, D24, D25, D26, D29, D30, D31, D32, D33, D34, D35, D36, D37, D38, D39, D40, D41	M2, M3, M4, M5, M6, M7, M9, M10, M11, M13, M14, M15, M16, M17, M18, M19, M20, M23, M24, M25, M26, M27, M28, M30	P1, P2, P3, P4, P5, P6, P7, P8, P9, P10, P11, P12, P13, P14, P15, P16, P17, P18, P19, P20, P21, P22, P23, P24, P25, P26, P27, P28, P29, P30, P31, P33, P34, P36, P37, P38, P39, P40	Hap01	0.910
		P32	Hap02	0.009
		P35	Hap03	0.009
D10			Hap04	0.009
D20			Hap05	0.009
D21			Hap06	0.009
D27			Hap07	0.009
D28			Hap08	0.009
	M1		Hap09	0.009
	M22		Hap10	0.009
	M29		Hap11	0.009

addition, *CytB* has also been considered as the best mitochondrial gene for phylogenetic analysis (Farias *et al.*, 2001; Megarani *et al.*, 2020). *CytB* gene is ideal for examining the systematic diversity of phylogeny due to its slowly evolved codon position and variable domains (Kumazawa & Nishida, 2000; Farias *et al.*, 2001; Shrestha *et al.*, 2019; Megarani *et al.*, 2020; Elvyra *et al.*, 2020). MSN has indicated that the *T. toli* population across Daro, Mukah, and Pusa demonstrated that Hap01

(0.910) was the most dominant and shared by 98 individuals from all three populations. The other 10 haplotypes were unique to each population. There is the presence of unique haplotypes at each of the three localities, which are Daro (n = 5), Mukah (n = 3), and Pusa (n = 2). Daro population had the highest haplotype (0.232) and nucleotide (0.00321) diversities, followed by Mukah at 0.214 and 0.00095, and the Pusa population recorded the lowest value at 0.099 and 0.00021, respectively (Table 4).

Table 4: Genetic diversity for *CytB* gene in *T. toli* showing the number of individuals (n), number of haplotypes (H), haplotype diversity (*h*) and nucleotide diversity ( $\pi$ ), and neutrality tests for each population *Tenualosa* spp. from three study sites in Sarawak

Population	n	H	Genetic Diversity	
			<i>h</i>	$\pi$
Daro	41	6	0.232	0.00321
Mukah	27	4	0.214	0.00095
Pusa	40	3	0.099	0.00021
<b>Total</b>	<b>108</b>	<b>13</b>	-	-

Genetic diversity can greatly influence the species or population's ability to cope and adapt to environmental conditions in new habitats (Frankham *et al.*, 2002; Spielman *et al.*, 2004; Gandra *et al.*, 2021). The value of haplotype diversity ( $h$ ) and nucleotide diversity ( $\pi$ ) are vital indicators for genetic diversity, as a higher the values, the higher the genetic diversity in the populations (Liu *et al.*, 2020; Zhang *et al.*, 2023). The NJ tree is constructed based on the value of the distance of sequences. Meanwhile, the ML tree is constructed based on the maximum values of characters from gene sequences (Amiroch *et al.*, 2018; Hong *et al.*, 2021; Chen & Wang, 2022).

The sharing of haplotype at 91% frequency suggests that the species could probably have originated from a common origin and shared close relationships. The finding are supported by a study by Lee *et al.* (2021), which proposed that the sharing of haplotypes at 81% frequency might come from similar origins. The other 10 haplotypes were unique to each population. There were five, three, and two unique haplotypes of Daro, Mukah, and Pusa, respectively, that were private to single localities. The result of haplotype diversity, ( $h$ ), and nucleotide diversity, ( $\pi$ ) from this study were as follows: Daro ( $h = 0.232$ ,  $\pi = 0.00321$ ), Mukah ( $h = 0.214$ ,  $\pi = 0.00095$ ), and Pusa ( $h = 0.099$ ,  $\pi = 0.00021$ ). These results overall indicated that both haplotype diversity and nucleotide diversity from the three localities have low genetic diversity ( $h < 0.5$ ,  $\pi < 0.005$ ) (Grant & Bowen, 1998; Fang *et al.*, 2022; Trang *et al.*, 2022; Yao *et al.*, 2022). Findings from this study were consistent with previous studies on *Tenualosa* spp. indicated by low values of both haplotype and nucleotide diversity in *T. toli* species at Sarawak rivers, Daro ( $h = 0.232$ ,  $\pi = 0.00032$ ) and Mukah ( $h = 0.178$ ,  $\pi = 0.00097$ ) (Abdul Aziz *et al.*, 2015). The other study on *T. ilisha* species from Indian waters also demonstrated relatively low haplotype and nucleotide diversity in Indian rivers, which were Ganga River ( $h = 0.064$ ,  $\pi = 0.00020$ ) and Hooghly River ( $h = 0.051$ ,  $\pi = 0.0002$ ) (Brahmane *et al.*, 2013). Low values of haplotype and

nucleotide diversity have also been reported in populations of several other different species of fish using the *CytB* gene such as in *Macruronus* species (Olavarria *et al.*, 2006), *Macropodus ocellatus* (Zhang *et al.*, 2020), and *Konosirus punctatus* (Liu *et al.*, 2020). Abdul Aziz *et al.* (2015) also mentioned that the low value of both haplotype and nucleotide diversity might be associated with sharing identical haplotypes (Hap01) among all the samples. As in this study, only 11 haplotypes were detected among 108 individuals with Hap01 being the most common haplotype shared by 98 individuals.

The large number of individuals sharing the same haplotype, Hap01 from the three populations of *COI* and *CytB* genes can indicate there is gene flow between Daro, Mukah, and Pusa. This condition can lead to population homogenisation, leading to many individuals from all three localities sharing the same haplotype (Zhang *et al.*, 2019; Boulanger *et al.*, 2021). The presence of gene flow can be explained by the migratory behaviour of the *T. toli* itself, which is an anadromous fish when it migrates to the river during spawning seasons (Blaber *et al.*, 1996; Roomiani *et al.*, 2014; Puvanasundram *et al.*, 2018; Aiman *et al.*, 2020). This situation can cause the gene flow from the three localities, causing the sharing of a single haplotype among many individuals (Puvanasundram *et al.*, 2018; Baetscher *et al.*, 2019).

Both NJ and ML trees inferred from *CytB* suggested that all 11 haplotypes formed a single cluster. The possible reason for the clustering might be that these 11 haplotypes came from a single population despite the fact that the samples collected were sourced from three different locations. The reasons can be supported by the study conducted by Arai *et al.* (2019) on *T. ilisha*, who discovered that all the haplotypes were grouped into a single clade and deduced that the samples originated from a single population. Abdullah *et al.* (2021; 2022) also reported that the same pattern was discovered as all the samples were grouped into a single clade and mentioned that the individuals possibly came from the same population.

The outcome of the phylogenetic tree, low haplotypes, and nucleotide diversity suggested that the individuals of *T. toli* originated from a single population. However, this population has undergone a bottleneck effect recently, which may have been caused by human activity (Zhang *et al.*, 2020). Moreover, habitat destruction and overfishing might be the two possible reasons for this finding (Chong *et al.*, 2010; Abdul Aziz *et al.*, 2015; Mohd Yusoff *et al.*, 2021). Chong *et al.* (2010) and Hon *et al.* (2016) also reported that there were some peat swamp areas in Sarawak that have been transformed into oil palm plantations, leading to the possibility of habitat destruction of some local species, including *T. toli*. This was since *Tenuulosa* spp. inhabits foreshore areas, estuaries, brackish water, lakes, and freshwater rivers (Roomiani *et al.*, 2014; Hon *et al.*, 2016). In addition, *T. toli* also exhibits migratory behaviour in that it migrates to fresh water sources for spawning but feeds and grows mainly in the sea (Roomiani *et al.*, 2014; Hon *et al.*, 2016). In addition, overfishing events can possibly occur during the spawning migration, which can reduce the number of spawners, leading to low genetic diversity (Abdul Aziz *et al.*, 2015). The impact can reduce the population size, leading to inbreeding and loss of genetic diversity and at worst can result in the extinction of populations (Lakra *et al.*, 2007; Verma *et al.*, 2016).

Phylogenetic trees of NJ and ML were constructed to resolve the relationship amongst 108 *CytB* sequences of *Tenuulosa* species. All haplotypes were clustered in one clade supported by 100% bootstrap value in both trees (Figure 3). The phylogenetic tree revealed that the haplotypes of *T. toli* formed only a single cluster.

#### **Population Structure of *T. toli* Inferred by mtDNA *CytB***

An AMOVA of *T. toli* was almost perfect within the population variations (100.02%) and low variations were detected among the populations (-0.02%). There were no variations among groups due to the lack of populations used (only

three). All hierarchical group analyses were insignificant at  $p > 0.05$  (Table 5).

Pairwise  $\Phi_{ST}$  values between the three populations ranged from -0.00599 between Daro and Mukah to 0.00877 between Pusa and Mukah. According to Chanthran *et al.* (2020), pairwise  $\Phi_{ST}$  values are often used to infer gene flow, in which a lower  $\Phi_{ST}$  value indicates low genetic divergence and higher gene flow. The highest gene flow was between Mukah and Daro (40.93) and the lowest was between Pusa and Mukah (25.28). The average gene flow for all three locations was 26.47.

IBD, based on the Mantel test, discovered that there was an insignificant correlation between genetic distance (pairwise  $\Phi_{ST}$  values) and geographic distance (km) (Figure 4). This indicates that the genetic distance (pairwise  $\Phi_{ST}$  values) was not influenced by geographic distance (km). Overall, the findings on genetic structure indicated panmixia with high gene flow among all three populations of *T. toli* in Sarawak Rivers (Daro, Mukah, and Paso).

The genetic structure of all three populations (Daro, Mukah, and Pusa) was examined by several methods, which were AMOVA F-statistics ( $F_{ST}$ ), pairwise  $\Phi_{ST}$  values, gene flow (Nm), IBD, and MSN. The amount and distribution of genetic variation within and between populations is defined as genetic structure (Pastorino & Marchelli, 2022). Efficiently identifying the genetic structure and related factors of a species can lead to an understanding of the evolutionary history of the species (Ju *et al.*, 2019).

The AMOVA  $F_{ST}$  result suggested no significant genetic differentiation ( $F_{ST} = 0.42620$ ,  $p > 0.05$ ) between all three populations. Pairwise  $\Phi_{ST}$  values further confirmed no significant genetic differentiation for each population. This condition was suspected due to the presence of gene flow (Yin *et al.*, 2013; Parmaksidz & Ekma, 2017; Sarker *et al.*, 2021). Gene flow analysis proved that there was high gene flow between the three locations. Cheng *et al.* (2020) mentioned that the values of gene flow ( $Nm < 1$ )




Figure 3: Phylogenetic analysis of *CytB* haplotypes with *T. toli*, *T. thibaudeaui*, *T. macrura*, and *T. ilisha* as sister group and *Dussumieria elopsoides* as outgroup based on (A) Neighbour-Joining (NJ) and (B) Maximum Likelihood (ML)

Table 5: Pairwise  $\Phi_{ST}$  (below diagonal) and gene flow (Nm) (upper diagonal) between populations of *T. toli* and AMOVA and F-statistics

Populations	Pairwise $\Phi_{ST}$ and Gene Flow			AMOVA and F-statistics			
	Mukah	Daro	Pusa	Variance Components	% Variations	F-statistics	p-value
Mukah	-	40.93	25.28	Among populations	-0.02	-0.00023	0.42620
Daro	-0.00599	-	34.94	Within populations	100.02		
Pusa	0.00877	0.00321	-				

Note: All pairwise comparisons are not significant at  $p > 0.05$ ; all  $p$ -values are not significant at  $p > 0.05$ .




Figure 4: *T. toli* population structure analysis: (A) Isolation by Distance (IBD) correlation between the genetic and geographic distance of *T. toli*. All pairwise comparisons are not significant at  $p < 0.05$ ; (B) Minimum Spanning Network (MSN) of *T. toli* across Daro, Mukah, and Pusa. The size of each circle indicates the frequency of the corresponding haplotype in the whole dataset

is interpreted as low to no Nm. The high Nm from this study was consistent with a previous study that indicated high to infinity Nm on *T. ilisha* between the Ganga and Hooghly rivers in India and in populations of *Rhabdosargus sarba* from Taiwan. This suggests that the free flow of the gene pool and the populations should be considered as one homogenous population (Brahmane et al., 2013; Hsu et al., 2020).

Another study by Cheng et al. (2020) revealed similar results as the Nm, valued between 9.00 and 24.00, indicating a high Nm rate. Previous research highlighted that the reason that led to the absence of genetic structure and population differentiation in the *T. toli* population was the migration behaviour of the *Tenuulosa* spp. increasing the chance for *T. toli* to migrate and spawn at other Sarawak rivers (Blaber et al., 1996; Roomiani et al., 2014; Abdul Aziz et al., 2015; Aiman et al., 2020). This is attributable to the fish's natural behaviour, leading to local population clusters or a mixture with other populations (Alvarado et al., 2005; Sarker et al., 2021).

In addition, the presence of high Nm between the three locations supported the findings of sharing the single haplotype by many individuals between the three populations above. IBD results from this study were consistent with the above findings, as there was no significant correlation between

the genetic distance (pairwise  $\Phi_{ST}$ ) and the geographic distance (km). The clustering of the MSN also indicates that the juvenile *T. toli* originated from similar origins and shared close relationships among the individuals, explaining the non-significant pairwise value  $\Phi_{ST}$  and high gene flow rate. Similar results were obtained by Yu et al. (2016) and Nazia et al. (2010), where the *Hyporhamphus sajori* and *Clarias macrocephalus*, respectively, came from a single population due to the absence of obvious philographic structure of the MSN.

#### Demographic History of *T. toli*

Two neutrality tests, Tajima's D and Fu's test revealed that each population achieved the following values: Daro (-2.62384; -0.03533), Mukah (-2.06816; -1.56486), and Pusa (-1.48662; -2.72950), respectively. Tajima's D values demonstrated a significant negative for all populations at  $p < 0.05$ . Meanwhile, in Fu's test, only Daro was insignificant at  $p < 0.05$  (Table 6). Significant negative values of Tajima's D and Fu's Fs would indicate a significant population expansion (Tajima, 1989; Fu, 1997).

Population mismatch distribution estimation of pairwise differences revealed that the population has a multimodal distribution pattern, as in Figure 5. A unimodal distribution pattern indicates that the population has undergone recent expansion while a bimodal and

Table 6: Neutrality tests for *T. toli* populations from Daro, Mukah, and Pusa

Population	n	H	Neutrality's Test	
			Tajima's D	Fu's Fs
Daro	41	6	<b>-2.62384</b>	-0.03533
Mukah	43	4	<b>-2.06816</b>	<b>-1.56486</b>
Pusa	35	3	<b>-1.48662</b>	<b>-2.72950</b>
<b>Total</b>	<b>119</b>	<b>13</b>	-	-

Note: Bold indicated significant at  $p < 0.05$ .




Figure 5: Multimodal mismatch distributions (pairwise differences) of *T. toli* showing the expected and observed pairwise differences between sequences with the respective frequencies

multimodal distribution pattern indicates that a population is in an equilibrium state (Rogers & Harpending, 1992; Jenkins *et al.*, 2018).

In addition, the MSN had a star-like shape. The finding further supports the occurrence of a recent demographic expansion in the population.

Tajima's negative D values suggest a balanced selection and the recent directional selection of the population has increased with an excess of uncommon alleles (Tajima, 1989). Based on Ewens' sampling distribution, Fu's  $F_{ST}$  is a sophisticated test that leverages information on haplotype distribution to detect previous population size fluctuations (Ewens, 1972;

Ramos-Onsins & Rozas, 2002). Under an excess of mutations, a significant negative value of Fu's  $F_s$  statistics is considered proof of population expansion.

Consistent with haplotype and nucleotide diversity, Tajima's D values were significantly negative for all populations at  $p < 0.05$ . Meanwhile, in Fu's test, only Daro was insignificant at  $p < 0.05$ . Significant negative values of Tajima D and Fu's  $F_s$  would indicate a rapid, significant population expansion in the past (Tajima, 1989; Fu, 1997; Gui *et al.*, 2021). However, for Daro, since this site has a significant negative value for Tajima's D but

an insignificant negative for  $F_u$ 's  $F_s$  value, the results reveal that the population of *T. toli* in this area has not expanded recently. However, the expansion that was experienced before was probably caused by the bottleneck effect that leads to a small effective early population size (Kashani et al., 2021). Furthermore, the results were supported by population demographic mismatch distribution analysis of *T. toli* as it matched the multimodal distribution.

Kai et al. (2014) and Shi et al. (2014) stated that a population with multimodal distribution indicates that the population is demographically equilibrium. In addition, the result from the MSN demonstrates the presence of a star-like shape structure. The star-like shape in the MSN suggests that there is an occurrence of significant population expansion due to the bottleneck effect (Yu et al., 2016; Landvik et al., 2017; Supmee et al., 2020; Zhou et al., 2023). This bottleneck effect might be caused by habitat loss due to the changing sea temperature, current, and coastline morphology during the late Pleistocene (the past 2.5 million years ago) (Tschá et al., 2016; Yu et al., 2016; Divya et al., 2017).

## Conclusions

In conclusion, this study revealed that all the samples taken from the three localities were confirmed as *T. toli* using *COI* and *CytB* genes. The genetic diversity of *T. toli* in Sarawak is relatively low and consists of one homogeneous population from three different population. This suggested the species could probably have originated from a common origin and shared a close relationship. Genetic structure analysis indicates non-significant genetic differentiation and pairwise  $\Phi_{ST}$  with high gene flow. IBD also exhibits a non-significant correlation between the pairwise  $\Phi_{ST}$  and the geographical distances (km). MSN had similar results as ML and NJ phylogenetic trees, which is the *T. toli* came from a similar origin. The neutrality test confirmed that all the samples have experienced population expansion and the population is

currently in an equilibrium state, probably due to bottleneck as a result of habitat loss during the late Pleistocene period.

It is suggested that collecting research data on genetic diversity and population dynamics is crucial and could serve as a monitoring tool for the sustainable management of resources. The genetic pools for a sustainable harvest must be preserved and replenished, which requires intensive management of fishery resources. Collecting more samples from different regions would make it possible to study the population genetics of a migratory species dispersed over regional boundaries such as the *T. toli*, if we can start working together, primarily with the related agencies like the DOF and Department of Agriculture Sarawak (DOA). A comprehensive strategy uses a wide range of complementary methods and more sensitive and adaptable markers like microsatellites for more conclusive results. Various programmes can be implemented such as a comprehensive conservation plan, fisheries management programmes, and foundation for sustainable fishery activities, together with other related information.

## Acknowledgements

The author would like to thank the Faculty of Fisheries and Food Science and Universiti Malaysia Terengganu for providing the research facilities. The authors would also like to express their gratitude to the Department of Agriculture, Sarawak for assisting in the collection of samples. Thank you also to Dr. Shahreza Md. Sheriff, project leader for Fundamental Research Grant Scheme (FRGS) Vot no. 59245 and the Ministry of Higher Education Malaysia for funding the research. The authors would like to thank Prof. Dr. Muhammad Ikhwanuddin Abdullah for his valuable comments on the article. Last but not least, we are deeply grateful and would like to appreciate the late Mr. Awang Alim Awang Kassim for his valuable contributions to the success of this project.

### Conflict of Interest Statement

The authors declare that they have no conflict of interest.

### References

- Aaron, C. C., Aziz, A. H. A., Meriam, S. A. S. T., Seah, Y. G., & Asma, A. N. (2018). Morphological and molecular identification of mullet species (Mugilidae) from Setiu Wetland, Terengganu, Malaysia. *Aquaculture, Aquarium, Conservation & Legislation*, 11(2), 429-438.
- Abdul Kadar, N., Mustafa, S., Hamzah, A., Nguyen, T. T. T., & Siti Azizah, M. N. (2010). No genetic differentiation between geographically isolated populations of *Clarias macrocephalus* (Gunther) in Malaysia revealed by sequences of mtDNA cytochrome B and D-loop gene regions. *Journal of Applied Ichthyology*, 26, 568-570.
- Abdullah, T. A., Al-Noor, S. S., & Javadmanesh, A. (2021). Genetic distance of Anadromous *Tenualosa ilisha* (Hamilton, 1882) in Shatt al-Arab River based on partial sequencing of mitochondrial DNA. *Agriculture and Natural Resources*, 55(5), 849-854. <https://doi.org/10.34044/j.anres.2021.55.5.15>
- Abdullah, T. A., Javadmanesh, A., & Al-Noor, S. S. (2022). Genetic indications for anadromous Hilsa Shad (*Tenualosa ilisha*) in Shatt al-Arab river using mtDNA Cytochrome B gene. *Basrah Journal of Agricultural Sciences*, 35(1), 71-80. <https://doi.org/10.37077/25200860.2022.35.1.06>
- Abdul Aziz, A. H., Tarmizi, Z., Ali, N. M., Ariffin, N. A., Abdullah, M. D., Lian, W. L., Giat, S. Y., Jaafar, T. N., Bolong, A. M., Kasim, A. A., & Sheriff, S. M. (2015). Mitochondrial DNA diversity of *Terubok* (*Tenualosa toli*) from Daro and Mukah, Sarawak inferred by partial Cytochrome B (Cyt-B). *Journal of Fisheries and Aquatic Science*, 10(2), 92-101. <https://doi.org/10.3923/jfas.2015.92.101>
- Aiman, W. M., Yusoff, F. D., Arshad, A., Kamal, A. H., Ismail, J., Idris, M. H., Karim, N. U., & Asif, A. A. (2020). Distribution of zooplankton community in Toli shad (*Tenualosa toli*) habitats, Sarawak, Malaysia. *Biodiversitas Journal of Biological Diversity*, 21(9). <https://doi.org/10.13057/biodiv/d210913>
- Alvarado Bremer, J. R., Mejuto, J., Gómez-Márquez, J., Boán, F., Carpintero, P., Rodríguez, J. M., Viñas, J., Greig, T. W., & Ely, B. (2005). Hierarchical analyses of genetic variation of samples from breeding and feeding grounds confirm the genetic partitioning of northwest Atlantic and South Atlantic populations of swordfish (*Xiphias gladius* L.). *Journal of Experimental Marine Biology and Ecology*, 327(2), 167-182. <https://doi.org/10.1016/j.jembe.2005.06.022>
- Ambak, M. A. (2012). *Fishes of Malaysia*. Terengganu: Penerbit UMT, Universiti Malaysia Terengganu.
- Amiroch, S., Pradana, M. S., Irawan, M. I., & Mukhlash, I. (2018). Maximum likelihood method on the construction of phylogenetic tree for identification the spreading of SARS epidemic. *2018 International Symposium on Advanced Intelligent Informatics (SAIN)*. <https://doi.org/10.1109/sain.2018.8673334>
- Arai, T., & Amalina, R. (2014). New record of a tropical Shad *Tenualosa ilisha* (Teleostei: Clupeidae) in Malaysian Waters. *Marine Biodiversity Records*, 7, 1-4. <https://doi.org/10.1017/s1755267214000736>
- Arai, T., Taha, H., Amalina, R., Iizuka, Y., & Chang, C. W. (2019). Anadromy and heterogenous population of a tropical shad *Tenualosa ilisha* in Malaysia, as revealed by otolith microchemistry and molecular evidence. *Journal of Fish Biology*, 95(6), 1506-1511. <https://doi.org/10.1111/jfb.14154>

- Arjunaidi, N. N., Zakaria, M. F., Aziz, A. H. A., Sheriff, M. S., Jaafar, T. N. A., Seah, Y. G., & Ariffin, N. A. (2016). Authentication of *Tenualosa* species in Perak River, Malaysia: Application of morphological measurement and molecular analysis of partial *COI* and 16S genes to resolve species ambiguity. *AACL Bioflux*, 9(6), 1355-1363.
- Baetscher, D. S., Anderson, E. C., Gilbert-Horvath, E. A., Malone, D. P., Saarman, E. T., Carr, M. H., & Garza, J. C. (2019). Dispersal of a nearshore marine fish connects marine reserves and adjacent fished areas along an open coast. *Molecular Ecology*, 28(7), 1611-1623. <https://doi.org/10.1111/mec.15044>
- Baker, R. J., & Bradley, R. D. (2006). Speciation in mammals and the genetic species concept. *Journal of Mammalogy*, 87(4), 643-662. <https://doi.org/10.1644/06-mammf-038r2.1>
- Bakri, A. M., & Esa, Y. (2021). Analysis of genetic diversity in five captive population of Asian Seabass (*Lates calcarifer*) for selective breeding in Malaysia. In F. Abu Bakar, M. F. Mat Isa., K. S. Rajoo., S. D. Ramiya., M. I. Asmuni., & S. S. Leong (Eds.), *1<sup>st</sup> Postgraduate Seminar on Agriculture and Forestry 2021 (PSAF 2021), 6 October 2021* (pp. 71-75). [https://www.researchgate.net/publication/356811193\\_Analysis\\_of\\_Genetic\\_Diversity\\_in\\_5\\_Captive\\_Population\\_of\\_Asian\\_Seabass\\_Lates\\_calcarifer\\_for\\_Selective\\_Breeding\\_in\\_Malaysia](https://www.researchgate.net/publication/356811193_Analysis_of_Genetic_Diversity_in_5_Captive_Population_of_Asian_Seabass_Lates_calcarifer_for_Selective_Breeding_in_Malaysia)
- Basith, A., Abinawanto, Kusriani, E., & Yasman. (2022). DNA barcode characterization of chocolate hind grouper (*Cephalopholis Boenak*) in several Indonesia waters with the new sequences record from Madura Island. *HAYATI Journal of Biosciences*, 29(6), 733-741. <https://doi.org/10.4308/hjb.29.6.733-741>
- Bingpeng, X., Heshan, L., Zhilan, Z., Chunguang, W., Yanguo, W., & Jianjun, W. (2018). DNA barcoding for identification of fish species in the Taiwan Strait. *PLOS ONE*, 13(6), e0198109. <https://doi.org/10.1371/journal.pone.0198109>
- Blaber, S. J., Milton, D. A., Pang, J., Wong, P., Boon-Teck, O., Nyigo, L., & Lubim, D. (1996). The life history of the tropical Shad *Tenualosa toli* from Sarawak: First evidence of Protandry in the Clupeiformes? *Environmental Biology of Fishes*, 46(3), 225-242. <https://doi.org/10.1007/bf00004998>
- Blaber, S. J. M., Brewer, D. T., Milton, D. A., Merta, G. S., Efizon, D., Fry, G., & van der Velde, T. (1999). The life history of the Protandrous Tropical Shad *Tenualosa macrura* (Alosinae: Clupeidae): Fishery implications. *Estuarine, Coastal and Shelf Science*, 49(5), 689-701. <https://doi.org/10.1006/ecss.1999.0545>
- Bohonak, A. J. (2002). IBD (Isolation by Distance): A program for analyses of isolation by distance. *Journal of Heredity*, 93(2), 153-154. <https://doi.org/10.1093/jhered/93.2.153>
- Boulanger, E., Benestan, L., Guerin, P., Dalongeville, A., Mouillot, D., & Manel, S. (2021). Climate differently influences the genomic patterns of two sympatric marine fish species. *Journal of Animal Ecology*, 91(6), 1180-1195. <https://doi.org/10.1111/1365-2656.13623>
- Brahmane, M. P., Kundu, S. N., Das, M. K., & Sharma, A. P. (2013). Low genetic diversity and absence of population differentiation of hilsa (*Tenualosa ilisha*) revealed by mitochondrial DNA Cytochrome B region in Ganga and Hooghly rivers. *African Journal of Biotechnology*, 12(22), 3383-3389. <https://doi.org/https://doi.org/10.5897/AJB2013.12364>
- Chantran, S. S., Lim, P.-E., Li, Y., Liao, T.-Y., Poong, S.-W., Du, J., Hussein, M. A., Sade, A., Rumpet, R., & Loh, K.-H. (2020). Genetic diversity and population structure of *Terapon jarbua* (Forskål, 1775) (Teleostei, Terapontidae) in Malaysian

- Waters. *ZooKeys*, 911, 139-160. <https://doi.org/10.3897/zookeys.911.39222>
- Chen, W., & Wang, H. (2022). Phylogenetic tree selection by testing substitution number in clade. *Diversity*, 14(7), 543. <https://doi.org/10.3390/d14070543>
- Chen, W., Li, C., Chen, F., Li, Y., Yang, J., Li, J., & Li, X. (2020). Phylogeographic analyses of a migratory freshwater fish (*Megalobrama terminalis*) reveal a shallow genetic structure and pronounced effects of sea-level changes. *Gene*, 737, 144478. <https://doi.org/10.1016/j.gene.2020.144478>
- Cheng, J., Kao, H., & Dong, S. (2020). Population genetic structure and gene flow of rare and endangered *Tetraena mongolica* Maxim revealed by reduced representation sequencing. *BMC Plant Biology*, 20(391), 1-13. <https://doi.org/10.21203/rs.3.rs-19749/v2>
- Chong, V. C., Lee, P. K., & Lau, C. M. (2010). Diversity, extinction risk and conservation of Malaysian fishes. *Journal of Fish Biology*, 76(9), 2009-2066. <https://doi.org/10.1111/j.1095-8649.2010.02685.x>
- Department of Fisheries Malaysia. (2021). Annual Fisheries Statistics 2020. Department of Fisheries, Ministry of Agriculture and Food Industry. Accessed November 20, 2022, from <https://www.dof.gov.my/en/resources/fisheries-statistics-i/>
- Derycke, S., Backeljau, T., & Moens, T. (2013). Dispersal and gene flow in free-living marine nematodes. *Frontiers in Zoology*, 10(1), 1. <https://doi.org/10.1186/1742-9994-10-1>
- Divya, P. R., Linu, J., Mohitha, C., Kathirvelpandian, A., Manoj, P., Basheer, V. S., & Gopalakrishnan, A. (2017). Deciphering demographic history and fine-scale population structure of Cobia, *Rachycentron canadum* (Pisces: Rachycentridae) using microsatellite and mitochondrial markers. *Marine Biodiversity*, 49(1), 381-393. <https://doi.org/10.1007/s12526-017-0817-x>
- Dong, Y.-wei, Wang, H.-shan, Han, G.-D., Ke, C.-huan, Zhan, X., Nakano, T., & Williams, G. A. (2012). The impact of Yangtze River discharge, ocean currents and historical events on the biogeographic pattern of *Cellana toreuma* along the China Coast. *PLOS ONE*, 7(4). <https://doi.org/10.1371/journal.pone.0036178>
- Drew, J. A., & Barber, P. H. (2012). Comparative phylogeography in Fijian coral reef fishes: A multi-taxa approach towards marine reserve design. *PLOS ONE*, 7(10). <https://doi.org/10.1371/journal.pone.0047710>
- Duong, T.-Y., Uy, S., Chheng, P., So, N., Tran, T.-H. T., Nguyen, N.-T. T., Pomeroy, R., & Egna, H. (2019). Genetic diversity and structure of striped snakehead (*Channa striata*) in the Lower Mekong Basin: Implications for aquaculture and fisheries management. *Fisheries Research*, 218, 166-173. <https://doi.org/10.1016/j.fishres.2019.05.014>
- Elbrecht, V., Feld, C. K., Gies, M., Hering, D., Sondermann, M., Tollrian, R., & Leese, F. (2014). Genetic diversity and dispersal potential of the stonefly *Dinocras cephalotes* in a Central European low mountain range. *Freshwater Science*, 33(1), 181-192. <https://doi.org/10.1086/674536>
- Elvyra, R., Solihin, D. D., Affandi, R., Junior, M. Z., & Suhendra, M. (2020). Short communication: Molecular characteristics and phylogenetic relationships of silurid catfishes (*Kryptopterus*, *ompok* and *phalacronotus*) from the Kampar River, Indonesia, based on the Cytochrome B gene. *Biodiversitas Journal of Biological Diversity*, 21(8), 3539-3546. <https://doi.org/10.13057/biodiv/d210816>
- Ewens, W. J. (1972). The sampling theory of selectively neutral alleles. *Theoretical Population Biology*, 3(1), 87-112. [https://doi.org/10.1016/0040-5809\(72\)90035-4](https://doi.org/10.1016/0040-5809(72)90035-4)
- Excoffier, L., & Lischer, H. E. (2010). Arlequin Suite Ver 3.5: A new series of programs to perform population genetics analyses under

- linux and windows. *Molecular Ecology Resources*, 10(3), 564-567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Fang, D.-an, Luo, H., He, M., Mao, C., Kuang, Z., Qi, H., Xu, D., Tan, L., & Li, Y. (2022). Genetic diversity and population differentiation of naked carp (*Gymnocypris przewalskii*) revealed by Cytochrome Oxidase subunit I and D-loop. *Frontiers in Ecology and Evolution*, 10. <https://doi.org/10.3389/fevo.2022.827654>
- Faria, R., Weiss, S., & Alexandrino, P. (2006). A molecular phylogenetic perspective on the evolutionary history of *Alosa* spp. (Clupeidae). *Molecular Phylogenetics and Evolution*, 40(1), 298-304. <https://doi.org/10.1016/j.ympev.2006.02.008>
- Farias, I. P., Ortí, G., Sampaio, I., Schneider, H., & Meyer, A. (2001). The Cytochrome B gene as a phylogenetic marker: The limits of resolution for analyzing relationships among cichlid fishes. *Journal of Molecular Evolution*, 53(2), 89-103. <https://doi.org/10.1007/s002390010197>
- Fernandes, T. J. R., Costa, J., Oliveira, M. B., & Mafra, I. (2017). DNA barcoding coupled to HRM analysis as a new and simple tool for the authentication of Gadidae fish species. *Food Chemistry*, 230, 49-57. <https://doi.org/10.1016/j.foodchem.2017.03.015>
- Firidin, S., Ozturk, R. C., Alemdag, M., Eroglu, O., Terzi, Y., Kutlu, I., Duzgunes, Z. D., Cakmak, E., & Aydin, I. (2020). Population genetic structure of Turbot (*Scophthalmus maximus* L., 1758) in the Black Sea. *Journal of Fish Biology*, 97(4), 1154-1164. <https://doi.org/10.1111/jfb.14487>
- Frankham, R., Briscoe, D. A., & Ballou, J. D. (2002). *Introduction to conservation genetics*. New York, USA: Cambridge University Press.
- Fu, Y.-X. (1997). Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics*, 147(2), 915-925. <https://doi.org/10.1093/genetics/147.2.915>
- Gandra, M., Assis, J., Martins, M. R., & Abecasis, D. (2021). Reduced global genetic differentiation of exploited marine fish species. *Molecular Biology and Evolution*, 38(4), 1402-1412. <https://doi.org/10.1093/molbev/msaa299>
- Ghouri, M. Z., Ismail, M., Javed, M. A., Khan, S. H., Munawar, N., Umar, A. B., Mehr-un-Nisa, Aftab, S. O., Amin, S., Khan, Z., & Ahmad, A. (2020). Identification of edible fish species of Pakistan through DNA barcoding. *Frontiers in Marine Science*, 7. <https://doi.org/10.3389/fmars.2020.554183>
- Grant, W. S., & Bowen, B. W. (1998). Shallow population histories in deep evolutionary lineages of marine fishes: Insights from sardines and anchovies and lessons for conservation. *Journal of Heredity*, 89(5), 415-426. <https://doi.org/10.1093/jhered/89.5.415>
- Groeneveld, M., Klein, J., Bennett, R., Abdulla, A., Bond, M., Ebert, D., Fernando, S., Gledhill, K., Jaquemet, S., Kiszka, J., Macdonald, A., Mann, B., Nevill, J., Price, A., Rumbelow, J., Siteo, J., van Staden, M., Wueringer, B., & Wueringer, B. (2024). Population genetic structure of bottlenose and whitespotted wedgefishes from the Southwest Indian Ocean using a dual marker approach. *Endangered Species Research*, 53, 409-427. <https://doi.org/10.3354/esr01311>
- Gui, Z., Wu, L., Cai, H., Mu, L., Yu, J.-F., Fu, S.-Y., & Si, X.-Y. (2021). Genetic diversity analysis of *Dermacentor nuttalli* within Inner Mongolia, China. *Parasites & Vectors*, 14(1). <https://doi.org/10.1186/s13071-021-04625-5>
- Gwak, W.-S., Lee, Y.-D., & Nakayama, K. (2014). Population structure and sequence divergence in the mitochondrial DNA control region of Gizzard Shad *Konosirus punctatus* in Korea and Japan. *Ichthyological Research*, 62(3), 379-385. <https://doi.org/10.1007/s10228-014-0450-7>

- Hale, M. L., Burg, T. M., & Steeves, T. E. (2012). Sampling for microsatellite-based population genetic studies: 25 to 30 individuals per population is enough to accurately estimate allele frequencies. *PLOS ONE*, 7(9), e45170. <https://doi.org/10.1371/journal.pone.0045170>
- Hon, J., Mohd-Azlan, J., Apui, S. A., & Lip, B. (2016). Not wastelands, let's manage our peat swamp properly through a systemic conservation. *15<sup>th</sup> International Peat Congress 2016*, 546-550.
- Hong, X., Zhu, X., Chen, K., Pan, D., & Li, K. (2013). Morphometric analysis of *Tenualosa ilisha*, *Alosa sapidissima* and *Tenualosa reevesii*. *Journal of South China Agricultural University*, 34(2), 203-206.
- Hong, Y., Guo, M., & Wang, J. (2021). ENJ algorithm can construct triple phylogenetic trees. *Molecular Therapy - Nucleic Acids*, 23, 286-293. <https://doi.org/10.1016/j.omtn.2020.11.004>
- Hsu, T.-H., Huang, C.-W., Lee, H.-T., Kuo, Y.-H., Liu, K.-M., Lin, C.-H., & Gong, H.-Y. (2020). Population genetic analysis for stock enhancement of silver sea bream (*Rhabdosargus sarba*) in Taiwan. *Fishes*, 5(2), 19. <https://doi.org/10.3390/fishes5020019>
- Jenkins, T. L., Castilho, R., & Stevens, J. R. (2018). Meta-analysis of northeast Atlantic marine taxa shows contrasting phylogeographic patterns following post-LGM expansions. *PeerJ*, 6. <https://doi.org/10.7717/peerj.5684>
- Ju, M.-M., Feng, L., Yang, J., Yang, Y.-C., Chen, X.-D., & Zhao, G.-F. (2019). Evaluating population genetic structure and demographic history of *Quercus spinosa* (Fagaceae) based on specific length amplified fragment sequencing. *Frontiers in Genetics*, 10, 1-12. <https://doi.org/10.3389/fgene.2019.00965>
- Kai, Y., Ueda, Y., Fujiwara, K., Itoh, M., Yamasaki, A., & Nakabo, T. (2014). Population structure and demographic history of *Davidijordania poecilimon* (Perciformes: Zoarcidae). *Species Diversity*, 19(1), 15-19. <https://doi.org/10.12782/sd.19.1.015>
- Kashani, E., Rezaei, H. R., Naderi, M., & Khorasani, N. (2021). Genetic diversity, population structure and demographic history of Brandt and long eared hedgehogs in Iran. *Pakistan Journal of Zoology*, 53(2). <https://doi.org/10.17582/journal.pjz/20190803220849>
- Kasuma, J., Farhana, N. N., Kamaruddin, H. M., Rusuli, M. S., & Yacob, Y. (2019). Identifying the challenges of the Sarawak malay *Terubok Ikan Masin* (salted fish) entrepreneur: Qualitative study. *Research in World Economy*, 10(2), 30. <https://doi.org/10.5430/rwe.v10n2p30>
- Kenchington, E., Heino, M., & Nielsen, E. E. (2003). Managing marine genetic diversity: Time for action? *ICES Journal of Marine Science*, 60(6), 1172-1176. [https://doi.org/10.1016/s1054-3139\(03\)00136-x](https://doi.org/10.1016/s1054-3139(03)00136-x)
- Khairul Adha, A. R., Hambali, T. M., Shabdin, M. L., Esa, Y., & Husaini, A. S. A. (2014). The status of *Tenualosa toli* (Valenciennes, 1847) in the Southwest Coast of Sarawak, Malaysia. *Kuroshio Science*, 8(1), 39-44.
- Kochzius, M., Seidel, C., Antoniou, A., Botla, S. K., Campo, D., Cariani, A., Vazquez, E. G., Hauschild, J., Hervet, C., Hjörleifsdottir, S., Hreggvidsson, G., Kappel, K., Landi, M., Magoulas, A., Marteinson, V., Nölte, M., Planes, S., Tinti, F., Turan, C., Venugopal, M. N., Weber, H., & Blohm, D. (2010). Identifying fishes through DNA barcodes and microarrays. *PLOS ONE*, 5(9). <https://doi.org/10.1371/journal.pone.0012620>
- Kumar, S., Stecher, G., & Tamura, K. (2016). Mega7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874. <https://doi.org/10.1093/molbev/msw054>

- Kumazawa, Y., & Nishida, M. (2000). Molecular phylogeny of Osteoglossoids: A new model for Gondwanian origin and plate tectonic transportation of the Asian arowana. *Molecular Biology and Evolution*, 17(12), 1869-1878. <https://doi.org/10.1093/oxfordjournals.molbev.a026288>
- Lakra, W. S., Mohindra, V., & Lal, K. K. (2007). Fish genetics and conservation research in India: Status and perspectives. *Fish Physiology and Biochemistry*, 33(4), 475-487. <https://doi.org/10.1007/s10695-007-9168-z>
- Landvik, M., Miraldo, A., Niemelä, P., Valainis, U., Cibulskis, R., & Roslin, T. (2017). Evidence for geographic substructuring of mtDNA variation in the east European hermit beetle (*Osmoderma barnabita*). *Nature Conservation*, 19, 171-189. <https://doi.org/10.3897/natureconservation.19.12877>
- Lee, K. H., Jeong, J. S., Park, J. S., Kim, M. J., Jeong, N. R., Jeong, S. Y., Lee, G. S., Lee, W., & Kim, I. (2021). Tracing the invasion and expansion characteristics of the flatid planthopper, *Metcalfa pruinosa* (Hemiptera: Flatidae), in Korea using mitochondrial DNA sequences. *Insects*, 12(1), 4. <https://doi.org/10.3390/insects12010004>
- Li, L., Wu, X., & Yu, Z. (2013). Genetic diversity and substantial population differentiation in *Crassostrea hongkongensis* revealed by mitochondrial DNA. *Marine Genomics*, 11, 31-37. <https://doi.org/10.1016/j.margen.2013.06.001>
- Librado, P., & Rozas, J. (2009). DnaSP V5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25(11), 1451-1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Liu, B., Zhang, K., Zhu, K., Shafi, M., Gong, L., Jiang, L., Liu, L., Muhammad, F., & Lü, Z. (2020). Population genetics of *Konosirus punctatus* in Chinese coastal waters inferred from two mtDNA genes (*COI* and *CytB*). *Frontiers in Marine Science*, 7. <https://doi.org/10.3389/fmars.2020.00534>
- Madduppa, H. H., Timm, J., & Kochzius, M. (2018). Reduced genetic diversity in the clown anemonefish *Amphiprion ocellaris* in exploited reefs of Spermonde Archipelago, Indonesia. *Frontiers in Marine Science*, 5, 1-8. <https://doi.org/10.3389/fmars.2018.00.080>
- Megarani, D. V., Nugroho, H. A., Andarini, Z. P., Surbakti, Y. D., & Widayanti, R. (2020). Genetic characterization and phylogenetic study of Indonesian indigenous catfish based on mitochondrial Cytochrome B gene. *Veterinary World*, 13(1), 96-103. <https://doi.org/10.14202/vetworld.2020.96-103>
- Mohd Radzi, B. (2019, May 21). Telur Ikan Terubok RM 500 sekilogram. *Harian Metro*. Retrieved October 10, 2022, from <https://www.hmetro.com.my/mutakhir/2019/05/458066/telur-ikan-Terubok-rm500-sekilogram>
- Mohd Yusoff, N. I., Mat Jaafar, T. N., Vilasri, V., Mohd Nor, S. A., Seah, Y. G., Habib, A., Wong, L. L., Danish-Daniel, M., Sung, Y. Y., Mazlan, A. G., Mat Piah, R., Ismail, S. I., & Tan, M. P. (2021). Genetic diversity, population structure and historical demography of the two-spined yellowtail stargazer (*Uranoscopus cognatus*). *Scientific Reports*, 11(1). <https://doi.org/10.1038/s41598-021-92905-6>
- Moreira, C., Correia, A. T., Vaz-Pires, P., & Froufe, E. (2019). Genetic diversity and population structure of the Blue Jack Mackerel *Trachurus picturatus* across its western distribution. *Journal of Fish Biology*, 94(5), 725-731. <https://doi.org/10.1111/jfb.13944>
- Munroe, T. A., Wongratana, T., & Nizinski, M. S. (1999) Clupeidae. Herrings (also, sardines, shads, sprats, pilchards, and menhadens). In: K. E. Carpenter, & V. H. Niem (Eds.), *The living marine resources of Western Central Pacific*. (Volume 3. Batoid fishes, chimeras and bony fishes part1 (Elopidae to Liophryniidae), pp. 1775-1821). FAO Rome.

- Mwita, C. J., & Chuhila, Y. J. (2023). Fish DNA: Advances and challenges. In W. S. Lakra, M. Goswami & V. L. Trudeau (Eds.), *Frontiers in aquaculture biotechnology* (pp. 171-185). Academic Press.
- Naeem, Z., Masud, S., Hassan, S., & Naeem, M. (2020). Molecular approach for identification of *Catla catla* using mitochondrial *COI* from Pakistan. *Mitochondrial DNA Part B*, 5(3), 3000-3003. <https://doi.org/10.1080/23802359.2020.1768913>
- Narejo, N. T., Lashari, P. K., & Jafri, S. I. H. (2008). Morphometric and meristic differences between two types of Palla, *Tenualosa ilisha* (Hamilton) from River Indus, Pakistan. *Pakistan Journal of Zoology*, 40(1), 31-35.
- Nei, M. (1987). *Molecular evolutionary genetics*. New York Chichester, West Sussex: Columbia University Press. <https://doi.org/10.7312/nei-92038>
- Nor Ain, M. R. (2021, December 13). Brisk business for 'terubok masin' traders. *New Straits Times*. Retrieved October 10, 2022, from <https://www.nst.com.my/news/nation/2021/12/753880/brisk-business-Terubok-masin-traders>
- Olavarría, C., Balbontín, F., Bernal, R., & Baker, C. S. (2006). Lack of divergence in the mitochondrial Cytochrome B gene between *Macruronus* species (Pisces: Merlucciidae) in the Southern Hemisphere. *New Zealand Journal of Marine and Freshwater Research*, 40(2), 299-304. <https://doi.org/10.1080/00288330.2006.9517422>
- Omer, A. S. (2017). Review on fish identification tools and their importance in biodiversity and fisheries assessments. *International Journal of Sciences: Basic and Applied Research*, 36(6), 118-126.
- Orsini, L., Vanoverbeke, J., Swillen, I., Mergeay, J., & De Meester, L. (2013). Drivers of population genetic differentiation in the wild: Isolation by dispersal limitation, isolation by adaptation and isolation by colonisation. *Molecular Ecology*, 22(24), 5983-5999. <https://doi.org/10.1111/mec.12561>
- Oszczka, P., Rola, K., Lenart-Boroń, A., & Boroń, P. (2014). High intraspecific genetic and morphological variation in the pioneer lichen *Cladonia rei* colonising slag dumps. *Open Life Sciences*, 9(5), 579-591. <https://doi.org/10.2478/s11535-014-0286-1>
- Panprommin, D., Soontornpravit, K., Tuncharoen, S., Pithakpol, S., Kannika, K., & Wongta, K. (2023). DNA barcoding for fish species identification and diversity assessment in the Mae Tam Reservoir, Thailand. *Fisheries and Aquatic Sciences*, 26(9), 548-557. <https://doi.org/10.47853/fas.2023.e47>
- Panprommin, D., Soontornpravit, K., Tuncharoen, S., Pithakpol, S., & Keereelang, J. (2019). DNA barcodes for the identification of species diversity in fish from Kwan Phayao, Thailand. *Journal of Asia-Pacific Biodiversity*, 12(3), 382-389.
- Parmaksız, A., & Eksi, E. (2017). Genetic diversity of the cyprinid fish *Capoeta trutta* (Heckel, 1843) populations from Euphrates and Tigris Rivers in Turkey based on mtDNA *COI* sequences. *Indian Journal of Fisheries*, 64(1), 18. <https://doi.org/10.21077/ijf.2017.64.1.62396-03>
- Pastorino, M. J., & Marchelli, P. (2022). Genetic resources: The base material for managing nature. In D. A DellaSala & M. I. Goldstein (Eds.), *Imperiled: The encyclopedia of conservation* (pp. 77-87). Elsevier.
- Prasertlux, S., Rongmung, P., Tang, S., Janpoom, S., Ninwichian, P., Khamnamtong, B., & Klinbunga, S. (2022). Assessing genetic diversity of wild populations and different color varieties for genetic improvement of Siamese Fighting Fish *Betta splendens* in Thailand. *Biochemical Genetics*. <https://doi.org/10.1007/s10528-022-10260-1>
- Puvanasundram, P., Esa, Y. B., Rahim, K. A. A., & Amin, S. M. N. (2018). Phylogeography and population structure of *Tenualosa toli*

- inferred from cytochrome B mitochondrial DNA fragment. *Journal of Environmental Biology*, 39(5), 895-906. [https://doi.org/10.22438/jeb/39/5\(si\)/26](https://doi.org/10.22438/jeb/39/5(si)/26)
- Ramos-Onsins, S. E., & Rozas, J. (2002). Statistical properties of new neutrality tests against population growth. *Molecular Biology and Evolution*, 19(12), 2092-2100. <https://doi.org/10.1093/oxfordjournals.molbev.a004034>
- Ratnasingham, S., & Hebert, P. D. N. (2013). A DNA-based registry for all animal species: The barcode index number (BIN) system. *PLOS ONE*, 8, e66213.
- Robert, A. A., Taha, H., Metali, F., Ahmad, N., & Arai, T. (2019). Validation of occurrence of tropical shads, *Tenualosa Ilisha* (Hamilton, 1822) and *T. toli* (valenciennes, 1847) (Teleostei, Clupeidae), in Malaysian waters. *Check List*, 15(1), 65-69. <https://doi.org/10.15560/15.1.65>
- Rogers, A. R., & Harpending, H. (1992). Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution*, 9(3), 552-569. <https://doi.org/10.1093/oxfordjournals.molbev.a040727>
- Roomiani, L., Sotudeh, A.M. & Hakimi Mofrad, R. (2014). Reproductive biology of Hilsa shad (*Tenualosa ilisha*) in coastal waters of the Northwest of Persian Gulf. *Iranian Journal of Fisheries Sciences*, 13(1), 201-215.
- Sá-Pinto, A., Branco, M. S., Alexandrino, P. B., Fontaine, M. C., & Baird, S. J. (2012). Barriers to gene flow in the marine environment: Insights from two common intertidal limpet species of the Atlantic and Mediterranean. *PLOS ONE*, 7(12). <https://doi.org/10.1371/journal.pone.0050330>
- Sarker, A., Jiang, J., Naher, H., Huang, J., Sarker, K. K., Yin, G., Baki, M. A., & Li, C. (2021). Cross-species gene enrichment revealed a single population of Hilsa Shad (*Tenualosa ilisha*) with low genetic variation in Bangladesh Waters. *Scientific Reports*, 11(1). <https://doi.org/10.1038/s41598-021-90864-6>
- Shi, S.-F., Li, M., Yan, S., Wang, M., Yang, C.-P., Lun, Z.-R., Brown, C. L., & Yang, T.-B. (2014). Phylogeography and demographic history of *Gotocotyla sawara* (Monogenea: Gotocotylidae) on Japanese Spanish mackerel (*Scomberomorus niphonius*) along the coast of China. *Journal of Parasitology*, 100(1), 85-92. <https://doi.org/10.1645/13-235.1>
- Shrestha, B. B., Khatiwada, J. R., & Thanet, D. R. (2019). MtDNA confirms the presence of *Moschus leucogaster* (Ruminantia, Moschidae) in Gaurishankar Conservation Area, Nepal. *Arxius De Miscel·Lània Zoològica*, 17, 209-218. <https://doi.org/10.32800/amz.2019.17.0209>
- Sout, N. M., Othman, R., Suliman, N. A., Murgaya, D., Harmin, S. A., & Zan, M. S. (2023). Phylogenetics analysis of blackwater fishes in peat swamp forest using Cytochrome B gene. *AIP Conference Proceedings*, 2625(1), 020011. <https://doi.org/10.1063/5.0129302>
- Spielman, D., Brook, B. W., & Frankham, R. (2004). Most species are not driven to extinction before genetic factors impact them. *Proceedings of the National Academy of Sciences*, 101(42), 15261-15264. <https://doi.org/10.1073/pnas.0403809101>
- Sun, P., Yu, J., Tang, B., & Liu, Z. (2021). Gene variation and population structure of *pampus chinensis* in the China coast revealed by mitochondrial control region sequences. *Mitochondrial DNA Part B*, 6(8), 2240-2245. <https://doi.org/10.1080/23802359.2021.1878963>
- Supmee, V., Sawusdee, A., Sangthong, P., & Suppapan, J. (2020). Population genetic structure of blue swimming crab (*Portunus pelagicus*) in the Gulf of Thailand. *Biodiversitas Journal of Biological Diversity*, 21(9). <https://doi.org/10.13057/biodiv/d210943>

- Taboun, Z. S., Walter, R. P., Ovenden, J. R., & Heath, D. D. (2021). Spatial and temporal genetic variation in an exploited reef fish: The effects of exploitation on cohort genetic structure. *Evolutionary Applications*, *14*(5), 1286-1300. <https://doi.org/10.1111/eva.13198>
- Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, *123*(3), 585-595. <https://doi.org/10.1093/genetics/123.3.585>
- Teacher, A. G., & Griffiths, D. J. (2010). HapStar: Automated haplotype network layout and visualization. *Molecular Ecology Resources*, *11*(1), 151-153. <https://doi.org/10.1111/j.1755-0998.2010.02890.x>
- Torquato, F., Range, P., Ben-Hamadou, R., Sigsgaard, E. E., Thomsen, P. F., Riera, R., Berumen, M. L., Burt, J. A., Feary, D. A., Marshall, A., D'Agostino, D., DiBattista, J. D., & Møller, P. R. (2019). Consequences of marine barriers for genetic diversity of the coral-specialist yellowbar Angelfish from the northwestern Indian Ocean. *Ecology and Evolution*, *9*(19), 11215-11226. <https://doi.org/10.1002/ece3.5622>
- Trang, V. T., Ha, T. T., Ngan, T. T., Vinh, N. D., Nhat, P. H., & Huyen, V. T. (2022). Genetic diversity of *Chanos chanos* (forsskål, 1775) from natural populations in Vietnam. *Vietnam Journal of Agricultural Sciences*, *5*(2), 1456-1463. <https://doi.org/10.31817/vjas.2022.5.2.03>
- Tschá, M. K., Baggio, R. A., Marteleto, F. M., Abilhoa, V., Bachmann, L., & Boeger, W. A. (2016). Sea-level variations have influenced the demographic history of estuarine and freshwater fishes of the coastal plain of Paraná, Brazil. *Journal of Fish Biology*, *90*(3), 968-979. <https://doi.org/10.1111/jfb.13211>
- Verma, R., Singh, M., & Kumar, S. (2016). Unravelling the limits of mitochondrial control region to estimate the fine scale population genetic differentiation in anadromous fish *Tenualosa ilisha*. *Scientifica*, *2016*, 1-9. <https://doi.org/10.1155/2016/2035240>
- Villanueva, B., Fernández, A., Peiró-Pastor, R., Peñaloza, C., Houston, R. D., Sonesson, A. K., Tsigenopoulos, C. S., Bargelloni, L., Gamsız, K., Karahan, B., Gökçek, E., Fernández, J., & Saura, M. (2022). Population structure and genetic variability in wild and farmed Mediterranean populations of gilthead seabream and European Seabass inferred from a 60K combined species SNP array. *Aquaculture Reports*, *24*, 101145. <https://doi.org/10.1016/j.aqrep.2022.101145>
- Waludin, S. N., Mohamed, A. A., Ing, C. P., & Mukai, Y. (2018). Genetic diversity of the orange-spotted grouper (*Epinephelus coioides*) in Terengganu Malaysia based on mitochondrial cytochrome B sequence data. *Malaysian Applied Biology*, *47*(6), 61-68.
- Wang, L., Liu, S., Zhuang, Z., Guo, L., Meng, Z., & Lin, H. (2013). Population genetic studies revealed local adaptation in a high gene-flow marine fish, the small yellow croaker (*Larimichthys polyactis*). *PLOS ONE*, *8*(12). <https://doi.org/10.1371/journal.pone.0083493>
- Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R., & Hebert, P. D. N. (2005). DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *360*(1462), 1847-1857. <https://doi.org/10.1098/rstb.2005.1716>
- Whitehead, P. J. P. (1985). *FAO Species Catalogue 7, Clupied Fishes of the World. Part I Chirocentridae, Clupeidae and Pristigasteridae* (Volume 7, pp. 303), FAO Fisheries.
- Williford, D., DeYoung, R. W., Honeycutt, R. L., Brennan, L. A., Hernández, F., Heffelfinger, J. R., & Harveson, L. A. (2014). Phylogeography of the Gambel's quail (*Callipepla gambelii*) of Western

- North America. *The Wilson Journal of Ornithology*, 126(2), 218-235. <https://doi.org/10.1676/13-111.1>
- Winters, K. L., van Herwerden, L., Choat, J. H., & Robertson, D. R. (2010). Phylogeography of the Indo-Pacific parrotfish *Scarus psittacus*: Isolation generates distinctive peripheral populations in two oceans. *Marine Biology*, 157(8), 1679-1691. <https://doi.org/10.1007/s00227-010-1442-4>
- Yao, G., Zhang, H., Xiong, P., Jia, H., Shi, Y., & He, M. (2022). Community characteristics and genetic diversity of macrobenthos in Haima Cold Seep. *Frontiers in Marine Science*, 9. <https://doi.org/10.3389/fmars.2022.920327>
- Yin, F., Gasser, R. B., Li, F., Bao, M., Huang, W., Zou, F., Zhao, G., Wang, C., Yang, X., Zhou, Y., Zhao, J., Fang, R., & Hu, M. (2013). Genetic variability within and among *Haemonchus contortus* isolates from goats and sheep in China. *Parasites & Vectors*, 6(1). <https://doi.org/10.1186/1756-3305-6-279>
- Yu, H. J., Kai, Y., & Kim, J.-K. (2016). Genetic diversity and population structure of *Hyporhamphus sajori* (Belontiiformes: Hemiramphidae) inferred from mtDNA control region and msDNA markers. *Journal of Fish Biology*, 89(6), 2607-2624. <https://doi.org/10.1111/jfb.13152>
- Zafar, S., Jabeen, F., Ali, M., & Sajjad, A. (2024). Assessment of genetic diversity of freshwater genus *Labeo* through DNA barcoding and phylogenetic analysis inhabited in the River Indus. *Journal of King Saud University - Science*, 36(4), 1031-1033. <https://doi.org/10.1016/j.jksus.2024.103123>
- Zhai, D., Zhang, Z., Zhang, F., Liu, H., Cao, W., & Gao, X. (2019). Genetic diversity and population structure of a cyprinid fish (*Ancherythroculter nigrocauda*) in a highly fragmented river. *Journal of Applied Ichthyology*, 35(3), 701-708. <https://doi.org/10.1111/jai.13897>
- Zhang, G., Chen, C., Lu, W., Li, J., Fang, T., Yang, K., Zhao, X., Gao, N., & Liang, Y. (2023). Genetic diversity and phylogeography of *Taenioides cirratus* in five geographical populations based on mitochondrial *COI* and *CytB* gene sequences. *Journal of Applied Ichthyology*, 2023, 1-9. <https://doi.org/10.1155/2023/4459823>
- Zhang, L., Zhou, Z., Li, H., Qiao, Y., & Zhang, Y. (2019). The mitochondrial COI gene reveals the genetic diversity of chub mackerel (*Scomber japonicus*) in the Taiwan Strait and its adjacent waters. *Pakistan Journal of Zoology*, 51(3), 935-942. <https://doi.org/10.17582/journal.pjz/2019.51.3.935.943>
- Zhang, Q., Sun, C., Zhu, Y., Xu, N., & Liu, H. (2020). Genetic diversity and structure of the round-tailed paradise fish (*Macropodus ocellatus*): Implications for population management. *Global Ecology and Conservation*, 21. <https://doi.org/10.1016/j.gecco.2019.e00876>
- Zheng, W., Zou, L., & Han, Z. (2015). Genetic analysis of the populations of Japanese anchovy *Engraulis japonicus* from the Yellow Sea and East China Sea based on mitochondrial cytochrome B sequence. *Biochemical Systematics and Ecology*, 58, 169-177. <https://doi.org/10.1016/j.bse.2014.12.007>
- Zhou, M.-Y., Wang, J.-J., Ren, J.-F., Li, F., Wu, J.-X., Zhou, J.-J., Li, J.-L., Yang, J.-Q., & Lin, H.-D. (2023). Historical landscape evolution shaped the phylogeography and population history of the cyprinid fishes of *Acrossocheilus* (Cypriniformes: Cyprinidae) according to mitochondrial DNA in Zhejiang Province, China. *Diversity*, 15(3), 425. <https://doi.org/10.3390/d15030425>