

FIRST RECORD OF INSECTS IN PULAU TINGGI, JOHOR, MALAYSIA

YIN HUI NG¹, AQILAH AWG ABDUL RAHMAN^{1,2*}, MD-ZAIRI ZAINAL³ AND MUHAMMAD ABU BAKAR ABDUL-LATIFF^{1,4*}

¹Department of Technology and Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (Pagoh Campus) 84600 Muar, Johor, Malaysia. ²Centre of Research for Sustainable Uses of Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (Pagoh Campus), 84600 Muar, Johor, Malaysia. ³Shaz Resort Pulau Tinggi Sdn. Bhd., Lot 44, Kampung Penaga, Pulau Tinggi, 86800 Mersing, Johor, Malaysia. ⁴Environmental Management and Conservation Research Unit (eNCORe), Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (Pagoh Campus), 84600 Muar, Johor, Malaysia.

*Corresponding authors: aqilah@uthm.edu.my, latiff@uthm.edu.my

Submitted final draft: 17 February 2022 Accepted: 26 March 2022

<http://doi.org/10.46754/jssm.2022.11.011>

Abstract: Species assembly of insects in Pulau Tinggi was extensively studied to obtain the first checklist of insect fauna on the island. Samples were collected from forested areas in Pulau Tinggi on three occasions (20-30 April, 1-10 May and 20-30 June 2019). Various sampling methods were used, including pitfall trap, light trap and active collection using sweep net and forceps. The specimens were dried and curated for identification. A total of 41 families from ten orders of insects representing 90 species and morphospecies were identified. Coleoptera was the most dominant order with 14 families, followed by seven families from the order Hemiptera, three from Hymenoptera, four from Orthoptera, three from Odonata, three from Diptera, two from Blattodea, one from Mantodea and one family from order Phasmatodea. This first inventory study has shown that this relatively unexploited island possesses many insect species. Hopefully, this data will assist policymakers and governing bodies, particularly the private landowners and Johor State Government in making conservation management decisions in the future.

Keywords: Coleoptera, Hemiptera, Mersing, Seribuat Archipelago, Malaysia, South China Sea.

Introduction

Insects are a crucial component for a functional natural and human-modified landscape. They play an important role in several ecosystem services that ensure the quality of agriculture, tourism and natural resources for humans (Saikim *et al.*, 2020; Hor *et al.*, 2021). Most insects have high survival and reproduction rate. They are a very successful group of organisms that can live in all types of ecosystems. They maintain the nutrient cycle by assisting the degradation of wood and leaf debris and diffusion of fungi, eliminating the decaying flesh and faeces, loosening the soil, disseminating plants or seeds, and being a food source for terrestrial and aquatic lives (Gullan & Cranston, 2010). Insects are also used worldwide as biomonitoring tools to measure diversity and ecosystem health (Rákossy & Schmitt, 2011; Shuey *et al.*, 2017). Globally, insect numbers are already in decline (Basset & Lamarre, 2019). If nothing is done to conserve

this group of organisms, Malaysia will soon see a significant insect number decline. Malaysia is already well known for its highly diverse fauna and iconic insect species (Nur Atiqah *et al.*, 2017; Hor *et al.*, 2021; Ridzuan *et al.*, 2021). Ranked as one of the most mega-diverse countries in the world, Malaysia is now focused on protecting its natural treasures, as stated in Malaysia's National Policy on Biological Diversity (2016-2025) (NRE, 2016). Within Malaysia, documentation of insect diversity has been carried out extensively in Borneo and some parts of Peninsular Malaysia (Nur Atiqah *et al.*, 2017; Chung *et al.*, 2020). New species of insects have been discovered quite frequently in recent years (Al-Talafha *et al.*, 2018; Gildenkov, 2019; Ng & Mound, 2020). However, knowledge of insect diversity in Malaysian island forests is still lacking. An island ecosystem provides a habitat for migratory, endemic and endangered species (Quek *et al.*, 1999; Francini & Ramos,

2014; Mohd-Taib *et al.*, 2019). Therefore, it makes sense to pay particular attention to the remarkable biodiversity in an island ecosystem.

Pulau Tinggi (02° 18' 0" N, 140° 07' 0" E) is located in the middle arc of Seribuat Archipelago, Mersing, Johor (Figure 1). The island has an area of 13.5 km² and an elevation of 610 m. Pulau Tinggi has a relatively low human disturbance and is covered with different vegetation types. The archipelago also consists of several main islands including Pulau Tioman, Pulau Pemanggil, Pulau Aur and Pulau Sibiu. Prior to the publication of this study, several inventory studies focusing on general insect groups or specific orders have been done in Malaysian islands (Quek *et al.*, 1999; Farizawati *et al.*, 2014; Rosmidi *et al.*, 2017; Mohamed *et al.*, 2019; Ridzuan *et al.*, 2021). However, none was done in Pulau Tinggi.

To our knowledge, no entomological research has been done on this relatively unexplored island. Therefore, this paper aims to record the various insect groups that inhabit the area. However, insects from the order Lepidoptera were not included in this paper. From the authors' preliminary observation, Lepidoptera of Pulau Tinggi requires a more

extensive discussion due to their relatively high abundance in Pulau Tinggi. Therefore, a separate paper will be published focusing on only the island's butterflies and moths.

Materials and Methods

Study Area

The study site is at Gunung Semudu in Tanjung Balang, Pulau Tinggi. Sampling was conducted within three hiking trails. Trail 1 and Trail 2 are located within a secondary forest while Trail 3 is located within a primary forest (Figure 2). The coordinates and elevation of each trail are recorded in Table 1.

Insect samples were collected from the forest area for three months within ten consecutive days each month (20-30 April, 1-10 May and 20-30 June 2019). The sampling was carried out in two replications. The sampling techniques consisted of active collection, pitfall trapping and light trapping. All samples were collected in three separate timings throughout the day: Morning (08:00 hours), afternoon (13:00 hours) and evening (17:00 hours), except light trapping, which was conducted at night (20:00 hours) for nocturnal insects.




Figure 1: Map of Pulau Tinggi and its location relative to Peninsular Malaysia and the Southern Seribuat Archipelago




Figure 2: Study trail located in Gunung Semudu

Table 1: Coordinate and elevation of each sampling trail used in this study

Trail	Sub-trail	Coordinates
1	1A	(02° 17' 06.1" N, 104° 06' 55.9" E, 4 m)
	1B	(02° 17' 07.7" N, 104° 06' 53.1" E, 19 m)
	1C	(02° 17' 38.5" N, 104° 07' 39.3" E, 56 m)
2	2A	(02° 17' 05.4" N, 104° 07' 18.9" E, 6 m)
	2B	(02° 17' 24.5" N, 104° 07' 16.3" E, 53 m)
	2C	(02° 17' 18.5" N, 104° 07' 05.8" E, 66 m)
3	3A	(02° 17' 27.6" N, 104° 07' 19.9" E, 80 m)
	3B	(02° 17' 42.3" N, 104° 07' 38.3" E, 118 m)
	3C	(02° 17' 31.1" N, 104° 07' 36.0" E, 142 m)

Sweep Net

The active collection was done opportunistically similar to the method by Chung *et al.* (2020). Flying insects were caught using sweep nets and put in glassine envelopes while crawling insects were sampled using fine-tipped forceps and kept in vials with 75% ethanol.

Light Trapping

Four sets of light traps were installed in light trapping from 20:00 hours until 01:00 hours.

A 2 × 2 m white fabric attached to a rope was hung vertically and lit by a 160-watt mercury bulb. The four sets of light traps were employed at three different trails. The traps were placed 300 m from the seashore (1A) at Trail A, at a mangrove area (2A) and shrub area (2B) at Trail 2, the entrance (3A) of Trail 3. Nocturnal insects attracted to the light were collected using a killing jar filled with a tissue soaked with ethyl acetate. The specimens were kept in 75% ethanol prior to drying and pinning.

Pitfall Trapping

In the pitfall trapping method, five sets of pitfall traps were set up randomly at different elevations ranging from 4 meters to 142 meters above sea level. Each set consisted of five traps with different bait types: Candy, stone, tofu, whiskey and salted fish. Cups with a width of 10 cm and depth of 15 cm were buried in the forest floor as traps. Some soapy water was poured into the trap to prevent the insects from escaping. Each pitfall trap was covered with dried leaves to prevent water from leaking into the trap. The five sets of pitfall traps were installed at a shrub area (1B) at Trail 1, at a mangrove area (2C) at Trail 2 and near the waterfall entrance (3A and 3B) at Trail 3.

Species Identification

Specimens were brought back to the taxonomy lab at the Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia for identification using stereo microscopes. Specimens were identified to species level using keys by Romoser and Stoffolano (1998), Borror *et al.* (2005), Buck *et al.* (2009), Choate (2011a), Choate *et al.* (2011), Choate (2011b) and Orr (2014). At the time of publication, identification of doubtful species without comparison with type specimens and museum collections were denoted as ‘unknown sp.’ in Table 2.

Table 2: Checklist of insects collected in Pulau Tinggi

Order	Family	Subfamily	Species
Coleoptera	Brentidae	Brentinae	Unknown sp. 1
	Lampyridae	Lampyrinae	<i>Pyrocoelia fumigata</i> (Gorham, 1880)
	Cantharidae	Silinae	<i>Laemoglyptus</i> sp.
	Chrysomelidae	Eumolpinae	<i>Abirus aeneus</i> (Wiedemann, 1821)
			<i>Abirus elegans</i> (Baly, 1864)
			<i>Basilepta anthracina</i> (Baly, 1867)
			<i>Basilepta lateripunctatum</i> (Baly, 1867)
			<i>Cleorina collaris</i> (Baly, 1867)
			<i>Colaspoides cuprea</i> (Baly, 1867)
			<i>Colaspoides parvula</i> (Baly, 1867)
	<i>Phytorus pinguis</i> (Baly, 1867)		
	<i>Platycorynus</i> sp.		
		Alticinae	<i>Chabria angulicollis</i> (Clark, 1865)
		Galerucinae	<i>Hoplosaenidea abdominalis</i> (Jacoby, 1884)
Erotylidae	Erotylinae	<i>Episcapha</i> sp.	
Carabidae	Cicindelinae	<i>Cicindela aurulenta</i> (Fabricius, 1801)	
		<i>Cylindera viduata</i> (Fabricius, 1801)	
		<i>Tricondyla</i> sp.	
	Lebiinae	<i>Catascopus phlogops</i> (Andrewes, 1926)	
Cleridae	Neorthopleurinae	<i>Allochotes scymnoides</i> (Westwood, 1875)	
Cerambycidae	Cerambycinae	<i>Hoplocerambyx spinicornis</i> (Newman, 1842)	

		Lamiinae	<i>Acalolepta</i> sp. <i>Gnoma</i> sp. <i>Epepeotes lateralis</i> (Guerin Meneville, 1831)	
	Coccinellidae	Coccinellinae	<i>Heteroneda billardieri</i> (Crotch, 1871)	
	Elateridae	Elaterinae	<i>Agonischius</i> sp.	
	Passalidae		<i>Odontotaenius</i> sp.	
	Scarabaeidae	Rutelinae	Unknown sp. 2	
		Scarabaeinae	<i>Onthophagus</i> sp. 1 <i>Onthophagus</i> sp. 2	
		Dynastinae	<i>Oryctes rhinoceros</i> (Linnaeus, 1758) <i>Oryctes</i> sp. <i>Xylotrupes gideon</i> (Linnaeus, 1767)	
	Staphylinidae	Aleocharinae	Unknown sp. 3	
	Tenebrionidae		Unknown sp. 4	
			Unknown sp. 5	
			Unknown sp. 6	
Blattodea	Blattidae	Blattinae	<i>Periplaneta americana</i> Unknown sp. 7	
		Epilamprinae	Unknown sp. 8	
	Termitidae	Nasutitermitinae	<i>Hospitalitermes umbrinus</i> (Haviland, 1898)	
		Macrotermitinae	<i>Macrotermes carbonarius</i> (Hagen, 1858)	
Diptera	Tephritidae	Trypetinae	<i>Euphranta maculifemur</i> (Meijere, 1924)	
	Muscidae	Muscinae	<i>Neomyia</i> sp.	
	Dolichopodidae	Dolichopodinae	<i>Paraclius</i> sp.	
Hemiptera	Reduviidae	Reduviinae	<i>Acanthaspis inermis</i> (Stahl, 1870)	
		Harpactorinae	Unknown sp. 9 Unknown sp. 10 <i>Campsolomus</i> sp.	
		Ectrichodiinae	Unknown sp. 11	
		Stenopodainae	<i>Aulacogenia patalungae</i> (Miller, 1940)	
	Membracidae	Membracinae	Unknown sp. 12	
	Pentatomidae	Pentatominae	<i>Antestiopsis anchora</i> (Thunberg, 1783)	
	Coreidae	Coreinae	<i>Physomerus grossipes</i> (Fabricius, 1794)	
	Largidae	Physopeltinae	<i>Physopelta gutta</i> (Burmeister, 1834)	
	Pyrrhocoridae	Pyrrhocorinae	Unknown sp. 13	
			<i>Dysdercus decussatus</i> (Boisduval, 1835)	
	Tessaratomidae	Tessaratominae	Unknown sp. 14	
	Phasmatodea			Unknown sp. 15

Hymenoptera	Formicidae	Formicinae	<i>Anoplolepis</i> sp. <i>Dinomyrmex gigas</i> (Latreille, 1802) <i>Camponotus</i> sp. <i>Oecophylla smaragdina</i> (Fabricius, 1775) <i>Polyrhachis</i> sp. <i>Polyrhachis proxima</i> (Roger, 1863)	
		Myrmicinae	<i>Crematogaster</i> sp.	
		Pseudomyrmecinae	<i>Tetraponera</i> sp.	
	Vespidae	Eumeninae	<i>Delta</i> sp. (Fabricius, 1775)	
	Apidae	Apinae	Unknown sp. 16	
		Xylocopinae	<i>Ceratina</i> sp.	
Mantodea	Mantidae	Mantinae	<i>Rhombodera</i> sp.	
Odonata	Aeshnidae	Aeshninae	<i>Anax guttatus</i> (Burmeister, 1839)	
		Libellulidae	<i>Orthetrum testaceum</i> (Burmeister, 1839) <i>Orthetrum glaucum</i> (Brauer, 1865) <i>Lathrecista asiatica</i> (Fabricius, 1798) <i>Neurothemis fluctuans</i> (Fabricius, 1793) <i>Neurothemis</i> sp. <i>Trithemis festiva</i> (Rambur, 1842) <i>Agrionoptera insignis</i> (Rambur, 1842)	
	Argiolestidae	Argiolestinae	Unknown sp. 17	
	Orthoptera	Acrididae	Gomphocerinae	<i>Gonista</i> sp.
		Pyrgomorphidae	Pyrgomorphinae	<i>Atractomorpha</i> sp.
Tettigoniidae		Phaneropterinae	<i>Ducetia malayana</i> (Heller, 2017) <i>Elimaea</i> sp.	
		Conocephalinae	<i>Euconocephalus varius</i> (Walker, 1869)	
		Gryllidae	Eneopterinae	<i>Lebinthus</i> sp.
Euscyrtinae			<i>Euscyrtus</i> sp.	
Landrevinae			<i>Endodrelanva jimini</i> (Tan & Kamaruddin, 2016)	
Trigonidiinae	<i>Amusurgus</i> sp.			

Results and Discussion

The first insect checklist for Pulau Tinggi produced from this study is shown in Table 2. Insects on this island are exceptionally diverse. A total of ten insect orders consisting of 41 families and 90 species were recorded. Most of the insects were collected from light traps.

Beetles (Coleoptera)

Coleoptera was the most dominant order with 14 families recorded. They are commonly

found in an open and disturbed area in Pulau Tinggi. A species of firefly was discovered here, namely *Pyrocoelia fumigata*. *P. fumigata* has also been recorded in Singapore (Jusoh *et al.*, 2021). The presence of firefly species is a good sign for the island as these charismatic insects provide a sense of allurements and wonder for conservation philanthropists. Further studies may evaluate the potential of firefly tourism in Pulau Tinggi. Lewis *et al.* (2021) estimated that over one million tourists had travelled annually for firefly watching worldwide in recent years.

Several rhinoceros beetles, namely *Xylotrupes Gideon* (brown rhinoceros beetle) and *Oryctes rhinoceros* (coconut palm rhinoceros beetle) were also recorded here. This is not surprising considering *O. rhinoceros* and *X. Gideon* are considered pests in an oil palm plantation which is abundantly available in the mainland of Johor (Pradipta *et al.*, 2020; Mat Ripin & Md. Latip, 2021).

True Bugs (Hemiptera)

Hemipterans were the second most abundant insect order discovered here in Pulau Tinggi. A total of seven families were recorded which includes 12 species and morphospecies. Notable species include *Acanthaspis inermis* (Stahl, 1870) and *Aulacogenia patalungae* (Miller, 1940) from the Reduviidae family. These species are well known to be distributed in Peninsular Malaysia, Singapore and Thailand (Tan *et al.*, 2015). Many other species in this order were not identified as species or even genera at the time of publication as no type specimens or museum collections of that particular morphospecies were found for comparison.

Termites and Cockroaches (Blattodea)

Both termites (Termitidae) and cockroaches (Blattidae) were recorded in Pulau Tinggi. One of the termite species recorded was *Macrotermes carbonarius*. While termites are an economical liability due to their ability to cause serious damage to structures in their natural habitat, termites help decompose organic matter, recycle nutrients and improve soil quality (Enagbonma & Babalola, 2019). The discovery of the American cockroach (*Periplaneta americana*) on this island was not shocking as this species is a common pest in human dwellings (Kamarol Zaman & Omar, 2019).

Ants, Bees and Wasps (Hymenoptera)

Three families of Hymenoptera were recorded which included ants (Formicidae), bees (Apidae) and wasps (Vespidae). The giant forest

ant, known as *Dinomyrmex gigas* was recorded in Endau Rompin Johor National Park (Shafiq & Maryati, 2021). This largely nocturnal insect is easily spotted in lowland tropical rainforests due to its large size and colour strikingly different from other ants. On the other hand, a more common ant species, the weaver ant (*Oecophylla smaragdina*) was also recorded here. The weaver ant is widely distributed throughout most of the Oriental region, from India to the Solomon Islands (Greenslade, 1972). On top of that, *O. smaragdina* is known to highly prefer food with higher protein content (Pimid *et al.*, 2019). In this survey, tuna was used in the bait traps which attracted the foraging behaviour of the nearby *O. smaragdina* colony.

Dragonfly (Odonata)

Three families from the order Odonata were recorded in Pulau Tinggi. Notable species include *Agrionoptera insignis* and *Neurothemis fluctuans* which have also been recorded in the forest of Pulau Pangkor (Farizawati *et al.*, 2014), Pulau Tioman (Choong *et al.*, 2017) and in more than eight localities in Johor mainland (Abdul Aziz *et al.*, 2018). It is worth noting that Pulau Tioman is also part of the Seribu Archipelago. Being a generalist species, *N. fluctuans* is more widespread, with additional records in Pulau Carey, Pulau Langkawi and Pulau Pinang (Farizawati *et al.*, 2014). In Pulau Tinggi and Pulau Sibul, *Orthetrum testaceum* and *N. fluctuans* were the most abundant species from the Libellulidae family (Azmi & Haris-Hussain, 2019). *O. testaceum* (Scarlet skimmer) and *N. fluctuans* (Grasshawk dragonfly) have also been reported in three National Parks in Sarawak (Kubah National Park, Lambir Hills National Park and Similajau National Park) (Dow *et al.*, 2013). Members from the Libellulidae family are regularly found as the most dominant in tropical and subtropical forests. Additionally, their distribution is common in various habitats due to their ability to breed in open and disturbed habitats (Azmi & Haris-Hussain, 2019).

Other Insect Orders (Diptera, Mantodea, Phasmatodea, Orthoptera)

Pulau Tinggi shares only six similar species with Pulau Sibul, even though they are part of the same Seribuat Archipelago (Azmi & Haris-Hussain, 2019). Overall, Pulau Sibul recorded 35 species of insects belonging to 12 orders and 24 families. The similar species were *Oryctes rhinoceros*, *Camponotus* sp., *Oecophylla smaragdina*, *Orthetrum testaceum*, *Lathrecista asiatica* and *Neurothemis fluctuans*. These are all common species found in human-modified landscapes due to their generalist capabilities (Pimid et al., 2019; Mat Ripin & Md. Latip, 2021). It has to be taken into consideration that most insects collected in this study were unidentifiable to the genus or species level. This shows that many studies and fundamental research still need to be done on the fauna of Pulau Tinggi. This is particularly true for insects since approximately one million have been described to date, with an estimated 5 million species left undescribed (Stork et al., 2015).

Issues and Implications Affecting Insect Diversity in Pulau Tinggi

Challenges in species-level identification are one of the major reasons why entomological data are not mainly used to justify conservation. The insect group is often considered too large and difficult to use. Any misidentification may compromise conservation research as these may result in overestimating or underestimating insect diversity. Hopefully, unidentified samples from this research will help advance the specific group's taxonomy in the future. This is crucial since to protect a taxon, one must identify and describe it first. It is impossible to protect something that is yet to be described.

Compared to other islands in Peninsular Malaysia, Pulau Tinggi is relatively unexplored. The forested and riverine area of Gunung Semudu is a source of water supply for nearby villages. The pre-existing terrestrial and marine ecosystems show great potential for naturalists and city dwellers to relax and immerse in nature (Akashah et al., 2021). As shown in this

study, a relatively high insect species richness was recorded. Therefore, we must continue protecting the forest and marine ecosystem in line with the National Policy of Biological Diversity (NPBD 2016-2025) (NRE, 2016).

Conclusion

The diverse forest of Malaysia is a habitat for many insect species. Insects play an important role in maintaining ecosystem balance. Unfortunately, they are the most ignored group due to their small size. This first documentation of insects in Pulau Tinggi is important as it has shown that this relatively unexploited island has many insect species. Conservation action should be taken immediately to preserve the insect diversity in Pulau Tinggi. Any development to be done on the island should consider the health of the ecosystem surrounding it. Consistent monitoring of insect diversity here is crucial to ensure the sustainability of the island's ecosystem. Perhaps, more intensive research on insects in the Seribuat Archipelago can be done in the future to open up future possibilities in the aspect of biodiversity conservation and tourism.

Acknowledgements

This project is funded by the Ministry of Higher Education Malaysia (MOHE) under the Malaysian Technical University Network (MTUN) grant scheme Vote K121 and Industrial Grant by Shaz Resort Sdn. Bhd. (UTHM-SHAZ-M004), both grants awarded to Associate Professor Ts. Dr. Muhammad Abdul Latiff Abu Bakar (UTHM) as the Principal Researcher. This research was supported by GPPS-UTHM-2018-H288 postgraduate grant from Universiti Tun Hussein Onn Malaysia (UTHM), Malaysia. We are grateful to Dato' Abdul Kadir Abu Hashim, Director General of the Department of Wildlife and National Parks, who provided us with the necessary facilities and assistance. This research was conducted under a research permit (JPHL & TN (IP): 600-6/1/4 (03)/M-00351-15-19). We are deeply indebted to the Department of Wildlife and National Parks Malaysia for

granting permission to carry out this research. The authors acknowledge the Ministry of Higher Education Malaysia, Universiti Tun Hussein Onn Malaysia and Shaz Resort Sdn. Bhd. for providing necessary funding, facilities and assistance.

References

- Abdul Aziz, M. A. A., Mohamed, M., & Tokiman, L. (2018). Faunistic Studies of Odonates (Insecta: Odonata) in Johor, Peninsular Malaysia. *Serangga*, 23(2), 14-35.
- Akashah N., Kamarudin K. R., Solehin, S. N., Rahman, N. S. A., Akma, U. N., Shahdan, F., Azman, H., Fadzil, S. N. M., Faid, N. H. M., Zaman, N. S. S., Badrulhisham, N. S., Bakar, M. A. L. A., Legiman, M. I., Rehan, A. M., Salleh, F. M., & Esa Y (2021). Species identification and relationship of sea cucumber species from Pulau Tinggi and Sedili Kechil, Johor based on ossicle shape. *Journal of Sustainable Natural Resources*, 2(1), 38-45.
- Al-Talafha, H. A., Yaakop, S., & Idris, A. B. (2018). Two New Species and Seven New Records of Horse Fly (Diptera: Tabanidae) from Malaysia, Including a Description of New Species and Modified Keys. *Journal of Medical Entomology*, 55(1), 112-121. <https://doi.org/10.1093/jme/tjx172>
- Azmi, W. A., & Haris-Hussain, M. (2019). A survey on diversity of insects in Pulau Sibul, Johor. *Malayan Nature Journal*, 71(3), 349-359.
- Basset, Y., & Lamarre, G. P. A. (2019). Toward a world that values insects: Rapid adoption of conservation measures is key to protecting insect populations. *Science*, 364(6447), 1230-1231. <https://doi.org/10.1126/science.aaw7071>
- Borror, D., Triplehord, C., & Johnson, N. (2005). *An Introduction to the Study of Insects* (7th ed.). Thomson Brooks Cole.
- Buck, M., Woodley, N. E., Borkent, A., Wood, M., Pape, T., Vockeroth, R., Michelsen, V., & Marshall, S. (2009). Key to diptera families - Adults. In B. Brown, Borkent A., Cumming J. M., Wood D. M., Woodley N. E., & Zumbado M. (Eds.), *Manual of Central American Diptera* (1st ed.). NRC Research Press.
- Choate, P. M. (2011a). A Dichotomous Key for the Identification of the Cockroach fauna (Insecta: Blattaria) of Florida. In *Florida Blattaria* (pp. 1-18). University of Florida.
- Choate, P. M. (2011b). Identification Key to the Principal Families of Florida Heteroptera. In *Identification - Insects of Florida*. University of Florida.
- Choate, P. M. (2011c). *Key to the Sub-orders of Hymenoptera* (pp. 1-4). University of Florida.
- Choong, C., Alwen, B., Izzat Husna, M., & Amirrudin, B. (2017). Odonata fauna of Tioman Island, Pahang, Peninsular Malaysia. *Journal of Wildlife and Parks*, 32, 31-40.
- Chung, A. Y. C., Paul, V., & Bosuang, S. (2020). The insect fauna of Tenompok forest reserve in Sabah, Malaysia. *Journal of Threatened Taxa*, 12(4), 15443-15459. <https://doi.org/10.11609/JOTT.5588.12.4.15443-15459>
- Dow, R. A., Reels, G., & Butler, S. (2013). Previously Unpublished Odonata Records from Sarawak, Borneo. *Journal of International Dragonfly Fund*, 6, 1-27.
- Enagbonma, B. J., & Babalola, O. O. (2019). Environmental sustainability: A review of termite mound soil material and its bacteria. *Sustainability (Switzerland)*, 11(14). <https://doi.org/10.3390/su11143847>
- Farizawati, S., Fauzi, M., Ruslan, M. Y., Norma-Rashid, Y., Ng, Y. F., & Idris, A. G. (2014). The Diversity of Odonates in Five Islands within the West Coast of Peninsular Malaysia. *Academic Journal of Entomology*, 7(3), 102-108. <https://doi.org/10.5829/idosi.aje.2014.7.3.855>
- Francini, R. B., & Ramos, R. R. (2014). Butterflies of Alcatrazes Island, São Paulo

- State, Brazil (Lepidoptera: Papilionoidea and Hesperioidea). *Check List*, 10(1), 28-32.
- Gildenkov, M. Y. (2019). Two new species of the genus *carpelimus* leach, 1819 from Malaysia (Coleoptera: Staphylinidae: Oxytelinae). *Russian Entomological Journal*, 28(4), 370-372. <https://doi.org/10.15298/rusentj.28.4.04>
- Greenslade, P. J. M. (1972). Comparative ecology of four tropical ant species. *Insectes Sociaux*, 19(3), 195-212. <https://doi.org/10.1007/BF02226626>
- Gullan, P. J., & Cranston, P. S. (2010). *The Insects: An Outline of Entomology* (4th ed.). Wiley-Blackwell.
- Hor, P. L., Norhazlini, M. Z., Nasihah, M., & Zulhazman, H. (2021). Diurnal insect pollinators of *rafflesia kerri* Meijer at Lojing Highlands, Kelantan, Peninsular Malaysia. *Malaysian Forester*, 84(1), 7-15.
- Jusoh, W. F. A., Ballantyne, L., Chan, S. H., Wong, T. W., Yeo, D., Nada, B., & Chan, K. O. (2021). Molecular systematics of the firefly genus *luciola* (Coleoptera: Lampyridae: Luciolinae) with the description of a new species from Singapore. *Animals*, 11(3), 1-16. <https://doi.org/10.3390/ani11030687>
- Kamarol Zaman, N. A., & Omar, M. A. (2019). Repellent activity of plants extraction against cockroaches (*Periplaneta americana*, *Blattella germanica*). *E-Proceedings of 4th International Conference on Research in TVET Studies (ICOR-TVET 2019)*, 145-147.
- Lewis, S. M., Thancharoen, A., Wong, C. H., López-Palafox, T., Santos, P. V., Wu, C., Faust, L., De Cock, R., Owens, A. C. S., Lemelin, R. H., Gurung, H., Jusoh, W. F. A., Trujillo, D., Yiu, V., López, P. J., Jaikla, S., & Reed, J. M. (2021). Firefly tourism: Advancing a global phenomenon toward a brighter future. *Conservation Science and Practice*, 3(5). <https://doi.org/10.1111/csp.2.391>
- Mat Ripin, M. A. F., & Md. Latip, S. N. H. (2021). *Oryctes rhinoceros* infestation and its interaction with oil palm damage in selected oil palm plantation. *International Transaction Journal of Engineering, Management & Applied Sciences & Technologies*, 12(9), 1-10. <https://doi.org/10.14456/ITJEMAST.2021.187>
- Mohamed, S., Mohd Zamri, K., Zullkeflee, A., Sajili, M. H., Roseli, M., & Mahmud, K. (2019). Insects diversity and abundance in coastal and inland forest of Perhentian Island, Terengganu, Peninsular Malaysia. *Bioscience Research*, 16(1), 2272-2281.
- Mohd-Taib, F. S., David, G., Lee, A., Low, K. H., Lee, K. S., Yusoff, M. F., & Yaacob, A. R. (2019). Checklist of birds in Pulau Pangkor, Perak, Malaysia. *The Malaysian Forester*, 82(1), 217-226.
- Ng, Y. F., & Mound, L. A. (2020). Character state variation among species of Thrips genus (Thysanoptera) in Malaysia, with one new species and two new records. *Zootaxa*, 4852(2), 239-245.
- NRE. (2016). *National Policy on Biological Diversity 2016-2025*. Putrajaya: Ministry of Natural Resource and Environment (NRE).
- Nur Atiqah, A. R., Siti Nurul Amirah, M. D., Hafizzudin, A. R., Salmah, Y., Ng, Y. F., Ting, J. S., Asraf, B., & Zubaid, A. (2017). An assessment of the nocturnal insect diversity and abundance between an agricultural and suburban landscape in Peninsular Malaysia. *Serangga*, 22(1), 274-282.
- Orr, A. G. (2014). Odonata in Bornean tropical rain forest formations: Diversity, endemicy and implications for conservation management. In A. C. Riviera (Ed.). *Forest and Dragonflies*. Sofia: Pensoft Publishers.
- Pimid, M., Ahmad, A. H., & Krishnan, K. T. (2019). Food preferences and foraging

- activity of asian weaver ants, *Oecophylla smaragdina* (Fabricius) (Hymenoptera: Formicidae) *Tropical Life Sciences*, 30(2).
- Pradipta, A. P., Wagiman, F. X., & Witjaksono, W. (2020). The coexistence of *Oryctes rhinoceros* L. and *Xylotrupes gideon* L. (Coleoptera: Scarabaeidae) on immature plant in oil palm plantation. *Jurnal Perlindungan Tanaman Indonesia*, 24(1), 82. <https://doi.org/10.22146/jpti.52582>
- Quek, K. C., Sodhi, N. S., & Liow, L. H. (1999). New records of butterfly species for Pulau Tioman, Peninsular Malaysia. *Raffles Bulletin of Zoology*, 47(6 SUPPL.), 271-276.
- Rákósy, L., & Schmitt, T. (2011). Are butterflies and moths suitable ecological indicator systems for restoration measures of semi-natural calcareous grassland habitats? *Ecological Indicators*, 11(5), 1040-1045. <https://doi.org/10.1016/j.ecolind.2010.10.010>
- Ridzuan, N. H., Farouk, S. A., Razak, S. A., Avicor, S. W., Taib, N., & Hamzah, S. N. (2021). Insect biodiversity of urban green spaces in Penang Island, Malaysia. *International Journal of Tropical Insect Science*, 42, 275-284. <https://doi.org/10.1007/s42690-021-00543-2>
- Romoser, W. S., & Stoffolano, J. G. (1998). *The Science of Entomology* (4th ed.). McGraw Hill.
- Rosmidi, F. H., Zahidin, M. A., Adanan, A., Azizah, A., Pesiu, E., & Abdullah, M. T. (2017). Checklist of butterflies in Pulau Perhentian and Pulau Bidong, Terengganu. *Journal of Sustainability Science and Management*, 12(1), 40-48.
- Saikim, F. H., Le, G., & Dawood, M. M. (2020). Tourists' Perceptions of Insects as the Determinants of Insect Conservation through Entomological Ecotourism. *Journal of Tropical Biology and Conservation*, 17, 79-95.
- Shafiq, H. M., & Maryati, M. (2021). *Dinomyrmex gigas* (Latreille, 1802): A Potential Icon for Taman Negara Johor Endau Rompin. *IOP Conference Proceeding: Earth and Environmental Science*, 736, 012044 <https://doi.org/10.1088/1755-1315/736/1/012044>
- Shuey, J., Labus, P., Carneiro, E., Dias, F. M. S., Leite, L. A. R., & Mielke, O. H. H. (2017). Butterfly communities respond to structural changes in forest restorations and regeneration in lowland Atlantic Forest, Paraná, Brazil. *Journal of Insect Conservation*, 21(3), 545-557. <https://doi.org/10.1007/s10841-017-9994-y>
- Stork, N. E., McBroom, J., Gely, C., & Hamilton, A. J. (2015). New approaches narrow global species estimates for beetles, insects, and terrestrial arthropods. *Proceedings of the National Academy of Sciences of the United States of America*, 112(24), 7519-7523. <https://doi.org/10.1073/pnas.1502408112>
- Tan, M. K., Yeo, Huiqing, & Lee, J. X. Q. (2015). Diversity of entomofauna (Orthoptera, Reduviidae and Aculeata) in the Mandai Lake Road Area, Singapore. *Nature in Singapore*, 8, 37-51.