

BATS OF PULAU TINGGI, JOHOR, MALAYSIA

MOHAMAD HAZARUL SYAFIQ ABDULLAH¹, MOHD-RIDWAN ABD RAHMAN^{2*}, FATIN NUR ATIKAH ZAHARI^{1,3}, NURSYUHADA OTHMAN^{1,3}, HIDAYAH HARIS^{1,3}, MOHD FAUDZIR NAJMUDDIN^{1,3}, MD-ZAIRI ZAINAL⁴ AND MUHAMMAD ABU BAKAR ABDUL-LATIFF^{1,3*}

¹Department of Technology and Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (Pagoh Campus), 84600 Muar, Johor, Malaysia. ²Centre for Pre-University Studies, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia. ³Environmental Management and Conservation Research Unit (eNCORE), Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (Pagoh Campus), 84600 Muar, Johor, Malaysia. ⁴Shaz Resort Pulau Tinggi Sdn. Bhd., Lot 44, Kampung Penaga, Pulau Tinggi, 86800 Mersing, Johor, Malaysia.

*Corresponding authors: armridwan@unimas.my, latiff@uthm.edu.my

Submitted final draft: 22 March 2022

Accepted: 9 April 2022

<http://doi.org/10.46754/jssm.2022.11.007>

Abstract: A series of bat surveys took place in Pulau Tinggi from July 2019 to September 2019. A total of 242 individuals representing 15 species from six families were captured from a cumulative effort of 462 trap nights. The most caught species for frugivorous bats was *Cynopterus brachyotis* (n = 150) while for insectivorous bats, it was *Hipposideros cervinus* (n = 20). Pulau Tinggi has a high density of bat species per area (1.01 sp/km²) when compared to other tourist islands on Peninsular Malaysia's east coast [e.g., Pulau Tioman (0.13 sp/km²), Pulau Perhentian (0.5 sp/km²) and Pulau Redang (0.4 sp/km²)]. As a result, high levels of disturbance such as tourism and habitat loss may negatively affect the diversity of bat species on the island. Because of time constraints and the study did not cover the entire island, information on the diversity of bats in Pulau Tinggi is still inadequate. Long-term monitoring of bat diversity covering forest areas throughout the island needs to be conducted immediately to identify the number of species, their distribution and ecology. This information is important in formulating a strategic plan for bat conservation in Pulau Tinggi.

Keywords: Bat, Chiroptera, small mammal, Malaysia, Johor, Seribuat Archipelago, South China Sea.

Introduction

Bats belong to the order Chiroptera, the second-largest group of mammals after Rodentia (Vaughan *et al.*, 2015). It is an important seed disperser, pollinator and insect population controller in ecosystems (Jones *et al.*, 2009). Malaysia is one of the biodiversity hotspots for bat species, with 143 species accounting for 10% of the world's total bat species (Senawi & Ahmad, 2021; Simmons & Cirranello, 2022). In Peninsular Malaysia alone, 110 bats have been described and mostly identified based on research conducted on the mainland (Lim *et al.*, 2017). Although many bat surveys have been conducted in Malaysia (Naharuddin *et al.*, 2015; Mohd-Hanif *et al.*, 2015; Shazali *et al.*, 2016; Lim *et al.*, 2017; Mohd-Ridwan *et al.*, 2018; Lim *et al.*, 2019; William-Dee *et al.*, 2019;

Senawi *et al.*, 2020; Lok *et al.*, 2021), studies on bat diversity in Peninsular Malaysia's offshore islands are poorly understood and not well documented (Roslan *et al.*, 2016).

Reports on the diversity of Malaysian bats on the islands continue to be limited by effort and funding due to its costly and unpredictable climatic events (Baqi *et al.*, 2021). Previously, offshore island bat diversity surveys, particularly in the South China Sea, took place on Pulau Bidong, Pulau Perhentian (Roslan *et al.*, 2016; Baqi *et al.*, 2021), Pulau Redang (Mohd-Yusof *et al.*, 2019) and Pulau Tioman (Medway, 1966; DWNP, 1995; Csorba *et al.*, 1997; Lim *et al.*, 1999). Pulau Tioman has the most bat species (19 species) while Pulau Bidong has the fewest (three species). According to those surveys, larger islands closer to the mainland had higher

rates of bat diversity than smaller islands with a greater distance from the mainland. The findings support the theory of island biogeography, which states that the richness of a species depends on the size and isolation of the island (MacArthur & Wilson, 2016; Valente *et al.*, 2020). Larger islands have a greater species richness than smaller islands due to abundant resources. However, as the island becomes more geographically isolated from the mainland and smaller, its species richness will decline (MacArthur & Wilson, 2016; Valente *et al.*, 2020).

Habitat loss and human disturbance are major threats island bats face (Jones *et al.*, 2009). The booming tourism industry across Malaysia's islands has put significant pressure on natural resources and land use in the offshore islands. Therefore, extensive scientific surveys and studies must be conducted to determine the impact of development on species diversity, distribution and ecology of island bats. To date, no island bat surveys have occurred on Pulau Tinggi. This survey aims to provide baseline data on bat species in Pulau Tinggi. The results are compared to previous studies conducted on Malaysia's east coast islands. The data gathered is crucial in assisting authorities in developing a

strategic plan and policies to improve island bat conservation, particularly in Pulau Tinggi.

Materials and Methods

Study Site

The study occurred in Pulau Tinggi (2° 18' N, 104° 07' E), located near Mersing, about 30 km to the southeast of Johor. Pulau Tinggi is located in the district of Mersing in Johor (Figure 1). There are over 19 villages there (Manaf *et al.*, 2011). Pulau Tinggi and other 12 small islands have been listed under East Johor Island Archipelagos (EJIA), where Pulau Tinggi is considered the biggest island among the rest (Azman, 2008). The other 12 islands under EJIA are Pulau Mensirip, Pulau Goal, Pulau Harimau, Pulau Rawa, Pulau Tengah, Pulau Hujong, Pulau Mentigi, Pulau Sibul Hujong, Pulau Sibul, Pulau Besar, Pulau Aur and Pulau Pemanggil.

Field Methods, Identification, Processing and Preservation

Field samplings were done using ground netting, high pole netting, and harp trapping (Mohd-Ridwan *et al.*, 2018). A total of ten mist nets, five high poles and six harp traps were placed throughout several bat flyways, including above




Figure 1: Map showing the location of Pulau Tinggi in Peninsular Malaysia

streams, trails, along narrow paths, in cleared parts of the forest and at the forest edge. The total number of nets and traps multiplied by the number of sampling days was used to evaluate the amount of trapping effort (Mohd-Ridwan *et al.*, 2018). Bats were identified based on Kingston *et al.* (2006) and Francis *et al.* (2008). The morphological measurements of the bats were taken using a digital vernier calliper and weighted using Pesola spring balance (Mohd-Ridwan *et al.*, 2018). Following Kunz (1988), adult bats were determined by examining the epiphyseal-diaphyseal fusion on the third, fourth and fifth metacarpals. To avoid recapture and lessen stress, all trapped bats were kept until all nets were closed (a maximum of 12 hours). Selected voucher specimens were euthanized using chloroform and subjected to skinning procedures. The liver and pectoral muscle were excised and preserved in 75% ethanol. The samples were then deposited at the Universiti Tun Hussein Onn (UTHM) zoological museum.

Data Analysis

The total number of species caught was used to calculate the species richness. The relative abundance was estimated by the number of individuals per species divided by the number of individuals. The Shannon-Wiener diversity index (H') was used to estimate species diversity in Pulau Tinggi, sensitive to changes in the abundance of rare species in a community (Solow, 1993). A diversity t-test was conducted to compare the abundance and composition of bat assemblages between different trapping methods. The Shannon-Wiener index and diversity t-test analysis was calculated using PAST software (Hammer *et al.*, 2001). The phenogram was generated to compare the species richness of bats documented at four east coast islands of Peninsular Malaysia. The cluster analysis comparisons were based on the absence and presence matrix of species documented in all the localities (Tingga *et al.*, 2012). The selected sites include Pulau Perhentian (Tambryn *et al.*, 2005; Roslan *et al.*, 2016), Pulau Bidong (Roslan *et al.*, 2016), Pulau Redang (Mohd-

Yusof *et al.*, 2019) and Pulau Tioman (Medway, 1966; DWNP, 1995; Csorba *et al.*, 1997; Lim *et al.*, 1999). A Venn diagram was produced to determine the shared and unique bat species compared to other east coast islands.

Results

A total of 242 individuals representing 15 species from six families: Emballonuridae, Hipposideridae, Megadermatidae, Pteropodidae, Rhinolophidae and Vespertilionidae were recorded at Pulau Tinggi (Table 1). Pteropodidae has the greatest number of species (five species) on this island, followed by Hipposideridae (four species) and Vespertilionidae (three species), whereas Rhinolophidae, Emballonuridae and Megadermatidae each with one species. A total of 137 individuals representing nine species (220/net-night) were mist-netted, 24 individuals (five species) were captured in harp traps (132/trap-night) and 81 individuals (eleven species) were caught using a high pole (110/net-night). *Cynopterus brachyotis* (62%) was the most captured species in Pulau Tinggi followed by *Eonycteris spelaea* (9.5%). Four species were recorded as singletons: *Hipposideros galeritus*, *Taphozous melanopogon*, *Megaderma spasma* and *Myotis muricola*.

Comparison of Trapping Methods

All three trapping methods were successful in capturing bats (Table 2). The high pole nets recorded the highest number of bat species (11 species), followed by mist nets (nine species) and harp traps (five species). Family Pteropodidae showed the highest overall captured rate for both high poles (trapping effort/nights = 0.64) and mist nets (trapping effort/nights = 0.58). Family Hipposideridae (trapping effort/nights = 0.17) recorded the highest capture rate using harp traps. There was no significant difference in the species and relative abundance of bats captured using mist nets and harp traps ($t = 0.64$, $df = 38$, $p = 0.52$), mist nets and high poles ($t = 0.61$, $df = 154$, $p = 0.54$) and between high pole and harp traps ($t = 1.02$, $df = 55$, $p = 0.30$).

Table 1: Taxonomic list of bats species captured with their average measurements (minimum and maximum ranges) from both study areas

No.	Family Species	N	Forearm Mean (range) (mm)	Ear Mean (range) (mm)	Hindfoot Mean (range) (mm)	Tail Mean (range) (mm)
Pteropodidae						
1.	<i>Pteropus hypomelanus</i>	12	124.17 (116-145)	25.58 (24-27)	40.33 (34-45)	NA
2.	<i>Cynopterus brachyotis</i>	150	60.29 (52-66)	16.24 (14-17)	12.89 (10-15)	8.78 (8-9)
3.	<i>Penthetor lucasi</i>	7	61.86 (57-65)	15 (11-17)	14.86 (12-17)	8 (6-10)
4.	<i>Eonycteris spelaea</i>	23	61.22 (55-70)	16.78 (15-17)	15 (15-18)	11.96 (9-14)
5.	<i>Macroglossus minimus</i>	7	38.86 (37-42)	13.43 (11-15)	11.57 (10-13)	
Rhinolophidae						
6.	<i>Rhinolophus stheno</i>	4	47.25 (43-50)	17 (15-20)	11.5 (11-13)	17.75 (16-20)
Hipposideridae						
7.	<i>Hipposideros cervinus</i>	20	45.55 (45-47)	14.3 (12-15)	7.3 (6-9)	23.5(22-27)
8.	<i>Hipposideros diadema</i>	9	82.22 (77-90)	25 (23-30)	14.45 (10-16)	41 (39-49)
9.	<i>Hipposideros galeritus</i>	1	50	10	7	30
10.	<i>Hipposideros bicolor</i>	2	44.5	16.5	5.5	28.5
Megademartidae						
11.	<i>Megaderma spasma</i>	1	60	32	14	NA
Emballonuridae						
12.	<i>Taphozous melanopogon</i>	1	60	15	12	23
Vespertilionidae						
13.	<i>Miniopterus medius</i>	2	37 (32-42)	9.5 (7-12)	8 (7-9)	30.5(29-32)
14.	<i>Philetor brachypterus</i>	2	32.8 (32-33.5)	11.12 (9.4-12.45)	6.9 (5-8)	33.54 (32.08-35.1)
15.	<i>Myotis muricola</i>	1	34	11	7	40

Table 2: List of species caught at Pulau Tinggi using three different methods including mist nets, harp traps and high-poles

No.	Family Species	Harp Tarp	Mist Net	High Pole	RA (%)	Total	IUCN (2022)	DWNP (2017)
Pteropodidae								
1.	<i>Pteropus hypomelanus</i>	-	12	-	5	12	NT	LC
2.	<i>Cynopterus brachyotis</i>	-	94	56	62	150	LC	LC
3.	<i>Penthetor lucasi</i>	-	3	4	2.9	7	LC	LC
4.	<i>Eonycteris spelaea</i>	-	17	6	9.5	23	LC	LC
5.	<i>Macroglossus minimus</i>	-	3	4	2.9	7	LC	DD
Rhinolophidae								

6.	<i>Rhinolophus stheno</i>	-	2	2	1.7	4	LC	LC
Hipposideridae								
7.	<i>Hipposideros cervinus</i>	16	-	4	8.3	20	LC	LC
8.	<i>Hipposideros diadema</i>	5	3	1	3.7	9	LC	LC
9.	<i>Hipposideros galeritus</i>	1	-	-	0.4	1	LC	LC
10.	<i>Hipposideros bicolor</i>	1	-	1	0.8	2	LC	NT
Megademartidae								
11.	<i>Megaderma spasma</i>	1	-	-	0.4	1	LC	LC
Emballonuridae								
12.	<i>Taphozous melanopogon</i>	-	-	1	0.4	1	LC	LC
Vespertilionidae								
13.	<i>Miniopterus medius</i>	-	2	-	0.8	2	LC	DD
14.	<i>Philetor brachypterus</i>	-	1	1	0.8	2	LC	DD
15.	<i>Myotis muricola</i>	-	-	1	0.4	1	LC	LC
No. of individuals		24	137	81		242		
No. of species		5	9	11		15		
Sampling effort		132	220	110				
Capture rate		0.18	0.62	0.74				
Shannon Index (H')		0.99	1.14	1.25				

Note: RA=Relative Abundance, LC=Least Concern, DD=Data Deficient, NT=Near Threatened

Species Diversity Indices

The number of bat species captured in Pulau Tinggi increased exponentially till the 10th night (Figure 2). Starting from the 11th night, the species accumulation curve hit an asymptote,

indicating sampling saturation. Species diversity index ($H' = 1.47$) showed a comparatively high diversity of bat species in Pulau Tinggi (Table 3) in comparison to other east coast islands (e.g., Pulau Perhentian and Pulau Bidong; Roslan *et al.*, 2016).




Figure 2: Species cumulative curve of bats captured at Pulau Tinggi

Comparison with Other Malaysian East Coast Islands

Our findings are being compared with other east coast islands, Pulau Perhentian, Pulau Bidong, Pulau Redang and Pulau Tioman (Table 3). 29 species of bats can be found throughout the east coast islands. Pulau Tioman has the highest number of bats (19 species), followed by Pulau Tinggi (15 species), Pulau Perhentian (nine species), Pulau Redang (eight species) and Pulau Bidong (three species). Three species of bats

were distributed throughout east coast islands, namely *Pteropus hypomelanus*, *Cynopterus brachyotis* and *Megaderma spasma* (Figure 2). Eight and nine species are unique to the Pulau Tinggi and Pulau Tioman, respectively (Table 3, Figure 2). The comparisons of four different sites in cluster analysis resulted in two groups (Figure 3 and 4). Pulau Perhentian clustered together with Pulau Redang, followed by Pulau Bidong and Pulau Tioman. Pulau Tinggi is the most distant group to other east coast islands.

Table 3: Taxonomic comparison with other Malaysian east-coast islands

No.	Family Species	This Study	Pulau Perhentian ^{a,b}	Pulau Bidong ^a	Pulau Redang ^{c,d}	Pulau Tioman ^{e,f,g,h}
Pteropodidae						
1.	<i>Pteropus hypomelanus</i>	+	+	+	+	+
2.	<i>Cynopterus brachyotis</i>	+	+	+	+	+
3.	<i>Cynopterus horsfieldii</i>					+
4.	<i>Penthetor lucasi</i>	+				
5.	<i>Eonycteris spelaea</i>	+	+		+	+
6.	<i>Macroglossus minimus</i>	+				
Rhinolophidae						
7.	<i>Rhinolophus affinis</i>		+		+	+
8.	<i>Rhinolophus lepidus</i>		+			+
9.	<i>Rhinolophus luctus</i>					+
10.	<i>Rhinolophus macrotis</i>					+
11.	<i>Rhinolophus megaphyllus</i>					+
12.	<i>Rhinolophus pusillus</i>					+
13.	<i>Rhinolophus stheno</i>	+				+
14.	<i>Rhinolophus borneensis</i>					+
Hipposideridae						
15.	<i>Aselliscus stoliczkanus</i>					+
16.	<i>Hipposideros bicolor</i>	+	+			+
17.	<i>Hipposideros cervinus</i>	+				
18.	<i>Hipposideros cineraceus</i>		+		+	
19.	<i>Hipposideros diadema</i>	+				
20.	<i>Hipposideros galeritus</i>	+				
21.	<i>Hipposideros larvatus</i>		+			+

Megadermatidae					
22.	<i>Megaderma spasma</i>	+	+	+	+
Nycteridae					
23.	<i>Nycteris tragata</i>				+
Emballonuridae					
24.	<i>Emballonura monticola</i>			+	+
25.	<i>Taphozous melanopogon</i>	+		+	
Vespertilionidae					
26.	<i>Miniopterus medius</i>	+			
27.	<i>Philetor brachypterus</i>	+			
28.	<i>Myotis muricola</i>	+			
Mollosidae					
29.	<i>Cheiromeles torquatus</i>				+
No. of family		6	4	2	4
No. of species		15	9	3	8
No. of individuals		242	35	84	Na
Sampling effort		462	80	80	Na
Capture rate		0.52	0.44	1.05	Na
Diversity index (H')		1.47	1.95	0.59	Na

^aRoslan *et al.*, 2016, ^bTambryn *et al.*, 2005, ^cMNS 1990, ^dMohd-Yusof, ^eLim *et al.*, 1999, ^fCsorba *et al.*, 1997, ^gMedway 1966, ^hDWNP 1995. *Unique for the respective island




Figure 3: The Venn diagram illustrates the number of shared and unique bat species across east-coast islands




Figure 4: Phenogram of the comparison between Pulau Tinggi with other bats survey records in east-coast islands of Peninsular Malaysia

Discussion

Species Richness and Abundance

This study contributes 15 species, accounting for approximately 13.6% of the bats in Peninsular Malaysia (Lim *et al.*, 2017). A short-nosed fruit bat, *C. brachyotis* was frequently captured during the study period. Similarly, with high capture rates, this species is found on all east coast islands (e.g., Pulau Perhentian and Pulau Bidong) (Roslan *et al.*, 2016; Baqi *et al.*, 2021). This adaptable species inhabits a wide range of habitats on the mainland and islands (Mohd-Ridwan *et al.*, 2011; 2018). Trap traps may have contributed to the richness and abundance of frugivores bats species on Pulau Tinggi. Most species caught in the harp trap are insectivorous bats from the family Hipposideridae. *Hipposideros cervinus* is the most caught insectivorous species during this survey. In Borneo, this species moves in large groups resulting in high numbers of individuals caught in a single harp trap (Mohd-Ridwan *et al.*, 2018).

The year-round availability of fruits and the long fruiting season may be important factors in sustaining a frugivorous bat population in the tropics (Fleming, 1988). In Pulau Tinggi, our study site is surrounded by locally-owned fruit orchards and small-scale banana plantations (*Musa sapientum*). Among the crops grown by the villagers are durian (*Durio zibethinus*),

rambutan (*Nephelium lappaceum*), guava (*Psidium quajava*), jackfruit (*Artocarpus heterophyllus*) and mango (*Mangifera indica*). Frugivore bats like *P. hypomelanus*, *C. brachyotis*, *E. spelaea*, *P. lucasi* and *M. mininus* are likely to affect forest regeneration and plant dispersal across the island. At night, large groups of *P. hypomelanus* visited durian flowers and fig trees. *Pteropus hypomelanus* is an important seed disperser and pollinator, particularly for durians (Aziz *et al.*, 2017).

Species Checklist

Species composition of bats in Pulau Tinggi (15 species) seems to be comparatively lower than Pulau Tioman (19 species) but relatively higher to other east coast offshore islands of Peninsular Malaysia such as Pulau Perhentian (nine species), Pulau Redang (eight species) and Pulau Bidong (three species). Size of an island and its distance to the mainland influence the diversity of organisms (Valente *et al.*, 2020). The size of Pulau Tioman is ten times bigger (136 km², 32 km to the mainland) than Pulau Tinggi (14.85 km², 20 km to the mainland). Even though the species richness of bats at Pulau Tioman is higher than Pulau Tinggi, the rate of bats per area is considered lower (0.13 sp/km²) than Pulau Tinggi (1.01 sp/km²). Meanwhile, Pulau Perhentian (15.35 km², 19 km to the mainland), Pulau Redang (19.09 km², 25 km to the mainland) and Pulau Tinggi are considerably

similar in terms of size and their distance to the mainland. However, the richness and the rate of bats per area for Pulau Perhentian (0.5 sp/km²) and Pulau Redang (0.4 sp/km²) are less than Pulau Tinggi. The level of anthropogenic activity also affects the diversity of insular bats (Wiles & Brooke 2010). This is evident on well-known tourist islands such as Pulau Perhentian, Pulau Redang and Pulau Tioman, where many forested areas have been converted into chalets and resort buildings (Weng, 2009).

According to the IUCN (2022), 14 species of Pulau Tinggi were classified as Least Concern, with only one species classified as Near Threatened. However, some species are included in several conservation categories at the national level (DWNP, 2017). Four bat species are globally classified as Least Concern (IUCN, 2022) and at the national level, they are listed as Data Deficient and Near Threatened (DWNP, 2017). Bats are experiencing more serious threats nationally, and local conservation actions are urgently required. The main threat to the bat community on Pulau Tinggi is habitat loss due to deforestation. Some forest areas on Pulau Tinggi have been developed into settlements, chalets, and resorts. However, compared to other tourist islands, the rate of land use is still low because development is concentrated in coastal areas. Most of the forested areas on Pulau Tinggi are still intact and unaffected by human activity.

Conclusion

This study successfully provided a checklist of bats in Pulau Tinggi. A total of 15 species from five families were successfully recorded during the seven-week study on the island. Given the short study period and the fact that it does not cover the entire island region, information on the bat community in Pulau Tinggi is still incomplete. Long-term monitoring of unexplored forest areas on Pulau Tinggi is required to determine the number of species, distribution and ecology of bats on the island. Furthermore, this information contributes to a better understanding of how bats react to changes in the island's landscape. The research findings

can also be adapted to the tourism module and used for awareness programs and conservation activities in Pulau Tinggi.

Acknowledgements

This project is funded by the Ministry of Higher Education Malaysia (MOHE) under the Malaysian Technical University Network (MTUN) grant scheme Vote K121 and Industrial Grant by Shaz Resort Sdn. Bhd. (UTHM-SHAZ-M004), both grants awarded to Associate Professor Ts. Dr. Muhammad Abdul Latiff Abu Bakar (UTHM) as the Principal Researcher. This research was supported by GPPS-UTHM-2018-H288 postgraduate grant from Universiti Tun Hussein Onn Malaysia (UTHM), Malaysia. We are grateful to Dato' Abdul Kadir Abu Hashim, Director General of the Department of Wildlife and National Parks who provided us with the necessary facilities and assistance. This research was conducted under a research permit (JPHL & TN (IP): 600-6/1/4 (03) / M-00351-15-19). We are deeply indebted to the Department of Wildlife and National Parks Malaysia for granting permission to carry out this research. The authors acknowledge the Ministry of Higher Education Malaysia, Universiti Tun Hussein Onn Malaysia and Shaz Resort Sdn. Bhd. for providing necessary funding, facilities and assistance.

References

- Abdullah M. T., Hall, L. S., Khan, F. A. A & Azhar, M. (2022). *Photographic guide to the bats of Malaysia*. Penerbit UMT, Kuala Nerus.
- Aziz, S. A., Clements, G. R., McConkey, K. R., Sritongchuay, T., Pathil, S., Abu Yazid, M. N. H., Campos-Arceiz, A., Forget, P. M. & Bumrungsri, S. (2017). Pollination by the locally endangered island flying fox (*Pteropus hypomelanus*) enhances fruit production of the economically important durian (*Durio zibethinus*). *Ecology and Evolution*, 7(21), 8670-8684.

- Azman, B. A. R., Ramlan, O., Wan-Lotfi, W. M., Zaidi, C. C., & Othman, B. H. R. (2008). Seagrass biodiversity of Pulau Tinggi, Johor. Malaysia marine ecosystem: The studies of Johor Darul Takzim East Coast. *Research and Information Series of Malaysian Coasts, Malaysia*, 1-176.
- Baqi, A., Azhar, I., Chen, E. W., Khan, F. A. A., Lian, C. J., Nelson, B. R., & Kumaran, J. V. (2021). The diversity of small mammals in Pulau Perhentian Kecil, Terengganu, Malaysia. *Journal of Threatened Taxa*, 13(6), 18427-18440.
- Csorba, G. T., Fuisz, T., & Kelen, B. (1997). New birds and bats from Pulau Tioman, Malaysia. *Malayan Nature Journal*, 50, 197-200.
- DWNP. (1995). *Inventori kepelbagaian biologi Rezab Hidupan Liar Pulau Tioman. Pahang Darul Makmur*. Biokepelbagaian mamalia di Pulau Tioman. 27-38. Unpublished report.
- DWNP. (2017). *Red list of mammals for Peninsular Malaysia* (Version 2). Department of Wildlife and National Parks (PERHILITAN) Peninsular Malaysia, Kuala Lumpur.
- Fleming, T. H. (1988). The short-tailed fruit bat. University of Chicago Press, Chicago.
- Francis, C. M. (2008). A guide to the mammals of Southeast Asia. New Holland Publishers United Ltd., United Kingdom.
- Hammer, O., Harper, D. A., & Ryan, P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*, 4(1), 1-9.
- IUCN. (2022). *Red List of Threatened Species*. www.iucnredlist.org
- Jones, G., Jacobs, D. S., Kunz, T. H., Willig, M. R., & Racey, P. A. (2009). Carpe noctem: The importance of bats as bioindicators. *Endangered Species Research*, 8(1-2), 93-115.
- Kingston, T., Lim, B. L. & Zubaid A. (2006). *Bats of Krau Wildlife Reserve*. Penerbit Universiti Kebangsaan Malaysia, Bangi.
- Kunz, T. H., & Kurta, A. (1988). Capture methods and holding devices. In Thomas H. Kunz & Stuart Parsons (Eds.), *Ecological and Behavioural Methods for the Study of Bats*. Smithsonian Institution Press, Washington.
- Lim, B. L., Kelvin, K. P., & Yong, H. S. (1999). The terrestrial mammals of Pulau Tioman, Peninsular Malaysia, with a catalogue of specimens at the Raffles Museum, National University of Singapore. *Raffles Bulletin of Zoology*, 6, 101-124.
- Lim, L. S., Struebig, M. J., Zalipah, M. N., Mohd-Adnan, A., Senawi, J., Zubaid, A., Sah, S. A. M. & Rossiter, S. J., (2019). Bats from the understorey of lowland tropical rainforests across Peninsular Malaysia. *Journal of Bat Research and Conservation*, 12, 68-82.
- Lim, V. C., Ramli, R., Bhassu, S., & Wilson, J. J. (2017). A checklist of the bats of Peninsular Malaysia and progress towards a DNA barcode reference library. *PLOS ONE*, 12(7), e0179555.
- Lok, Y. C., Siau, V. G., Mohd Kanapiah, N. A. A., Lai, T. C., Haslan, N. N. H. H., Nukili, N. N., Daud, U. S., Amat, A., Gompoyo, J., Tuh, Y. Y. F. & Hasan, N. H. (2021). Bat species diversity trend along an elevation gradient: A study in Crocker Range Park, Sabah, Borneo. *Biodiversity Data Journal*, 9, e72651.
- MacArthur, R. H., & Wilson, E. O. (2016). *The theory of island biogeography*. Princeton University Press.
- Manaf, A. A., Shamsuddin, M., & Omar, M. (2011). Marine Park GAZETtement Impacts on Pulau Tinggi Community in Mersing, Johor. *Journal of Tropical Marine Ecosystem*, 1(1), 35-43.
- Medway, L. (1966). The mammals. Observations on the fauna of Pulau Tioman and Pulau

- Tulai. *Bulletin of the Singapore National Museum*, 34, 9-32.
- Mohd-Hanif, R. M. D., Nur-Aida, M. T., Zahirunisa, A. R., Mohd-Ridwan, A. R. & Abdullah, M. T., (2015). Contribution of regenerated forest in conservation of bats in Peninsular Malaysia. *Journal of Tropical Forest Science*, 27(4), 506-516.
- Mohd-Ridwan, A. R., Tingga, R. C. T., Azhar, M. I., Hassan, N. H., & Abdullah, M. T. (2011). Bats of the wind cave nature reserve, Sarawak, Malaysia Borneo. *Tropical Natural History*, 11, 159-175.
- Mohd-Ridwan, A. R., Tahir, N. F., Eshak, M. H., Csorba, G., Görföl, T., Khan, F. A. A., & Mohd-Azlan, J. (2018). Bats assemblage and lunar phase effect on bat activity at mixed dipterocarp forest, Gunung Gading National Park, Sarawak, Borneo. *Sains Malaysiana*, 47(7), 1349-1357.
- Mohd-Yusof, N., Senawi, J., Nor, S. M., & Md-Zain, B. M. (2019). Haplotype and network analysis of island flying fox (*Pteropus hypomelanus*) using D-loop region of mitochondrial DNA to confirm subspecies designation. *Mammal Research*, 65(2), 375-385.
- Naharuddin, N. M., Shazali, N., Libar, R., Karim, N. F., Mohd-Ridwan, A. R., Roslan, A., Azhar, I., Khalik, M. Z. and Khan, F. A. A., (2015). Bats of Bako National Park and additional notes on the rare partial albinism in fawn roundleaf bat (*Hipposideridae: Hipposideros cervinus*). *Borneo Journal of Resource Science and Technology*, 5(2), 44-52.
- Roslan, A., Gertrude, D., Nur-Izzah, I. A., Aisyah, N., Rahim, A., Elizabeth, P., Nor-Zalipah, M. & Mohd-Tajuddin, A. (2016). Notes of Bats in Pulau Bidong and Pulau Perhentian Besar, Terengganu, Malaysia. *Journal of Sustainability Science and Management Special Issue Number 1: The International Seminar on the Straits of Malacca*, 26-35.
- Senawi, J. & Ahmad, N. (2021). *Kelawar Malaysia: Malaysia Biodiversity Information System MyBIS*. Kementerian Tenaga dan Sumber Asli. Kuala Lumpur, Malaysia.
- Senawi, J., Mahyudin, A., Daud, U. S., Amat, A., Lagundi, S., Gondilang, E., Sutail, E., Narimin, S. & Azhar, I., 2020. Bat diversity in Imbak canyon conservation area: Note on their echolocation calls and ectoparasites. *Journal of Tropical Biology & Conservation*, 17, 217-232.
- Shazali, N., Khalik, M. Z., Rahman, S. H., Azhar, I., Khan, F. A. A., Tahir, N. A., Murni, R., McArthur, E., Latip, N. A., Rahman, M. R. A. & Naharuddin, N., (2016). Small mammals from Miri, Northeastern region of Sarawak, Malaysian Borneo: Note on new locality records. *Check List*, 12(2), 1863.
- Simmons, N. B., & A. L. Cirranello. (2022). *Bat Species of the World: A taxonomic and geographic database*. Accessed on 21 March 2022.
- Solow, A., Polasky, S., & Broadus, J. (1993). On the measurement of biological diversity. *Journal of Environmental Economics and Management*, 24(1), 60-68.
- Tamblyn, A., Turner, C., O'Malley, R., Weaver, N., Hughes, T., Hardingham, S., & Roberts, H. (2005). *Malaysia tropical forest conservation project report of the Perhentian phase 2005*. Final Report. Coral Cay Conservation, London.
- Tingga, R. C. T., Anwarali, F. A., Mohd-Ridwan, A. R., Senawi, J., & Abdullah, M. T. (2012). Small mammals from Kuala Atok, Taman Negara Pahang, Malaysia. *Sains Malaysiana*, 41(6), 659- 669.
- Valente, L., Phillimore, A. B., Melo, M., Warren, B. H., Clegg, S. M., Havenstein, K., Tiedemann, R., Illera, J. C., Thebaud, C., Aschenbach, T. & Etienne, R. S. (2020). A simple dynamic model explains the diversity of island birds worldwide. *Nature*, 579, 92-96.

- Vaughan, T. A., Ryan, J. M., & Czaplewski, N. J. (2015). *Mammalogy* (6th ed.). Jones & Bartlett, Burlington.
- Weng, C. N. (2009). Ecotourism and environmental conservation in small islands in the East Coast of Peninsular Malaysia. *Malaysian Journal of Environmental Management*, 10(2), 53-69.
- Wiles, G. J., Brooke, A. P., Fleming, T. H., & Racey, P. A. 2009. Conservation threats to bats in the tropical Pacific islands and insular Southeast Asia. In *Island bats: Evolution, ecology, and conservation* (pp. 405-459). University of Chicago Press, Chicago, Illinois.
- William-Dee, J., Khan, F. A. A., Rosli, Q., Morni, M. A., Azhar, I., Lim, L. S., Tingga, R. C. T. and Rahman, M. R. A. (2019). Comparative distribution of small mammals diversity in protected and non-protected area of Peninsular Malaysia. *Tropical Life Sciences Research*, 30(2), 131-147.