

## CHLOROPHYTA MICROALGA DIVERSITY IN MESAT RIVER LUBUK LINGGAU CITY, SOUTH SUMATERA PROVINCE, INDONESIA

SEPRIYANINGSIH, SEPRIYANINGSIH<sup>1</sup>, HARMOKO, HARMOKO<sup>1\*</sup> AND NUR HAYATI<sup>2</sup>

<sup>1</sup>University PGRI Silampari, Mayor Toha Street, Lubuklinggau City, South Sumatera Province, Indonesia 31628. <sup>2</sup>University Hasyim Asy'ari Tebuireng Jombang, Jombang Regency, East Java Province, Indonesia 61471.

\*Corresponding author: [putroharmoko@gmail.com](mailto:putroharmoko@gmail.com)

Submitted/final draft: 8 November 2022 Accepted: 20 January 2023

<http://doi.org/10.46754/jssm.2023.03.004>

**Abstract:** The condition of the Mesat River in Lubuk Linggau City, Indonesia is not good, considering the large amount of waste from household activity. The purpose of this study is to analyse the diversity of microalgae in the Mesat River in Lubuklinggau City. This research paper used a survey method and was carried out between February and April 2020. Microalgae samples were collected at three stations along the Mesat River. Each station consisted of three water sampling points, to wit the left, middle, and right of the river. Water samples were taken three times, with a span of one week. Microscopic observation of the resulting microalgae for clear qualitative analysis was conducted. The results of the study found Chlorophyta microalgae which consisted of 3 classes, 8 orders, 12 families, 15 genera and 19 species. Diversity index ranges between 0.98 and 1.27, which indicated that Chlorophyta is in the low diversity category. The evenness index value of Chlorophyta being 0.36 indicates that the community is in a depressed condition but there is no dominance because the average value of the dominance index is 0.011. Abiotic factors measured in the Mesat River in this category is still reasonable and in accordance with second-class water quality standards. Overall, green algae diversity in the Mesat River was good, although the level of diversity was low.

Keywords: Chlorophyta, microalgae, diversity, Mesat River.

### Introduction

The Mesat River is located at Lubuk Linggau City in the South Sumatra Province of Indonesia has a length of approximately 10 kilometres with a width of between three and four metres and a depth of between one and two metres, starting from the Mesat Jaya and Mesat Seni villages and ending at the Wirakarya Village. The flow of the Mesat River is influenced by the amount of household and industrial waste from the tofu and tempeh making process. The condition of the Mesat River will appear clear if the water flow is less but if the water discharge is more, one will notice a lot of garbage in the river and the water is noticeably cloudy.

Lubuk Linggau City has several rivers, including the Kelingi River, Mesat River, Malus River, Kati River, and Kasie River. Water pollution is the ingress and functioning of organisms, substances, energy, or other constituents of the air that degrades water quality and does not meet its designation

(Government Regulation of the Republic of Indonesia Number 82 of 2001). Pollutants include pesticides, artificial fertilisers, garbage, oil spills and detergents (Thyagaraju, 2016; Bat *et al.*, 2018). In addition to affecting water quality, it will also cause the ecosystem in these waters to be disturbed.

Microalgae can be found in fresh water and seawater, and are included in the types of photoautotroph living things (Mourelle *et al.*, 2017; Coêlho *et al.*, 2019). Algae acts as one of the ecological parameters that can provide an overview of water conditions and includes important biotic components in the metabolism of water bodies because they are primary producers in the food chain of aquatic ecosystems (Prata *et al.*, 2019; Rasal *et al.*, 2019). Chlorophyta has an important role in public waters as the main producers or phytoplankton. Chlorophyta converts inorganic compounds into organic compounds through photosynthesis thus becoming a food source for zooplankton and fish

larvae (Kaparapu, 2018; Lomartire *et al.*, 2021). Some research has been done on Chlorophyta microalgae in the Kelingi River (Harmoko & Sepriyaningsih, 2018) and in the Kasie River (Harmoko & Sepriyaningsih, 2020). A study on microalgae in the Mesat River was carried out by Harmoko *et al.* (2018), but is only limited to Lubuk Linggau Timur II District which found 17 Chlorophyta species. The purpose of this study was to analyze the diversity of microalgae in the Mesat River in Lubuklinggau City.

## Materials and Methods

### Research Location and Time

The study was conducted on the Mesat River in Lubuk Linggau Timur I District during February–April 2020. Microalgae samples were taken at three different stations depending on the river flow. The sampling location is shown in Figure 1.

Determination of stations refers to the research results of Goldyn and Kippen (2017) while various microhabitats offer different environmental conditions. Thus, we tested the effect of different habitat types typically found within small ponds on the microalgae

and zooplankton communities. We found that submerged macrophytes have the strongest impact on microalgae and zooplankton communities out of all the analysed habitats. Some epontic diatoms (e.g., *Fragilaria dilatata*, *Cymbella affinis*, which was based on differences in the environment around the river, Station 1 is near a bridge which has shallow water characteristics, Station 2 is near houses that has characteristics of deep water with garbage along the water flow, and Station 3 is near a rubber plantation, which has a tree-covered water flow. Sampling was carried out three times.

### Sample Collection

Sampling was carried out in the morning, one station consists of three points, namely the right bank, left bank, and the middle of the river. The water quality parameters measured consisted of temperature (stem thermometer, acidity (pH meter Risantec brand), dissolved oxygen (DO meter Lutron DO-5510), and water clarity (Secchi disk). Water samples were taken using a plankton net with a mesh size of 20, to filter 5 L of river water. 10 ml of stored water was then transferred to a bottle and four drops of alcohol were added.



Figure 1: Sampling locations on the Mesat River Lubuklinggau City, South Sumatera Province, Indonesia

### Data Collection

The observation of river water samples using an Olympus CX22 binocular microscope with 400× magnification was done at the University PGRI Silampari Biology Laboratory. Observations were made by taking water samples with a dropper pipette (one drop of about 0.05 ml) and made up to five observations. The Chlorophyta samples obtained were then photographed for further identification and analysis. Identification of Chlorophyta microalgae species was done with reference to a book (Wehr & Sheath, 2003; Bellinger & Sigeo, 2010).

### Data Analysis

The diversity index was calculated using the Shannon-Wiener formula (Odum, 1971), namely:

$$H' = - \sum_{i=1}^s p_i \ln p_i$$

Information:

$H'$  = diversity index

$p_i$  =  $n_i/N$

$n_i$  = number of species  $i$

$N$  = total number of species

$S$  = number of species in the sample

The criteria for diversity are as follows:  $H' \leq 2$ : "Small diversity",  $2 < H' \leq 3$ : "Medium diversity", and  $H' > 3$ : "High diversity".

The dominance index is calculated according to (Odum, 1971) as follows:

$$C = \sum_{i=1}^s (p_i)^2$$

Information:

$C$  = dominance index

$p_i$  = proportion of the  $i$ th species

$S$  = number of species found

The dominance index values were grouped into three criteria, namely:  $0 < C \leq 0.5$ : "Low dominance",  $0.5 < C \leq 0.75$ : "Medium dominance", and  $0.75 < C \leq 1$ : "High dominance".

The evenness index was calculated according to (Odum, 1971), as follows:

$$E = \frac{H'}{\ln S}$$

Information:

$E$  = evenness index

$H'$  = species diversity index

$S$  = number of species found

Evenness index values were categorised as follows:  $0 < E \leq 0.5$ : "Community is depressed",  $0.5 < E \leq 0.75$ : "Community is unstable", and  $0.75 < E \leq 1$ : "Community is stable".

### Results and Discussion

Chlorophyta microalgae found in the Mesat River consisted of 3 classes, 6 orders, 11 families, 15 genera and 19 species as shown in Table 1.

The species of microalgae Chlorophyta are *Draparnaldia* sp., *Stigeoclonium lubricum*, *Chlamydomonas* sp., *Chlorococcum* sp., *Eudorina* sp., *Volvox* sp., *Pediastrum duplex*, *Pediastrum simplex*, *Tetraedron* sp., *Coelastrum* sp., *Scenedesmus armatus*, *Scenedesmus dimorphus*, *Scenedesmus quadricauda*, *Ankistrodesmus* sp., *Chlorella pyrenoidosa*, *Chlorella vulgaris*, *Oocystis* sp., *Rhizoclonium* sp., and *Ulothrix* sp.

Chlorophyceae has the highest number of species, namely 14 species (74%), then, Trebouxiophyceae class 3 (16%) and Ulvophyceae class 2 species (10%). The percentage composition of microalgae species can be seen in Figure 2.

The composition of microalgae species based on the sampling location, namely Station 1 has the highest composition with 45%, then, Station 2 with a value of 33% and finally, Station 3 with a value of 22%. The composition of microalgae by station can be seen in Figure 3.

The results of the analysis of the diversity, dominance, and evenness of Chlorophyta microalgae in the Mesat River are shown in Table 2.

Chlorophyta diversity in all stations was in the low category with the highest diversity index found at Station 1 of 1.27 while at Station

Table 1: Recapitulation of microalgae types in Mesat River Lubuklinggau City, South Sumatera Province, Indonesia

No.	Class	Order	Family	Genera	Species	
1	Chlorophyceae	Chaetophorales	Chaetophoraceae	Draparnaldia	<i>Draparnaldia</i> sp.	
2				Stigeoclonium	<i>Stigeoclonium lubricum</i>	
3		Chlamydomonadales	Chlamydomonadaceae	Chlamydomonas	<i>Chlamydomonas</i> sp.	
4			Chlorococcaceae	Chlorococcum	<i>Chlorococcum</i> sp.	
5			Volvocaceae	Eudorina	<i>Eudorina</i> sp.	
6				Volvox	<i>Volvox</i> sp.	
7				Selenastraceae	Ankistrodesmus	<i>Ankistrodesmus</i> sp.
8			Hydrodictyaceae	Pediasstrum		<i>Pediasstrum duplex</i>
9						<i>Pediasstrum Simplex</i>
10		Tetraedron			<i>Tetraedron</i> sp.	
11		Sphaeropleales	Coelastrum		<i>Coelastrum</i> sp.	
12				Scenedesmaceae		<i>Scenedesmus armatus</i>
13					Scenedesmus	<i>Scenedesmus dimorphus</i>
14						<i>Scenedesmus quadricauda</i>
15		Trebouxiophyceae	Chlorellales	Chlorellaceae	Chlorella	<i>Chlorella pyrenoidosa</i>
16						<i>Chlorella vulgaris</i>
17					Oocystaceae	Oocyst
18		Ulvophyceae	Cladophorales	Cladophoraceae	Rhizoclonium	<i>Rhizoclonium</i> sp.
19			Ulotrichales	Ulotrichaceae	Ulothrix	<i>Ulothrix</i> sp.

2 of 0.98, it was the lowest. The dominance index value at all stations was also in the low category which indicates that if the dominance index value (C) is close to zero, then, there is no dominant biota in the water. The highest evenness index value at Station 1 was 0.42 and the lowest at Station 2 of 0.32 was included in the “depressed community” category. The smaller the uniformity index, the smaller the uniformity of the population, which proves that

the distribution of the number of individuals of each species is not the same, so, there is a tendency for one type of population to dominate. The higher the homogeneity value, the more descriptive the number of populations of the organisms in each species is either the same or not significantly different.

Measurements of temperature, dissolved oxygen, acidity, and clarity in the Mesat River are shown in Table 3.




Figure 2: Percentage composition of microalgae species in Mesat River Lubuklinggau City, South Sumatera Province, Indonesia




Figure 3: Percentage of microalgae composition by station in Mesat River Lubuklinggau City, South Sumatera Province, Indonesia

Table 2: Diversity, dominance and evenness of microalgae species in Mesat River Lubuklinggau City, South Sumatera Province, Indonesia

No.	Component	Station 1	Station 2	Station 3	Mean	Std. Deviation
1	Diversity	1.27	0.98	1.04	2.35	0.22
2	Dominance	0.01	0.008	0.006	0.11	0.01
3	Evenness	0.42	0.32	0.34	0.79	0.07

Table 3: Abiotic factors in Mesat River Lubuklinggau City, South Sumatera Province, Indonesia

No.	Component	Station 1	Station 2	Station 3	Mean	Std. Deviation
1	Temperature (°C)	27.2	26.5	26.8	26.83	0.35
2	Dissolved oxygen (mg/L)	4.6	4.2	4.5	4.43	0.20
3	Acidity	6.7	6.6	6.6	6.63	0.05
4	Clarity (cm)	16.3	22	17.5	18.60	3.00

Dissolved oxygen levels and pH still meet water quality standards in accordance with Government Regulation No. 82 of 2001. Dissolved oxygen belongs to class II, namely

with a value of 4 mg/L, acidity or pH is included in all classes, both classes I–IV with values between 6 and 9. The clarity as measured by the Secchi disk and the water temperature as



measured by the thermometer in the regulations are not clearly written as to what the standard numbers should be.

The number of species of Chlorophyta microalgae found in the Mesat River was 19 species. Based on Table 1, the highest diversity index is at Station 1 with a value of 1.27 and the lowest is at Station 2 with a value of 0.98. The diversity index, which ranged from 0.98–1.27 is considered low because according to Odum (1971), the criteria for diversity is as follows:  $H' \leq 2$ : “Small diversity”,  $2 < H' \leq 3$ : “Medium diversity”, and  $H' > 3$ : “High diversity”.

Species diversity in a waters can provide information about the level of pollution of the water (Chen *et al.*, 2019; Nabi *et al.*, 2019). The diversity index value ( $H'$ ) ranged from 0–3. The level of diversity will be high if the value of  $H'$  is close to 3, which indicates good water conditions. Conversely, if the value of  $H'$  is close to 0, then, the diversity is low and the water conditions are not good (Odum, 1971).

The highest dominance index is Station 1 with a value of 0.019 and the lowest was at Station 3 with a value of 0.006. The dominance index is used to see the most common species found in a habitat and describes the presence of the dominant species (Li *et al.*, 2021; Lin *et al.*, 2021). The dominance of Chlorophyta microalgae is included in the criteria for low dominance  $0 < C < 0.5$  (Krebs, 2001). Based on this, in general in the Mesat River during the study there was no dominant species in the Chlorophyta microalgae community. However, there are species that are often found, namely from the Genera Chlorella.

The highest evenness index value (0.42) is at Station 1 which is included in the medium criteria and the lowest (0.32) is at Station 2 which is included in the low criteria. The level of evenness of species is a description of the distribution of individuals in the community (Engel *et al.*, 2020). Most Chlorophyta are autotrophs meaning they can synthesize food directly from inorganic compounds (Calbet *et al.*, 2014; Khalili *et al.*, 2020). If environmental conditions are favourable, Chlorophyta will

grow rapidly which can fill water bodies and cause an algae population explosion (Odufuwa & Ajaba, 2019; Chaffai *et al.*, 2022).

The average temperature in the Mesat River is 26.9°C, dissolved oxygen content is 4 mg/L, water pH is 6.7 and clarity is 18.63 cm (Table 3). The presence of Chlorophyta in waters will vary depending on water quality conditions which are grouped into physical and chemical factors (Gil-Guarín *et al.*, 2020; Pala *et al.*, 2021). Temperature is one of the environmental factors that greatly affects the life of aquatic organisms, including microalgae (Harmoko & Krisnawati, 2018). The Mesat River temperature range is ideal for microalgae growth, the optimum temperature limit for microalgae growth as found in several previous studies, is around 20–30°C (Singh & Singh, 2015; Corredor *et al.*, 2021; Soriano *et al.*, 2021).

The highest dissolved oxygen level in the Mesat River is at Station 1 was 4.59 mg/L and the lowest is at Station 2 with a value of 4 mg/L. The average dissolved oxygen level in the Mesat River was 4.46 mg/L when adjusted for the dissolved oxygen criteria by Young *et al.* (2014), indicating the criteria for not being considered polluted. This can be seen from the condition of the Mesat River which is still clean but during the rainy season, there is a lot of garbages and it gets cloudy. Based on PP No. 82 of 2001 for the DO and pH parameters of water in the Mesat River, it is classified as class II. Water in class II criteria is water that can be used for water recreation facilities/infrastructure, freshwater fish farming, agriculture, animal husbandry, and other uses requiring water quality equivalent to those specified.

The pH value in the Mesat River ranged from 6.6 to 6.7. Organisms have different tolerance limits to acidity, most natural waters have acidity ranging from 6–9 (Ying *et al.*, 2014; Furuhashi *et al.*, 2019). Based on this, the Mesat River has a pH that is within the normal range, so that microalgae can grow well.

Clarity with a value of 18.63 cm is included in the cloudy category. This is in accordance with Arthington *et al.* (2006), namely cloudy

waters if the clarity value is between 0.25 and 1 m. If the water clarity is less than 25 cm, it can endanger the life of plankton because photosynthesis is inhibited (Buchanan, 2020) taxonomic composition, biomass. High turbidity inhibits the penetration of sunlight in the photosynthesis process of phytoplankton and can cause siltation. Penetration of light into the water is influenced by the intensity and angle of incidence of light, water surface conditions, and dissolved and suspended materials in the water (Atta et al., 2013).

Chlorophyta microalgae found in the Mesat River in Lubuklinggau City found 19 species and included many categories when compared to previous studies. In the Kelingi River, Lubuklinggau City, 15 species of chlorophytic microalgae were found (Harmoko & Sepriyaningsih, 2018) while in the Kasie River, 9 types of Chlorophyta were found (Harmoko & Sepriyaningsih, 2020). This difference is caused by different river conditions, the Mesat River is a river that is polluted with organic waste, causing the presence of quite lots of Chlorophyta.

### Conclusion

Microalgae Chlorophyta found in the Mesat River includes 3 classes, 6 orders, 11 families, 15 genera and 19 species. The Mesat River has a diversity of Chlorophyta microalgae in a small category with a low level of dominance and the Genera that is often found in the Mesat River is *Chlorella*. This research can be used as data for related services, management and protecting the environment in the Mesat River in Lubuklinggau City, considering that Chlorophyta is one the reasons that algae blooms in the water.

### Acknowledgements

The author would like to thank the of the Republic of Indonesia, Research and Technology Ministry, Research and Community Service Directorate that funded this research paper through a Beginner Lecturer Research scheme in 2020 under Contract Number: 815/SP2H/LT/MONO/LL2/2020.

### References

- Arthington, A. H., Bunn, S. E., Poff, N. L. R., & Naiman, R. J. (2006). The challenge of providing environmental flow rules to sustain river ecosystems. *Ecological Applications*, 16(4), 1311-1318. <https://doi.org/10.1890/1051-0761>
- Atta, M., Idris, A., Bukhari, A., & Wahidin, S. (2013). Intensity of blue LED light: A potential stimulus for biomass and lipid content in fresh water microalgae *Chlorella vulgaris*. *Bioresource Technology*, 148(3), 373-378. <https://doi.org/10.1016/j.biortech.2013.08.162>
- Bat, L., Öztekin, A., Şahin, F., Arıcı, E., & Özsandıkçı, U. (2018). An overview of the Black Sea pollution in Turkey. *Mediterranean Fisheries and Aquaculture Research*, 1(2), 67-86. <https://doi.org/medfar.v1i37150.397536>
- Bellinger, E. G., & Sigeo, D. C. (2010). *Freshwater algae identification and use as bioindicators*. Wiley-Blackwell.
- Buchanan, C. (2020). A water quality binning method to infer Phytoplankton community structure and function. *Estuaries and Coasts*, 43(4), 661-679. <https://doi.org/10.1007/s12237-020-00714-3>
- Calbet, A., Sazhin, A. F., Nejstgaard, J. C., Berger, S. A., Tait, Z. S., Olmos, L., Sousoni, D., Isari, S., Martínez, R. A., Bouquet, J. M., Thompson, E. M., Båmstedt, U., & Jakobsen, H. H. (2014). Future climate scenarios for a coastal productive planktonic food web resulting in microplankton phenology changes and decreased trophic transfer efficiency. *PLOS ONE*, 9(4), 1-16. <https://doi.org/10.1371/journal.pone.0094388>
- Celewicz-Goldyn, S., & Kuczynska-Kippen, N. (2017). Ecological value of macrophyte cover in creating habitat for microalgae (diatoms) and zooplankton (rotifers and crustaceans) in small field and forest water bodies. *PLOS ONE*, 12(5), 1-14. <https://doi.org/10.1371/journal.pone.0177317>

- Chen, J., McIlroy, S. E., Archana, A., Baker, D. M., & Panagiotou, G. (2019). A pollution gradient contributes to the taxonomic, functional, and resistome diversity of microbial communities in marine sediments. *Microbiome*, 7(1), 1-12. <https://doi.org/10.1186/s40168-019-0714-6>
- Corredor, L., Barnhart, E. P., Parker, A. E., Gerlach, R., & Fields, M. W. (2021). Effect of temperature, nitrate concentration, pH and bicarbonate addition on biomass and lipid accumulation in the sporulating green alga PW95. *Algal Research*, 53, 102148. <https://doi.org/10.1016/j.algal.2020.102148>
- de Freitas Coêlho, D., Tundisi, L. L., Cerqueira, K. S., da Silva Rodrigues, J. R., Mazzola, P. G., Tambourgi, E. B., & de Souza, R. R. (2019). Microalgae: Cultivation aspects and bioactive compounds. *Brazilian Archives of Biology and Technology*, 62(2), 1-13. <https://doi.org/10.1590/1678-4324-2019180343>
- Engel, F. G., Dini-Andreote, F., Eriksson, B. K., Salles, J. F., de Lima Brossi, M. J., & Matthiessen, B. (2020). Dispersal mitigates bacterial dominance over microalgal competitor in metacommunities. *Oecologia*, 193(3), 677-687. <https://doi.org/10.1007/s00442-020-04707-8>
- Odum, P. E. (1971). *Fundamentals of Ecology*. W.B. Saunders Company.
- Furuhashi, Y., Hond, R., Noguchi, M., Hara-Yamamura, H., Kobayashi, S., Higashimine, K., & Hasegawa, H. (2019). Optimum conditions of pH, temperature and preculture for biosorption of europium by microalgae *Acutodesmus acuminatus*. *Biochemical Engineering Journal*, 143(3), 58-64. <https://doi.org/10.1016/j.bej.2018.12.007>
- Geraldine Soriano, H. A., Baldia B, B., & Fernando, S. (2021). Preliminary culture studies of microalgae isolated from the freshwater tributaries of Carigara bay, Leyte, Philippines. *Asian Journal of Biological and Life Sciences*, 9(3), 408-415. <https://doi.org/10.5530/ajbls.2020.9.62>
- Gil-Guarín, I. C., Villabona-González, S. L., Parra-García, E., & Echenique, R. O. (2020). Environmental factors driving phytoplankton biomass and diversity in a tropical reservoir. *Revista de La Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 44(171), 423-436. <https://doi.org/10.18257/racefyn.1052>
- Harmoko, H., & Sepriyaningsih, S. (2018). Keanekaragaman mikroalga chlorophyta di Sungai Kelingi Kota Lubuklinggau, Sumatera Selatan. *Jurnal Pro Life*, 5(3), 666-676. <https://doi.org/10.33541/jpvol6Iss2pp102>
- Harmoko, H., & Sepriyaningsih, S. (2020). Keanekaragaman mikroalga chlorophyta di Sungai Kasie Kota Lubuklinggau Provinsi Sumatera Selatan. *Quagga: Jurnal Pendidikan dan Biologi*, 12(1), 52. <https://doi.org/10.25134/quagga.v12i1.2142>
- Harmoko, H., Triyanti, M., & Aziz, L. (2018). Eksplorasi mikroalga di Sungai Mesat Kota Lubuklinggau. *Biodidaktika, Jurnal Biologi dan Pembelajarannya*, 13(2), 19-23. <https://doi.org/10.30870/biodidaktika.v13i2.3366> [in Indonesia].
- Harmoko, & Krisnawati, Y. (2018). Keanekaragaman mikroalga divisi cyanobacteria di danau aur kabupaten musi rawas. *Jurnal Biodjati*, 3(1), 8-14. <https://doi.org/10.15575/biodjati.v3i1.1638> [in Indonesia].
- Kaparapu, J. (2018). Application of microalgae in aquaculture. *Phykos*, 48(1), 21-26.
- Khalili, Z., Jalili, H., Noroozi, M., Amrane, A., & Ashtiani, F. R. (2020). Linoleic-acid-enhanced astaxanthin content of *Chlorella sorokiniana* (Chlorophyta) under normal and light shock conditions. *Phycologia*, 59(1), 54-62. <https://doi.org/10.1080/00318884.2019.1670012>
- Krebs, C. J. (2001). *Ecology: The experimental analysis of distribution and abundance* (5th Edition), Benjamin Cummings.



- Li, Y., Yu, Z., Ji, S., Meng, J., Kong, Q., Wang, R., & Liu, J. (2021). Diverse drivers of phytoplankton dynamics in different phyla across the annual cycle in a freshwater lake. *Journal of Freshwater Ecology*, 36(1), 13-29. <https://doi.org/10.1080/02705060.2020.1868586>
- Lin, L., Wang, F., Chen, H., Fang, H., Zhang, T., & Cao, W. (2021). Ecological health assessments of rivers with multiple dams based on the biological integrity of phytoplankton: A case study of North Creek of Jiulong River. *Ecological Indicators*, 121, 106998. <https://doi.org/10.1016/j.ecoind.2020.106998>
- Lomartire, S., Marques, J. C., & Gonçalves, A. M. M. (2021). The key role of zooplankton in ecosystem services: A perspective of interaction between zooplankton and fish recruitment. *Ecological Indicators*, 129(1), 147-152. <https://doi.org/10.1016/j.ecoind.2021.107867>
- M. Rasal, V., G. Yadre, S., Shukla, S. P., Ravi, P. M., Ravi, P. M., Kumar Mishra, M., Munilkumar, S., Kumar Pal, A., Lakra, W. S., & Dasgupta, S. (2019). Microalgae distribution and diversity in the Narmada River Basin around Chutka, Madhya Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*, 8(09), 1488-1501. <https://doi.org/10.20546/ijcmas.2019.809.171>
- Mourelle, M. L., Gómez, C. P., & Legido, J. L. (2017). The potential use of marine microalgae and cyanobacteria in cosmetics and thalassotherapy. *Cosmetics*, 4(4). <https://doi.org/10.3390/cosmetics4040046>
- Nabi, G., Ali, M., Khan, S., & Kumar, S. (2019). The crisis of water shortage and pollution in Pakistan: risk to public health, biodiversity, and ecosystem. *Environmental Science and Pollution Research*, 26(11), 10443-10445. <https://doi.org/10.1007/s11356-019-04483-w>
- Pala, G., Caglar, M., FARUQ, R., & Selamoglu, Z. (2021). Research article: Chlorophyta algae of Keban Dam Lake Gülüşkür region with aquaculture criteria in Elazığ, Turkey. *Iranian Journal of Aquatic Animal Health*, 7(1), 32-46. <https://doi.org/10.52547/ijaah.7.1.32>
- Presiden Republik Indonesia. (n.d.). *Peraturan pemerintah Republik Indonesia Nomor 82 Tahun 2001, Peraturan pemerintah Republik Indonesia I (2001)*. Regulasip.
- Prata, J. C., Costa, J. P. da, Lopes, I., Duarte, A. C., & Rocha-Santos, T. (2019). Effects of microplastics on microalgae populations: A critical review. *Science of the Total Environment*, 665(1), 400-405. <https://doi.org/10.1016/j.scitotenv.2019.02.132>
- Singh, S. P., & Singh, P. (2015). Effect of temperature and light on the growth of algae species: A review. *Renewable and Sustainable Energy Reviews*, 50, 431-444. <https://doi.org/10.1016/j.rser.2015.05.024>
- Thyagaraju, N. (2016). Water pollution and its impact on environment of society. *International Research Journal of Management, IT & Social Sciences*, 3(5), 1-7. <https://doi.org/10.21744/irjmis.v3i5.10>
- Wehr, J. D., & Sheath, R. G. (2003). *Freshwater algae of north america ecology and classification*. Academic Press.
- Ying, K., Gilmour, D. J., & Zimmerman, W. B. (2014). Effects of CO<sub>2</sub> and pH on growth of the microalga *Dunaliella salina*. *Journal of Microbial and Biochemical Technology*, 6(3), 167-173. <https://doi.org/10.4172/1948-5948.1000138>
- Young, S. Sen, Yang, H. N., Huang, D. J., Liu, S. M., Huang, Y. H., Chiang, C. T., & Liu, J. W. (2014). Using benthic macroinvertebrate and fish communities as bioindicators of the Tanshui river basin around the greater Taipei area - Multivariate analysis of spatial variation related to levels of water pollution. *International Journal of Environmental Research and Public Health*, 11(7), 7116-7143. <https://doi.org/10.3390/ijerph110707116>